











FROM
THE LIBRARY
OF
SIR WILLIAM OSLER, BART.
OXFORD




















THE
PHYSICIAN'S
VISITING LIST
LINDSAY & BLAKISTON'S
FOR
1885.

THIRTY FOURTH YEAR OF ITS PUBLICATION.

PHILADELPHIA:
P. BLAKISTON, SON & CO.

(SUCCESSORS TO LINDSAY & BLAKISTON),

1012 WALNUT STREET.

SOLD BY ALL BOOKSELLERS AND DRUGGISTS.

CONTENTS.

	PAGE
ALMANAC,	3
TABLE OF SIGNS,	4
MARSHALL HALL'S READY METHOD IN ASPHYXIA,	5
POISONS AND ANTIDOTES,	6
THE METRIC OR FRENCH DECIMAL SYSTEM OF WEIGHTS AND MEASURES,	8
POSOLOGICAL TABLE, rewritten in accordance with the new (sixth) Revision of the U. S. Pharmacopœia,	10
TABLE FOR CALCULATING THE PERIOD OF UTERO- GESTATION,	14
LIST OF NEW REMEDIES,	25
SYLVESTER'S METHOD FOR ARTIFICIAL RESPIRATION,	26
DIAGRAM OF THE CHEST.	27

BLANK LEAVES, for Visiting List.

"	"	"	Monthly Memoranda.
"	"	"	Addresses of Patients and others.
"	"	"	" " Nurses, their references, etc.
"	"	"	Accounts asked for.
"	"	"	Memoranda of Wants.
"	"	"	Obstetric Engagements.
"	"	"	Vaccination Engagements.
"	"	"	Record of Births.
"	"	"	" " Deaths.
"	"	"	General Memoranda, etc.

Entered according to the Act of Congress, in the year 1884,

BY P. BLAKISTON, SON & CO.,

In the Office of the Librarian of Congress at Washington.

PRESS OF HENRY B. ASHMEAD

ALMANAC FOR 1885.

	S	M	T	W	T	F	S		S	M	T	W	T	F	S		
January .	...	...	...	...	...	1	2	3	July ...	...	...	...	...	1	2	3	4
	4	5	6	7	8	9	10			5	6	7	8	9	10	11	
	11	12	13	14	15	16	17			12	13	14	15	16	17	18	
	18	19	20	21	22	23	24			19	20	21	22	23	24	25	
	25	26	27	28	29	30	31			26	27	28	29	30	31	..	
February.	1	2	3	4	5	6	7		August ..	...	...	...	...	...	1		
	8	9	10	11	12	13	14			2	3	4	5	6	7	8	
	15	16	17	18	19	20	21			9	10	11	12	13	14	15	
	22	23	24	25	26	27	28			16	17	18	19	20	21	22	
										23	24	25	26	27	28	29	
March...	1	2	3	4	5	6	7		September	...	...	...	...	...	...	...	
	8	9	10	11	12	13	14			...	...	...	...	...	...	...	
	15	16	17	18	19	20	21			...	...	...	...	...	...	...	
	22	23	24	25	26	27	28			...	...	...	...	...	...	...	
	29	30	31	...	...	...	...			...	...	...	...	...	...	...	
April....	...	...	...	1	2	3	4		October ..	...	...	...	...	...	1	2	3
	5	6	7	8	9	10	11			...	...	...	...	...	4	5	6
	12	13	14	15	16	17	18			...	...	...	...	...	7	8	9
	19	20	21	22	23	24	25			...	...	...	...	...	10	11	12
	26	27	28	29	30	...	...			...	...	...	...	...	13	14	15
May	...	...	...	...	1	2			November	...	...	...	...	...	16	17	18
	3	4	5	6	7	8	9			...	...	...	...	...	19	20	21
	10	11	12	13	14	15	16			...	...	...	...	...	22	23	24
	17	18	19	20	21	22	23			...	...	...	...	...	25	26	27
	24	25	26	27	28	29	30			...	...	...	...	...	28	29	30
	31	...	...	...	...	...	...			...	...	...	...	...	...	...	...
June	...	1	2	3	4	5	6		December	...	...	...	...	...	1	2	3
	7	8	9	10	11	12	13			...	...	...	...	...	4	5	6
	14	15	16	17	18	19	20			...	...	...	...	...	7	8	9
	21	22	23	24	25	26	27			...	...	...	...	...	10	11	12
	28	29	30	...	...	...	...			...	...	...	...	...	13	14	15
										...	...	...	...	...	16	17	18
										...	...	...	...	...	19	20	21
										...	...	...	...	...	22	23	24
										...	...	...	...	...	25	26	27
										...	...	...	...	...	28	29	30
										...	...	...	...	...	31	...	...

1886—January to July.

	S	M	T	W	T	F	S		S	M	T	W	T	F	S
January .	...	...	...	...	...	1	2	April	...	...	...	...	1	2	3
	3	4	5	6	7	8	9		4	5	6	7	8	9	10
	10	11	12	13	14	15	16		11	12	13	14	15	16	17
	17	18	19	20	21	22	23		18	19	20	21	22	23	24
	24	25	26	27	28	29	30		25	26	27	28	29	30	...
February.	31	...	...	...	...	...	...	May	...	...	...	...	...	...	1
	1	2	3	4	5	6	7		2	3	4	5	6	7	8
	8	9	10	11	12	13	14		9	10	11	12	13	14	15
	15	16	17	18	19	20	21		16	17	18	19	20	21	22
	22	23	24	25	26	27	28		23	24	25	26	27	28	29
	29	30	...	...	...	...	...		30	31	...	...	...	...	...
March . . .	...	1	2	3	4	5	6	June	...	...	1	2	3	4	5
	7	8	9	10	11	12	13		6	7	8	9	10	11	12
	14	15	16	17	18	19	20		13	14	15	16	17	18	19
	21	22	23	24	25	26	27		20	21	22	23	24	25	26
	28	29	30	31	...	...	...		27	28	29	30	...	...	...

TABLE OF SIGNS.

THE following Table explains itself. Thus each patient having a line, and each day its column. The marks in the margin, as below, are intended to be made in the column under the proper date, and immediately on a line with the patient's name. For instance, for a visit to be made, a dot thus •; when made, the dot will be crossed /; for a second visit, •/; when made, the dot is again X, and so on.

•	Visit to be made.
/	Visit made.
•/	Second visit to be made.
X	Second visit made.
10	Visit to be made at a certain hour A. M.
5	Visit to be made at a certain hour P. M.
C	Consultation.
✓	Consultation held.
L	Visit made, and medicine sent.
N	Night Visit.
V	Vaccination.
O	Obstetric case.
œ	Office Consultation.

MARSHALL HALL'S

READY METHOD IN ASPHYXIA.

1st. TREAT the patient *instantly on the spot*, in the *open air*, freely exposing the face, neck, and chest to the breeze, except in severe weather.

2d. In order to *clear the throat*, place the patient gently on the face, with one wrist under the forehead, that all fluid, and the tongue itself, may fall forward, and leave the entrance into the windpipe free.

3d. To *excite respiration*, turn the patient slightly on his side, and apply some irritating or stimulating agent to the nostrils, as *Veratrine*, *dilute Ammonia*, etc.

4th. Make the face warm by brisk friction; then dash cold water upon it.

5th. If not successful, lose no time; but, to *imitate respiration*, place the patient on his face, and turn the body gently, but completely, *on the side, and a little beyond*; then again on the face, and so on, alternately. Repeat these movements deliberately and perseveringly, *fifteen times only* in a minute. (When the patient lies on the thorax, this cavity is *compressed* by the weight of the body, and *expiration* takes place. When he is turned on the side, this pressure is removed, and *inspiration* occurs.)

6th. When the prone position is resumed, make a uniform and efficient pressure *along the spine*, removing the pressure immediately, before rotation on the side. (The pressure augments the *expiration*; the rotation commences *inspiration*.) Continue these measures.

7th. Rub the limbs *upward*, with *firm pressure* and with *energy*. The object being to aid the return of venous blood to the heart.

8th. Substitute for the patient's wet clothing, if possible, such other covering as can be instantly procured, each bystander supplying a coat or cloak, etc. Meantime, and from time to time, to *excite inspiration*, let the surface of the body be *slapped* briskly with the hand.

9th. Rub the body briskly till it is dry and warm, then dash cold water upon it, and repeat the rubbing.

AVOID the immediate removal of the patient, as it involves a *dangerous loss of time*; also, the use of bellows, or any *forcing instrument*; also, the *warm bath*, and all *rough treatment*.

POISONS AND ANTIDOTES.

POISON.	ANTIDOTE.
ACIDS, MINERAL.....	(Chalk, magnesia (plaster off wall in emergency), solution carbonate of soda, emollient drinks, fixed oils.
ACONITE.....	(Emetics, stimulants external and internal.
ANTIM. TART.....	(Vegetable acids, such as tannic acid, catechu.
ARSENIC.....	(Moist peroxide of iron (obtained from perchloride of iron and calcined magnesia), charcoal, ammonia, artificial respiration, cold affusion.
ATROPIA.....	Same as Belladonna.
ARGENTI NIT.....	(Solution of common salt in demulcent drinks.
BELLADONNA.....	(Emetics—chiefly sulphate of copper (gr. x.), cold to head, active cathartics, ammonia externally and internally, stimulants, opium.
CANNIBIS INDICA.....	(Hot brandy and water, vegetable acids, lemon juice, vinegar, etc.; allow patient to sleep; blister to nape of neck.
CANTHARIDES.....	(Emetics, emollient drinks, opiates by mouth and rectum.
CARBOLIC ACID.....	Castor oil, olive oil.
CHLORINE WATER.....	Albumen, white of egg, milk, flour.
CHLOROFORM.....	(Fresh air, artificial respiration (inclining head down, pull tongue forward), galvanism to pneumogastric and through diaphragm, brandy and ammonia enemata. A case in which Tracheotomy was successful; performed by Mr. Howse, Guy's Hospital. The hypodermic injection of 1-10 gr. digitaline, followed in four hours after by 1-10 gr. atropin. by Professor Larabee, Louisville, recovered a patient after galvanism had failed.
COLCHICUM.....	(Emetics, followed by demulcent drinks. If coma be present, brandy, ammonia, coffee.
CONIUM.....	(Emetics; followed by stimulants external and internal.
CORROSIVE SUBLIMATE.....	(Albumen, white of egg (4 gr. sublim. require white of one egg), flour, milk, protochloride of tin.

POISONS AND ANTIDOTES.—Continued.

POISON.	ANTIDOTE.
CROTON OIL.....	{ Emetic of sulphate of copper 10 gr., followed by mucilaginous fluid, containing opium.
CUPRI SULPH.....	Albumen or white of egg.
DIGITALIS.....	{ Recumbent posture after emetics, stimulants internally and exter- nally.
ELATERIUM.....	{ Demulcent drinks and enemata, small doses of opium, and the warm bath.
HYDROCYANIC ACID.....	{ Fresh air and artificial respiration, with cold affusion; freshly precipi- tated oxide of iron, with an alkaline carbonate.
HYDROCYANIC.....	{ Stomach pump, emetics, stimulants external and internal, lemon juice.
IODINE.....	{ Emetics and demulcent drinks, starch or flour diffused in water.
LEAD SALTS.....	{ Sulphate or phosphate of soda, epsom salts, succeeded by emetics, and afterwards opium and milk.
LOBELIA.....	Stimulants externally and internally.
MORPHIA.....	Same as Opium.
NUX VOMICA.....	{ Enema of tobacco, $\frac{1}{2}$ oz. to 20 oz. of boiling water to be given till spasms abate; Nicotine $\frac{1}{4}$ drop in warm sherry and water.
OPIMUM.....	{ Emetic of 1 to 2 sulphate of copper soda, stomach pump, stimulants external and internal, cold and coffee, cold affusion, ammonia to nostrils, galvanic shocks compelling to move, artificial respiration, and camellina.
PHOSPHORUS.....	Magnesia, sulphuric, citric, and tartaric acids.
POTASH AND SODA SALTS.....	Dilute acetic acid (tri acid), lemon juice fixed as demulcents.
STRAMONIUM.....	Same as Belladonna.
STRYCHNIA.....	{ Camellina, belladonna, tincture of opium, morphia, tobacco, liberal use of hot doses, 5 gr. morphia increased to 1 gr. strychnia.
TOBACCO.....	{ Emetic, stimulants external and in- ternal, strychnia.
ZINC SALTS.....	Carbonate of soda, emetics, warm demulcent drinks.

METRIC OR FRENCH DECIMAL SYSTEM OF WEIGHTS AND MEASURES.

BY

OSCAR OLDBERG, PHARM. D.,
SAINT LOUIS.

The metric system is based upon the **METER**, which is the standard unit of length of that system, and equal to 39.370432 inches, or about 10 per cent. longer than the yard.

The metric unit of *fluid measure* is the **LITER**—the cube of $\frac{1}{10}$ Meter, or 1000 Cubic-centimeters—equal to about 34 fluid ounces.

The metric unit of *weight* is the **GRAM**, which represents the weight of one Cubic-centimeter of water at its maximum density. It is equal to about 15 grains.

One **CUBIC-CENTIMETER** is equal to about 16 minims.

IN WRITING PRESCRIPTIONS IT IS SUFFICIENTLY ACCURATE AND SAFE TO CONSIDER 1 GRAM AS EXACTLY EQUAL TO 15 TROY GRAINS, AND TO CONSIDER 1 CUBIC-CENTIMETER AS EQUAL TO 15 MINIMS.

We accordingly have:—

1 Gram equal to $\frac{1}{15}$ troy grains.

1 troy grain equal to $\frac{1}{15}$ Gram.

1 Cubic-centimeter equal to $\frac{1}{4}$ fluid drachm.

1 fluid drachm equal to $\frac{1}{4}$ Cubic-centimeter.

Hence:

1. TO CONVERT TROY GRAINS INTO GRAMS, OR MINIMS INTO CUBIC-CENTIMETERS:—

a. *Divide by 10, and from the quotient subtract one-third; or, b. Divide by 15; and*

2. TO CONVERT APOTHECARIES' DRACHMS INTO GRAMS, OR FLUID DRACHMS INTO CUBIC-CENTIMETERS, *multiply by 4.*

In writing prescriptions the "Gram" (abbreviated "Gm.") and "Cubic-centimeter," (abbreviated "C. c.," which may be called "fluigram," and written "i Gm") only, should be used.

All other terms, and units, and prefixes, used in the metric system, may be wholly ignored by the physician and the pharmacist.*

Example of a Metric Prescription:

R. Hydrarg. Chloridi. Corro.....0.25 Gm.
Potassii Iodidi.....10.00 Gm.
Aque.....100.00 C. c.
Tinct. Cinch. Comp.100.00 C. c.
Mix.

The use of a decimal line prevents possible errors.

To write a prescription for fifteen doses of any medicine, write it first for one dose in *grains* and *minims*, and then substitute the same number of "Grams" and "fluigrams," thus:

R. Opii gr.....gr. i
Camphore.....gr. i.
Make one pill,

* The prefixes are simply numerals, as follows:

myria, which means 10,000.	deci, which means 0.1.
kilo, " " 1,000.	centi, " " 0.01.
hecto, " " 100.	milli, " " 0.001.
deka, " " 10.	

and are quite unnecessary in the writing of prescriptions (if not in all cases), English numerals being more convenient, and at least equally explicit.

and to get fifteen such doses in metric terms, write:—

R. Opil.....1 Gm.

Camphore.....2 Gm.

Make fifteen pills.

The Gram and the Cubic-centimeter, (*Augram*) when referring to liquids, may be considered as equal quantities, except the liquids be very heavy (as in the case of chloroform) or very light (as in the case of ether).

Measures may be discarded and weights exclusively employed, if preferred. All quantities in a prescription would then be expressed in GRAMS.*

The average "drop" (water) may be considered equal to 0.05 C. c., or 0.05 Gm. An average TEASPOON holds 5 C. c., and an average TABLESPOON 20 C. c. Decimal numbers should be used as far as practicable without sacrifice of accuracy as to strength and dose of the preparation. It is safe to prescribe 30 Gm. for one troy ounce, and 250 C. c. for eight fluid ounces.

The above contains ALL THAT IT IS NECESSARY TO KNOW OR LEARN OF the metric system in ORDER TO WRITE METRIC PRESCRIPTIONS, without a metric pharmacological table, or with one.

To become familiar with the system, the rules given above for the conversion of Apothecaries' weights and measures into the corresponding metric quantities, may be profitably used, the results to be verified by comparison with the following

TABLE OF EQUIVALENTS.

APOTHECARIES' WEIGHTS (AND MEASURES).	METRIC WEIGHTS (AND MEASURES).
Troy grains (or minims).	Grams (or Cubic-centimeters).
$\frac{1}{64}$	0.001
$\frac{1}{32}$	0.002
$\frac{1}{16}$	0.004
$\frac{1}{8}$	0.008
$\frac{1}{4}$	0.016
$\frac{1}{2}$	0.033
1	0.066
2	0.133
5	0.333
10	0.666
15	1.000
20	1.333
30	2.000
Drachms (or Fluid drachms).	Grams (or Cubic-centimeters).
1	4
2	8
4	16
6	24
Troy ounces (or Fluid ounces).	
1	32
2	64
4	128
6	192
8	256
12	384
16	512

* As any liquid medicine must necessarily be administered to the patient in *measured*, and not in *weighed* doses, it will of course be more convenient to the physician to continue to make use of fluid measures in writing prescriptions, especially as he is already accustomed to this, and would not then have to bear in mind the specific gravity of *any* liquid ingredient in the prescription. To the pharmacist it makes but little difference, as he will have both weights and measures, and can use one or the other, as may be directed. If the physician discards measures, he must, of necessity, so adjust the proportion in his formula as to produce a mixture of which after all the dose must be a "teaspoonful," or some other convenient measure, and this is as unnecessary as it is difficult.

DOSE TABLE.

IN METRIC AS WELL AS APOTHECARIES' WEIGHTS AND MEASURES.

COMPILED FOR THE PHYSICIAN'S VISITING LIST.

By OSCAR OLDBERG, PHARM. D.,

SAINT LOUIS.

THOROUGHLY REVISED AND CORRECTED, ACCORDING TO THE NEW PHARMACOPEIAL STANDARDS (1883).

Remedies.	Dose in metric weights or measures.	Dose in apothecaries' weights or measures.
Abstract. aconiti.....	3 to 10 centigrams	$\frac{1}{2}$ to $1\frac{1}{2}$ grains.
Abstract. aspidosperme...	0.30 to 1.20 grams	5 to 20 grains.
Abstract. belladonnæ.....	3 to 10 centigrams	$\frac{1}{2}$ to $1\frac{1}{2}$ grains.
Abstract. cannab. ind.....	6 to 20 centigrams	1 to 3 grains.
Abstract. conii.....	6 to 20 centigrams	1 to 3 grains.
Abstract. coto.....	6 to 20 centigrams	1 to 3 grains.
Abstract. digitalis.....	6 to 20 centigrams	1 to 3 grains.
Abstract. gelsemii.....	6 to 20 centigrams	1 to 3 grains.
Abstract. hyoscyami.....	12 to 40 centigrams	2 to 6 grains.
Abstract. ignatiæ.....	6 to 20 centigrams	1 to 3 grains.
Abstract. ipecac.....	0.20 to 2 grams	3 to 30 grains.
Abstract. jalapæ.....	0.40 to 2 grams	6 to 30 grains.
Abstract. nuc. vom.....	6 to 20 centigrams	1 to 3 grains.
Abstract. phytolaccae.....	0.30 to 1 gram	5 to 15 grains.
Abstract. pilocarpi.....	0.40 to 2 grams	6 to 30 grains.
Abstract. podophylli.....	25 to 60 centigrams	1 to 10 grains.
Abstract. sanguin. alterat.	6 to 20 centigrams	1 to 3 grains.
Abstract. sanguin. emet...	0.60 to 1.20 Gm.....	10 to 15 grains.
Abstract. senegæ.....	25 to 60 centigrams	4 to 10 grains.
Abstract. valeriana.....	0.60 to 1.20 Gm.....	10 to 15 grains.
Abstract. veratr. vir.....	6 to 20 centigrams	1 to 3 grains.
Acet. lobelia.....	1 to 4 C. c.....	15 to 60 minims.
Acet. opii.....	0.30 to 1 C. c.....	5 to 16 minims.
Acet. sanguinar.....	1 to 2 C. c.....	15 to 30 minims.
Acet. scillæ.....	0.60 to 2 C. c.....	10 to 30 minims.
Acid. acet. dil.....	4 to 6 C. c.....	60 to 90 minims.
Acid. arsenios.....	1 to 5 milligrams	$\frac{1}{2}$ to $\frac{1}{12}$ grain.
Acid. benzoic.....	0.30 to 1 Gm.....	5 to 15 grains.
Acid. boric.....	30 to 60 centigrams	5 to 10 grains.
Acid. carbolic.....	6 to 20 centigrams	1 to 3 grains.
Acid. gallic.....	0.20 to 1 Gm.....	3 to 15 grains.
Acid. gall. in albuminuria	0.60 to 4 Gm.....	10 to 60 grains.
Acid. hydrobrom. 34%.....	0.60 to 1 Gm.....	10 to 15 grains.
Acid. hydrobrom. dil.....	1 to 2.50 C. c.....	15 to 40 minims.
Acid. hydrochlor.....	20 to 60 centigrams	3 to 10 grains.

DOSE TABLE.—Continued.

Remedies.	Dose in metric weights or measures.	Dose in apothecaries weights or measures.
Acid. hydrochlor. dil.....	0.60 to 2 C. c.....	10 to 30 minims.
Acid. hydrocyan. dil.....	0.10 to 0.30 C. c.....	2 to 6 minims.
Acid. lactic.....	1 to 4 Gm.....	15 to 60 grains.
Acid. nitr.....	20 to 60 centigrams.....	3 to 10 grains.
Acid. nitr. dil.....	0.60 to 2 C. c.....	10 to 30 minims.
Acid. nitro-hydrochlor.....	20 to 60 centigrams.....	3 to 10 grains.
Acid. nitro-hydrochlor. dil.....	0.30 to 1.30 C. c.....	5 to 20 minims.
Acid. phosphoric (50.....)	0.20 to 1 Gm.....	3 to 15 grains.
Acid. phosphoric dil.....	0.60 to 4 C. c.....	10 to 60 minims.
Acid. salicyl.....	0.30 to 1 Gm.....	5 to 15 grains.
Acid. sulphuric.....	0.30 to 0.60 Gm.....	5 to 10 grains.
Acid. sulphuric dil.....	0.30 to 2 C. c.....	5 to 30 minims.
Acid. sulphuric arom.....	0.30 to 2 C. c.....	5 to 30 minims.
Acid. sulphuros.....	2 to 4 C. c.....	30 to 60 minims.
Acid. tannic.....	10 to 60 centigrams.....	2 to 10 grains.
Aconitina (white crystals).....	0.15 to 0.30 milligr.....	$\frac{1}{100}$ to $\frac{1}{200}$ grain.
Alum.....	10 to 30 centigrams.....	2 to 5 grains.
Aluminum.....	5 to 20 centigrams.....	1 to 3 grains.
Alumen.....	0.50 to 1 Gm.....	10 to 15 grains.
Ammonii benzoas.....	0.50 to 1.30 Gm.....	10 to 20 grains.
Ammonii bromid.....	0.30 to 2 Gm.....	5 to 30 grains.
Ammonii carb.....	20 to 60 centigrams.....	3 to 10 grains.
Ammonii chlorid.....	1 to 2 Gm.....	15 to 30 grains.
Ammonii iodid.....	0.20 to 1 Gm.....	3 to 15 grains.
Ammonii phosph.....	30 to 1.30 Gm.....	5 to 20 grains.
Ammonii picras.....	0.50 to 3 centigrams.....	$\frac{1}{4}$ to $\frac{1}{2}$ grain.
Ammonii sulph.....	0.20 to 1 Gm.....	3 to 15 grains.
Ammonii valer.....	0.20 to 1 Gm.....	3 to 15 grains.
Anyl nitris.....	0.10 to 0.40 C. c.....	2 to 5 minims.
Anhydum iodatum.....	0.20 to 2 Gm.....	3 to 30 grains.
Ant. et pot. tartr.; diaph.....	4 to 10 milligrams.....	$\frac{1}{10}$ to $\frac{1}{2}$ grain.
Ant. et pot. tartr.; emetic.....	6 to 10 centigrams.....	1 to 2 grains.
Antimonii oxid.....	3 to 10 centigrams.....	$1\frac{1}{2}$ to 2 grains.
Antimonii oxysulphuret.....	3 to 10 centigrams.....	$\frac{1}{2}$ to 2 grains.
Antimonii sulphid.....	3 to 10 centigrams.....	$\frac{1}{2}$ to 2 grains.
Antimonii sulphuret.....	3 to 10 centigrams.....	$\frac{1}{2}$ to 2 grains.
Apomorph. hydrochlor.....	3 to 6 milligrams.....	$\frac{1}{10}$ to $\frac{1}{20}$ grain.
Aqua ammoniac.....	0.30 to 2 C. c.....	6 to 30 minims.
Aqua amygd. amor.....	10 to 15 C. c.....	2 to 4 fl. drachms.
Aqua camphora.....	15 to 60 C. c.....	$\frac{1}{2}$ to 2 fl. oz.
Aqua chlori.....	4 to 15 C. c.....	1 to 4 fl. drachms.
Aqua creasoti.....	4 to 15 C. c.....	1 to 4 fl. drachms.
Aqua laurocerasi.....	0.30 to 2 C. c.....	6 to 30 minims.
Argenti iodidum.....	3 to 10 centigrams.....	$\frac{1}{2}$ to 2 grains.
Argenti nitras.....	1 to 2 centigrams.....	$\frac{1}{10}$ to $\frac{1}{20}$ grain.
Argenti oxid.....	3 to 10 centigrams.....	$\frac{1}{2}$ to 2 grains.
Arseni iodidum.....	1 to 2.50 milligrams.....	$\frac{1}{100}$ to $\frac{1}{20}$ grain.
Asafetida.....	0.30 to 1.30 Gm.....	5 to 20 grains.
Atropina.....	0.50 to 2 milligrams.....	$\frac{1}{100}$ to $\frac{1}{20}$ grain.
Atropina sulph.....	0.50 to 2 milligrams.....	$\frac{1}{100}$ to $\frac{1}{20}$ grain.
Auri et sodii chlorid.....	2 to 4 milligrams.....	$\frac{1}{100}$ to $\frac{1}{20}$ grain.
Bebeerina sulph.....	20 to 60 centigrams.....	3 to 10 grains.
Berberina and its salts.....	0.20 to 1 Gm.....	1 to 15 grains.
Bismuthi citras.....	0.20 to 1 Gm.....	3 to 15 grains.
Bismuthi et ammon. citr.....	0.10 to 1 Gm.....	1 to 15 grains.

DOSE TABLE.

IN METRIC AS WELL AS APOTHECARIES' WEIGHTS AND MEASURES.

COMPILED FOR THE PHYSICIAN'S VISITING LIST,

By OSCAR OLDBERG, PHARM. D.,

SAINT LOUIS.

THOROUGHLY REVISED AND CORRECTED, ACCORDING TO THE NEW PHARMACOPOEIAL STANDARDS (1883).

Remedies.	Dose in metric weights or measures	Dose in apothecaries' weights or measures
Abstract. aconiti.....	3 to 10 centigrams	$\frac{1}{2}$ to $1\frac{1}{2}$ grains.
Abstract. aspidosperme...	0.30 to 1.20 grams	5 to 20 grains.
Abstract. belladonnae.....	3 to 10 centigrams	$\frac{1}{2}$ to $1\frac{1}{2}$ grains.
Abstract. cannab. ind.....	6 to 20 centigrams	1 to 3 grains
Abstract. conii.....	6 to 20 centigrams	1 to 3 grains
Abstract. coto.....	6 to 20 centigrams	1 to 3 grains
Abstract. digitalis.....	6 to 20 centigrams	1 to 3 grains
Abstract. gelsemii.....	6 to 20 centigrams	1 to 3 grains
Abstract. hyoscyami.....	12 to 40 centigrams	2 to 6 grains
Abstract. ignatie.....	6 to 20 centigrams	1 to 3 grains
Abstract. ipecac.....	0.20 to 2 grams	3 to 30 grains
Abstract. jalape.....	0.40 to 2 grams	6 to 30 grains.
Abstract. nuc. vom.....	6 to 20 centigrams	1 to 3 grains.
Abstract. phytolaccae.....	0.30 to 1 gram	5 to 15 grains.
Abstract. pilocarpi.....	0.40 to 2 grams	6 to 30 grains.
Abstract. podophylli.....	25 to 60 centigrams	1 to 10 grains
Abstract. sanguin. alterat.	6 to 20 centigrams	1 to 3 grains
Abstract. sanguin. emet....	0.60 to 1.20 Gm.....	10 to 15 grains.
Abstract. senega.....	25 to 60 centigrams	4 to 10 grains
Abstract. valeriana.....	0.60 to 1.20 Gm.....	10 to 15 grains.
Abstract. veratr. vir.....	6 to 20 centigrams	1 to 3 grains.
Acet. lobelia.....	1 to 4 C. c.....	15 to 60 minims.
Acet. opii.....	0.30 to 1 C. c.....	5 to 16 minims.
Acet. sanguinar.....	1 to 2 C. c.....	15 to 30 minims.
Acet. scillae.....	0.60 to 2 C. c.....	10 to 30 minims.
Acid. acet. dil.....	4 to 6 C. c.....	60 to 90 minims.
Acid. arsenios.....	1 to 5 milligrams	$\frac{1}{2}$ to $\frac{1}{4}$ grain.
Acid. benzoic.....	0.30 to 1 Gm.....	5 to 15 grains
Acid. borie.....	30 to 60 centigrams	5 to 10 grains.
Acid. carbolic.....	6 to 20 centigrams	1 to 3 grains
Acid. gallic.....	0.20 to 1 Gm.....	3 to 15 grains
Acid. gall. in albuminuria	0.60 to 4 Gm.....	10 to 60 grains
Acid. hydrobrom. 34 %.....	0.60 to 1 Gm.....	10 to 15 grains
Acid. hydrobrom. dil.....	1 to 2.50 C. c.....	15 to 40 minims.
Acid. hydrochlor.....	20 to 60 centigrams	3 to 10 grains.

DOSE TABLE.—Continued.

Remedies.	Dose in metric weights or measures.	Dose in apothecaries weights or measures.
Acid hydrochlor. dil.....	0.60 to 2 C. c.....	10 to 30 minims
Acid. hydrocyan. dil.....	0.10 to 0.30 C. c.....	2 to 6 minims
Acid. lactic.....	1 to 4 Gm.....	15 to 60 grains
Acid. nitr.....	20 to 60 centigrams	3 to 10 grains
Acid. nitr. dil.....	0.60 to 2 C. c.....	10 to 30 minims
Acid. nitro-hydrochlor.....	20 to 60 centigrams	3 to 10 grains
Acid. nitro-hydrochlor. dil.....	0.30 to 1.30 C. c.....	5 to 20 minims
Acid. phosphoric (50 %).....	0.20 to 1 Gm.....	3 to 15 grains
Acid. phosphoric dil.....	0.60 to 4 C. c.....	10 to 60 minims
Acid. salicyl.....	0.30 to 1 Gm.....	5 to 15 grains
Acid. sulphuric.....	0.30 to 0.60 Gm.....	5 to 10 grains
Acid. sulphuric dil.....	0.30 to 2 C. c.....	5 to 30 minims
Acid. sulphuric arom.....	0.30 to 2 C. c.....	5 to 30 minims
Acid. sulphurous.....	2 to 4 C. c.....	30 to 60 minims
Acid. tannic.....	10 to 60 centigrams	2 to 10 grains
Aconitina (white crystals).....	0.15 to 0.30 milligr.	1/8 to 2/8 grain
Aloe.....	10 to 30 centigrams	2 to 5 grains
Molnum.....	6 to 20 centigrams	1 to 3 grains
Alumen.....	0.50 to 1 Gm.....	10 to 15 grains
Ammonii benzoas.....	0.50 to 1.30 Gm.....	10 to 20 grains
Ammonii bromid.....	0.30 to 2 Gm.....	5 to 30 grains
Ammonii carb.....	20 to 60 centigrams	3 to 10 grains
Ammonii chlorid.....	1 to 2 Gm.....	15 to 30 grains
Ammonii iodid.....	0.20 to 1 Gm.....	3 to 15 grains
Ammonii phosph.....	30 to 130 Gm.....	5 to 20 grains
Ammonii picras.....	0.50 to 3 centigrams	1/4 to 1 1/2 grain
Ammonii sulph.....	0.20 to 1 Gm.....	3 to 15 grains
Ammonii valer.....	0.20 to 1 Gm.....	3 to 15 grains
Amyl nitris.....	0.10 to 0.40 C. c.....	2 to 5 minims
Amylum iodatum.....	0.20 to 2 Gm.....	3 to 30 grains
Ant. et pot. tartr.; diaph.....	4 to 10 milligrams	1/8 to 1/4 grain
Ant. et pot. tartr.; emetic.....	6 to 10 centigrams	1 to 2 grains
Antimonii oxid.....	3 to 10 centigrams	1 1/2 to 2 grains
Antimonii oxysulphuret.....	3 to 10 centigrams	1 1/2 to 2 grains
Antimonii sulphid.....	3 to 10 centigrams	1 1/2 to 2 grains
Antimonii sulphuret.....	3 to 10 centigrams	1 1/2 to 2 grains
Apomorph. hydrochlor.....	3 to 6 milligrams.....	3/8 to 1/2 grain
Aqua ammonia.....	0.30 to 2 C. c.....	6 to 30 minims
Aqua amygd. amor.....	10 to 15 C. c.....	2 to 4 fl. drachms.
Aqua camphoræ.....	15 to 60 C. c.....	1/2 to 2 fl. oz.
Aqua chlori.....	4 to 15 C. c.....	1 to 4 fl. drachms.
Aqua creasoti.....	4 to 15 C. c.....	1 to 4 fl. drachms.
Aqua laurocerasi.....	0.30 to 2 C. c.....	6 to 30 minims
Argentii iodidum.....	3 to 10 centigrams	1/2 to 2 grains
Argentii nitras.....	1 to 2 centigrams	1/6 to 1/3 grain
Argentii oxid.....	3 to 10 centigrams	1/2 to 2 grains
Arsenii iodidum.....	1 to 2.50 milligrams	1/20 to 1/10 grain
Asafoetida.....	0.30 to 1.30 Gm.....	5 to 20 grains
Atropina.....	0.50 to 2 milligrams	1/28 to 1/12 grain
Atropinae sulph.....	0.50 to 2 milligrams	1/28 to 1/12 grain
Auri et sodii chlorid.....	2 to 4 milligrams.....	1/32 to 1/16 grain
Bebeerinae sulph.....	20 to 60 centigrams	3 to 10 grains
Berberina and its salts.....	0.20 to 1 Gm.....	3 to 15 grains
Bismuthi citras.....	0.20 to 1 Gm.....	3 to 15 grains
Bismuthi et ammon. citr.....	0.10 to 1 Gm.....	1 to 15 grains

DOSE TABLE—Continued.

Remedies.	Dose in metric weights or measures.	Dose in apothecaries' weights or measures.
Bismuthi subcarb.....	0.40 to 2 Gm.....	6 to 30 grains
Bismuthi subnitr.....	0.40 to 2 Gm.....	6 to 30 grains
Bismuthi tannas.....	0.40 to 2 Gm.....	6 to 30 grains.
Bismuthi valer.....	6 to 20 centigrams	1 to 3 grains.
Brayern.....	8 to 15 Gm.....	2 to 6 drachms
Bromum.....	3 to 10 centigrams	$\frac{1}{2}$ to 2 grains
Brucina.....	1 to 4 milligrams..	$\frac{1}{32}$ to $\frac{1}{16}$ grain.
Caffeina.....	6 to 30 centigrams	1 to 5 grains
Caffeina citras.....	6 to 30 centigrams.	1 to 5 grains.
Calcii bromidum.....	0.30 to 2 Gm.....	5 to 30 grains
Calcii carb.....	1 to 4 Gm.....	15 to 60 grains.
Calcii hypophosphis.....	0.20 to 1 Gm.....	3 to 15 grains.
Calcii iodidum.....	6 to 20 centigrams	1 to 3 grains
Calcii phosphas.....	1 to 2 Gm.....	15 to 30 grains.
Calx sulphurata.....	2 to 6 centigrams	$\frac{1}{32}$ to 1 grain.
Cambogium.....	6 to 25 centigrams	1 to 4 grains.
Camphora.....	20 to 60 centigrams	3 to 10 grains.
Camph. monobrom.....	10 to 30 centigrams	2 to 5 grains.
Cantharis.....	3 to 6 centigrams	$\frac{1}{2}$ to 2 grains.
Capsicum.....	6 to 20 centigrams	1 to 3 grains.
Castoreum.....	0.40 to 1 Gm.....	6 to 15 grains.
Catechu.....	1 to 2 Gm.....	15 to 30 grains.
Cerii nitras.....	6 to 20 centigrams	1 to 3 grains.
Cerii oxalas.....	6 to 20 centigrams	1 to 3 grains.
Chinoidinum.....	0.20 to 2 Gm.....	3 to 30 grains.
Chloral.....	0.20 to 1 Gm.....	3 to 15 grains.
Chloroformum.....	0.05 to 0.30 C. c.....	1 to 5 minims.
Chrysarobinum.....	0.20 to 1 Gm.....	3 to 15 grains.
Cinchona.....	1 to 4 Gm.....	15 to 60 grains.
Cinchonidina, and its salts.....	0.06 to 2 Gm.....	1 to 30 grains.
Cinchonina, and its salts.....	0.06 to 2 Gm.....	1 to 30 grains.
Cinnamomum.....	0.40 to 2 Gm.....	6 to 30 grains.
Codeina.....	3 to 10 centigrams	$\frac{1}{2}$ to 2 grains.
Confectio senna.....	4 to 10 grams.....	1 to 2 grains.
Conina, and its salts.....	1 to 2 milligrams	$\frac{1}{64}$ to $\frac{1}{32}$ grain.
Copaiba.....	1 to 4 C. c.....	15 to 60 minims.
Cotoina.....	1 to 3 centigrams	$\frac{1}{16}$ to $\frac{1}{8}$ grain.
Creasotum.....	0.05 to 0.20 C. c.....	1 to 3 minims
Creta præpar.....	1 to 5 Gm.....	15 to 75 grains.
Croton chloral.....	5 to 60 centigrams	1 to 10 grains.
Cubeba.....	1 to 4 Gm.....	15 to 60 grains.
Cupri acetas.....	3 to 40 centigrams	$\frac{1}{2}$ to 6 grains.
Cupri sulphas.....	3 to 60 centigrams	$\frac{1}{2}$ to 10 grains.
Cuprum ammon.....	1 to 6 centigrams	$\frac{1}{16}$ to 1 grain.
Curare.....	2 to 10 milligrams	$\frac{1}{32}$ to $\frac{1}{16}$ grain.
Curarina.....	1 to 3 milligrams	$\frac{1}{64}$ to $\frac{1}{32}$ grain.
Decoct. aloes comp.....	15 to 60 C. c.....	$\frac{1}{2}$ to 2 fl. ozs.
Decoct. sarsap. comp.....	50 to 200 C. c.....	2 to 6 fl. ozs
Digitalinum.....	1 to 2 milligrams	$\frac{1}{16}$ to $\frac{1}{8}$ grain.
Digitalis.....	3 to 10 centigrams	$\frac{1}{32}$ to 2 grains.
Duboisina, and its salts.....	0.50 to 1 milligram	$\frac{1}{128}$ to $\frac{1}{64}$ grain.
Elaterium; U. S. P., 1880.....	1 to 4 milligrams	$\frac{1}{64}$ to $\frac{1}{16}$ grain.
Elaterium; U. S. P., 1870.....	4 to 30 milligrams	$\frac{1}{16}$ to $\frac{1}{2}$ grain.
Emetina, and its salts, emetic.....	8 to 15 milligrams	$\frac{1}{16}$ to $\frac{1}{8}$ grain.
Emetina, and salts, diaph.....	0.50 to 2 milligr'm	$\frac{1}{64}$ to $\frac{1}{32}$ grain.

DOSE TABLE.—*Continued.*

Remedies.	Dose in metric weights or measures.	Dose in apothecaries' weights or measures.
Emulsio hydrocyan.....	2 to 4 C. c.....	$\frac{1}{2}$ to 1 fl. drachm.
Ergota.....	1 to 4 Gm.....	15 to 60 grains.
Ergotinum.....	10 to 50 centigrams	2 to 8 grains.
Eserina and its salts.....	1 to 3 milligrams	$\frac{1}{2}$ to $\frac{1}{20}$ grain.
Extr. absinthii.....	10 to 60 centigrams	2 to 6 grains.
Extr. absinthii fluid.....	1 to 2 C. c.....	15 to 30 minims.
Extr. achilleæ.....	20 to 60 centigrams	3 to 10 grains.
Extr. achilleæ fluid.....	1 to 4 C. c.....	15 to 60 minims.
Extr. aconi fol. Engl.....	2 to 6 centigrams	$\frac{1}{2}$ to 1 grain.
Extr. aconi fol. U. S. P. 70.....	3 to 12 centigrams	$\frac{1}{2}$ to 2 grains.
Extr. aconi fol. fluid.....	0.05 to 0.30 C. c.....	1 to 5 minims.
Extr. aconi rad. U. S. P. 80.....	5 to 15 milligrams	$\frac{1}{2}$ to 1 grain.
Extr. aconi rad. fluid.....	0.03 to 0.15 C. c.....	2 to 2½ minims.
Extr. alchædis fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. alchædis rad. fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. alchædis aquos.....	3 to 20 centigrams	$\frac{1}{2}$ to 3 grains.
Extr. alston. constr. fl.....	4 to 15 C. c.....	1 to 4 fl. drachms.
Extr. angusturæ fl.....	1 to 3 C. c.....	15 to 45 minims.
Extr. angusturæ rad. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. anthemidis.....	10 to 60 centigrams	2 to 10 grains.
Extr. anthemidis fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. apocyn. andros fl.....	0.50 to 2 C. c.....	8 to 50 minims.
Extr. apocyn. cannabin. fl.....	0.50 to 2 C. c.....	8 to 30 minims.
Extr. aralia hisp. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. aralia nudic. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. aralia racem. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. aralia spin. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. arisa. fl.....	3 to 5 C. c.....	45 to 75 minims.
Extr. arisa. fol.....	20 to 50 centigrams	3 to 8 grains.
Extr. arisa. fl.....	0.50 to 1 C. c.....	5 to 15 minims.
Extr. arisa. rad.....	10 to 30 centigrams	2 to 5 grains.
Extr. arisa. rad. fl.....	0.50 to 1 C. c.....	5 to 15 minims.
Extr. arisa. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. ari triphylli fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. asari fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. asclep. incarn. fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. asclep. syr. fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. asclep. tuber. fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. aspidii fl.....	5 to 15 C. c.....	1 to 4 fl. drachms.
Extr. aspidosperme fl.....	1 to 3 C. c.....	15 to 45 minims.
Extr. auranti cort. fl.....	1 to 10 C. c.....	$\frac{1}{4}$ to 2½ fl. drachms.
Extr. azedarach fl.....	1 to 5 C. c.....	15 to 75 minims.
Extr. baptisæ fl.....	0.50 to 2 C. c.....	7 to 30 minims.
Extr. bellad. fol. (Eng.).....	1 to 4 centigrams	$\frac{1}{8}$ to $\frac{1}{20}$ grain.
Extr. bellad. alcohol.....	1 to 3 centigrams	$\frac{1}{8}$ to $\frac{1}{20}$ grain.
Extr. bellad. fol. fl.....	0.20 to 0.40 C. c.....	3 to 6 minims.
Extr. bellad. rad.....	8 to 15 milligrams	$\frac{1}{8}$ to $\frac{1}{4}$ grain.
Extr. bellad. rad. fl.....	0.10 to 0.20 C. c.....	1 to 3 minims.
Extr. berber. aquifol. fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. berber. vulg. fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. boldi fl.....	0.20 to 1 C. c.....	3 to 15 minims.
Extr. braveræ fl.....	4 to 15 C. c.....	2 to 4 fl. drachms.
Extr. bryoniæ fl.....	1 to 4 C. c.....	15 to 60 minims.
Extr. buchu fl.....	2 to 10 C. c.....	$\frac{1}{2}$ to 2½ fl. drachms.
Extr. calami fl.....	1 to 4 C. c.....	15 to 60 minims.

DOSE TABLE.—*Continued.*

Remedies.	Dose in metric weights or measures.	Dose in apothecaries' weights or measures.
Extr. calend. fl.	1 to 4 C. c.	15 to 60 minims.
Extr. calumbæ.	20 to 60 centigrams	3 to 10 grains.
Extr. calumbæ fl.	1 to 4 C. c.	15 to 60 minims.
Extr. canella fl.	1 to 4 C. c.	15 to 60 minims.
Extr. cannab. Amer. fl.	0.20 to 1 C. c.	3 to 15 minims.
Extr. cannab. ind.	1 to 3 centigrams	$\frac{1}{6}$ to $\frac{1}{2}$ grain.
Extr. cannab. ind. fl.	0.20 to 0.40 C. c.	3 to 6 minims.
Extr. cantharidis fl.	0.06 to 0.20 C. c.	1 to 3 minims.
Extr. capsici fl.	0.06 to 0.20 C. c.	1 to 3 minims.
Extr. cardam. comp. fl.	1 to 3 C. c.	15 to 45 minims.
Extr. cardui bened. fl.	1 to 4 C. c.	15 to 60 minims.
Extr. carui.	1 to 4 C. c.	15 to 60 grains.
Extr. cascarilla fl.	3 to 10 C. c.	$\frac{3}{4}$ to 2½ fl. drachms.
Extr. castaneæ fl.	3 to 10 C. c.	$\frac{1}{2}$ to 2½ fl. drachms.
Extr. catariæ fl.	1 to 5 C. c.	$\frac{1}{4}$ to 1¼ fl. drachms.
Extr. catechu liquid	0.06 to 2 C. c.	8 to 30 minims.
Extr. caulophylli fl.	1 to 2 C. c.	15 to 30 minims.
Extr. chelidonii fl.	1 to 2 C. c.	15 to 30 minims.
Extr. chelonis fl.	2 to 4 C. c.	30 to 60 minims.
Extr. chinaph. fl.	3 to 5 C. c.	$\frac{3}{4}$ to 1¼ fl. drachms.
Extr. chionanthi fl.	5 to 10 C. c.	$\frac{3}{4}$ to 2½ fl. drachms.
Extr. chirate fl.	2 to 5 C. c.	$\frac{1}{2}$ to 1¼ fl. drachms.
Extr. cimicifuge fl.	0.50 to 2 C. c.	8 to 30 minims.
Extr. cinchonæ.	1 to 2 Gm.	15 to 30 grains.
Extr. cinchonæ fl.	2 to 4 C. c.	30 to 60 minims.
Extr. cinchonæ arom. fl.	2 to 4 C. c.	30 to 60 minims.
Extr. cinchonæ comp. fl.	2 to 5 C. c.	$\frac{1}{2}$ to 1¼ fl. drachms.
Extr. cocculi fl.	0.05 to 0.20 C. c.	1 to 3 minims.
Extr. colch. rad.	2 to 10 centigrams	$\frac{1}{2}$ to 1½ grains.
Extr. colch. rad. fl.	0.20 to 1 C. c.	3 to 15 minims.
Extr. colch. sem. fl.	0.10 to 0.60 C. c.	$\frac{1}{2}$ to 10 minims.
Extr. collinsoniæ fl.	2 to 4 C. c.	30 to 60 minims.
Extr. colocynth.	10 to 30 centigrams	1½ to 5 grains.
Extr. colocynth comp.	10 to 30 centigrams	1½ to 5 grains.
Extr. condurango fl.	0.0 to 2 C. c.	8 to 30 minims.
Extr. conii fol. (Engl.)	5 to 20 centigrams	1 to 4 grains.
Extr. conii fol. alc.; U.S.P. 70	5 to 10 centigrams	1 to 1½ grains.
Extr. con. (fr. alc.; U.S.P. 80)	2 to 6 centigrams	$\frac{1}{2}$ to 1 grain.
Extr. conii fol. fl.	0.20 to 1 C. c.	3 to 15 minims.
Extr. con. (fr. fl.; U.S.P. 80)	0.10 to 0.30 C. c.	$\frac{1}{2}$ to 5 minims.
Extr. convallariæ rad. fl.	1 to 2 C. c.	15 to 30 minims.
Extr. coptidis fl.	2 to 4 C. c.	30 to 60 minims.
Extr. corn. flor. fl.	2 to 4 C. c.	30 to 60 minims.
Extr. corydalis fl.	1 to 2 C. c.	15 to 30 minims.
Extr. coto fl.	0.20 to 1 C. c.	3 to 15 minims.
Extr. cubebæ fl.	1 to 2 C. c.	15 to 30 minims.
Extr. cyripedil fl.	1 to 4 C. c.	15 to 60 minims.
Extr. damianæ fl.	2 to 10 C. c.	$\frac{1}{2}$ to 2½ fl. drachms.
Extr. delphinii fl.	0.06 to 0.20 C. c.	1 to 3 minims.
Extr. digitalis.	1 to 3 centigrams	$\frac{1}{6}$ to $\frac{1}{2}$ grain.
Extr. digitalis fl.	0.10 to 0.40 C. c.	1 to 6 minims.
Extr. dioscoreæ fl.	1 to 2 Gm.	15 to 30 minims.
Extr. ditæ fl.	5 to 15 C. c.	1 to 4 fl. drachms.
Extr. draconii fl.	2 to 4 C. c.	30 to 60 grains.
Extr. droseræ fl.	0.30 to 0.60 C. c.	5 to 10 minims.

DOSE TABLE.—Continued

Remedies.	Dose in metric weights or measures	Dose in apothecaries' weights or measures.
Extr. dulcamare.....	0.30 to 1 Gm.....	5 to 15 grains.
Extr. dulcamare fl.....	4 to 8 C.c.....	1 to 2 fl. drachms.
Extr. ergotæ.....	10 to 50 centigrams.....	1½ to 8 grains.
Extr. ergotæ fl.....	1 to 4 C.c.....	15 to 60 minims.
Extr. eryodictyi fl.....	1 to 2 C.c.....	15 to 30 minims.
Extr. erythroxyl fl.....	2 to 8 C.c.....	½ to 2 fl. drachms.
Extr. eucalypti fl.....	1 to 4 C.c.....	15 to 60 minims.
Extr. euonymi fl.....	1 to 4 C.c.....	15 to 60 minims.
Extr. eupatori fl.....	2 to 4 C.c.....	30 to 60 minims.
Extr. euphorb. spec. fl.....	0.30 to 2 C.c.....	5 to 30 minims.
Extr. ferri pom.....	0.20 to 1 Gm.....	3 to 15 grains.
Extr. frangulæ fl.....	2 to 10 C.c.....	½ to 2½ fl. drachms.
Extr. frankeniæ fl.....	0.50 to 1 C.c.....	8 to 15 minims.
Extr. galliæ fl.....	3 to 8 C.c.....	½ to 2 fl. drachms.
Extr. gelsemii.....	0.10 to 0.50 C.c.....	2 to 8 minims.
Extr. gelsemii fl.....	0.10 to 0.50 C.c.....	1 to 8 minims.
Extr. gent. fl.....	2 to 4 C.c.....	30 to 60 minims.
Extr. gent. comp. fl.....	2 to 4 C.c.....	30 to 60 minims.
Extr. gent. quinquefl. fl.....	1 to 2 C.c.....	15 to 30 minims.
Extr. geranii fl.....	1 to 2 C.c.....	15 to 30 minims.
Extr. ggi fl.....	1 to 2 C.c.....	15 to 30 minims.
Extr. gillenii fl.....	1 to 2 C.c.....	15 to 30 minims.
Extr. gossypii fl.....	1 to 3 C.c.....	15 to 45 minims.
Extr. graminæ rad. cort. fl.....	3 to 8 C.c.....	½ to 2 fl. drachms.
Extr. grind. rob. fl.....	2 to 4 C.c.....	30 to 60 minims.
Extr. grind. squari fl.....	2 to 4 C.c.....	30 to 60 minims.
Extr. guaiaci ligni fl.....	2 to 4 C.c.....	30 to 60 minims.
Extr. guaranæ fl.....	1 to 2 C.c.....	15 to 30 minims.
Extr. hamotoxyl.....	0.50 to 2 Gm.....	8 to 30 grains.
Extr. hamotoxyl fl.....	2 to 4 C.c.....	30 to 60 minims.
Extr. hamamelid fl.....	2 to 6 C.c.....	30 to 90 minims.
Extr. helleb. nigris.....	3 to 20 centigrams.....	½ to 3 grains.
Extr. helleb. nigris fl.....	0.30 to 1 C.c.....	5 to 15 minims.
Extr. helonii fl.....	0.50 to 2 C.c.....	8 to 30 minims.
Extr. hepaticæ fl.....	2 to 4 C.c.....	30 to 60 minims.
Extr. humuli.....	0.20 to 1 Gm.....	3 to 15 grains.
Extr. humuli fl.....	2 to 4 C.c.....	30 to 60 minims.
Extr. hydrangææ fl.....	2 to 4 C.c.....	30 to 60 minims.
Extr. hydrastis.....	20 to 60 centigrams.....	3 to 10 grains.
Extr. hydrastis fl.....	0.50 to 2 C.c.....	8 to 30 minims.
Extr. hyoscyami (Engl.).....	5 to 25 centigrams.....	1 to 4 grains.
Extr. hyoscyami alc.....	5 to 10 centigrams.....	1 to 2 grains.
Extr. hyoscyami sol. fl.....	0.20 to 1 C.c.....	3 to 15 minims.
Extr. hyoscyaminæ fl.....	0.10 to 0.50 C.c.....	2 to 8 minims.
Extr. ignatiæ.....	2 to 8 centigrams.....	½ to 1½ grains.
Extr. ignatiæ fl.....	0.05 to 0.30 C.c.....	1 to 6 minims.
Extr. ipocac fl.....	0.20 to 4 C.c.....	3 to 60 minims.
Extr. iridis versicol.....	0.20 to 0.30 Gm.....	3 to 6 grains.
Extr. irid. versicol. fl.....	1 to 2 C.c.....	15 to 30 minims.
Extr. jalapæ; u. s. p., 1870.....	30 to 60 centigrams.....	5 to 10 grains.
Extr. jalapæ alc.....	20 to 40 centigrams.....	3 to 6 grains.
Extr. jalapæ fl.....	1 to 4 C.c.....	15 to 60 minims.
Extr. juglandis.....	1 to 2 Gm.....	15 to 30 grains.
Extr. juglandis fl.....	3 to 8 C.c.....	½ to 2 fl. drachms.
Extr. junip. fl.....	2 to 4 C.c.....	30 to 60 minims.

DOSE TABLE.—Continued.

Remedies.	Dose in metric weights or measures.	Dose in apothecaries' weights or measures.
Extr. kamala fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. kino, liquid.....	1 to 2 C. c.....	15 to 30 minims.
Extr. krameria.....	0.30 to 1 Gm.....	5 to 15 grains.
Extr. krameria fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. lactuca.....	0.30 to 1 Gm.....	5 to 15 grains.
Extr. lactuca fl.....	1 to 4 C. c.....	15 to 40 minims.
Extr. lactucarii fl.....	0.50 to 2 C. c.....	8 to 30 minims.
Extr. lappa fl.....	4 to 8 C. c.....	1 to 2 fl. drachms.
Extr. laris fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. leonuri fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. leptandra.....	20 to 60 centigrams	3 to 10 grains.
Extr. leptandra fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. lobelia fl.....	0.30 to 2 C. c.....	5 to 30 C. c.
Extr. lobelia sem. fl.....	0.20 to 1 C. c.....	3 to 15 C. c.
Extr. lupulini fl.....	0.30 to 1 C. c.....	5 to 15 minims.
Extr. lycopi fl.....	0.30 to 2 C. c.....	5 to 30 minims.
Extr. mali.....	5 to 10 Gm.....	1 to $2\frac{1}{2}$ drachms.
Extr. manzanita fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. marrubii fl.....	4 to 8 C. c.....	1 to 2 fl. drachms.
Extr. matico fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. matricaria fl.....	0.50 to 2 C. c.....	8 to 30 minims.
Extr. menispermii fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. methystice fl.....	1 to 4 C. c.....	15 to 60 minims.
Extr. mezerel.....	3 to 6 centigrams	$\frac{1}{2}$ to 1 grain.
Extr. mezerel fl.....	0.20 to 0.60 C. c....	3 to 10 minims.
Extr. micromeria fl.....	1 to 4 C. c.....	15 to 60 minims.
Extr. mitchella fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. myrica fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. nectandra fl.....	4 to 15 C. c.....	1 to 4 fl. drachms.
Extr. nuc. vom.....	2 to 10 centigrams	$\frac{1}{2}$ to $1\frac{1}{2}$ grains.
Extr. nuc. vom. fl.....	0.06 to 0.30 C. c....	1 to 5 minims.
Extr. nuphar fl.....	0.30 to 1 C. c.....	5 to 15 minims.
Extr. nymphaea fl.....	0.30 to 1 C. c.....	5 to 15 minims.
Extr. oenothera fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. opii.....	1 to 3 centigrams	$\frac{1}{8}$ to $\frac{1}{2}$ grain.
Extr. papaveris.....	3 to 12 centigrams	$\frac{1}{2}$ to 2 grains.
Extr. papaveris fl.....	1 to 3 C. c.....	15 to 45 minims.
Extr. pareira fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. petroselinii fl.....	4 to 8 C. c.....	1 to 2 fl. drachms.
Extr. phellandri fl.....	4 to 8 C. c.....	1 to 2 fl. drachms.
Extr. phoradendri fl.....	2 to 4 C. c.....	$\frac{1}{2}$ to 1 fl. drachm.
Extr. physostigma.....	4 to 10 milligrams	$\frac{1}{16}$ to $\frac{1}{8}$ grain.
Extr. physostigma fl.....	0.06 to 0.20 C. c....	1 to 3 minims.
Extr. phytolaccae baccar fl	0.30 to 2 C. c.....	5 to 30 minims.
Extr. phytolaccae rad.....	6 to 20 centigrams	1 to 3 grains.
Extr. phytolaccae rad. fl.	0.30 to 2 C. c.....	5 to 30 minims.
Extr. pilocarpii fl.....	1 to 4 C. c.....	15 to 60 minims.
Extr. pimenta fl.....	1 to 3 C. c.....	15 to 45 minims.
Extr. piper. nigr. fl.....	1 to 3 C. c.....	15 to 45 minims.
Extr. piscidia fl.....	1 to 4 C. c.....	15 to 60 minims.
Extr. podophylli.....	3 to 10 centigrams	$\frac{1}{2}$ to $1\frac{1}{2}$ grains.
Extr. podophylli fl.....	0.50 to 2 C. c.....	8 to 30 minims.
Extr. polygoni fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. polygonati fl.....	0.30 to 1 C. c.....	5 to 15 minims.
Extr. populi fl.....	2 to 4 C. c.....	30 to 60 minims.

POST. TABLE.—*Continued*

Remedies.	Dose in metric weights or measures.	Dose in apothecaries' weights or measures.
Extr. prinos fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. prun. virg. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. ptelea fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. pulsatilla fl.....	0.10 to 0.30 C. c.....	2 to 5 minims.
Extr. quassia.....	6 to 30 centigrms.....	1 to 5 grains.
Extr. quassia fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. quercus fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. rhamni cath. fr. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. rhamni pursh. cort. fl.....	2 to 8 C. c.....	30 to 120 minims.
Extr. rhei.....	0.30 to 1 Gm.....	5 to 15 grains.
Extr. rhei fl.....	1 to 3 C. c.....	15 to 45 minims.
Extr. rhois arom. fl.....	1 to 4 C. c.....	15 to 60 minims.
Extr. rhois glabr. cort. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. rhois glabr. fruct. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. rhois toxicod. fl.....	0.06 to 0.40 C. c.....	1 to 6 minims.
Extr. ricini fol. fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. rose fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. rubi fl.....	1 to 1 C. c.....	15 to 60 minims.
Extr. rumicis fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. rutæ fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. sabatie fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. sabine fl.....	0.30 to 1 C. c.....	5 to 15 minims.
Extr. salicis fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. salvia fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. sambuci fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. sanguin fl.....	0.30 to 1 C. c.....	5 to 15 minims.
Extr. santali citr. fl.....	4 to 8 C. c.....	1 to 2 fl. drachms.
Extr. santonica fl.....	1 to 4 C. c.....	15 to 60 minims.
Extr. sarsap fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. comp. fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. sassapar. fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. seille fl.....	0.30 to 2 C. c.....	5 to 30 minims.
Extr. seille comp. fl.....	0.30 to 2 C. c.....	5 to 30 minims.
Extr. scoparii fl.....	2 to 4 C. c.....	$\frac{1}{2}$ to 1 fl. drachm.
Extr. scutellaria fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. senecionis fl.....	4 to 8 C. c.....	1 to 2 fl. drachms.
Extr. senega fl.....	0.30 to 1 C. c.....	8 to 15 minims.
Extr. senna fl.....	4 to 15 C. c.....	1 to 4 fl. drachms.
Extr. serpent. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. sinarubæ fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. solidag. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. spigelia fl.....	1 to 4 C. c.....	15 to 60 minims.
Extr. spigelia et senna fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. stillingia fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. stillingia comp. fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. stramonii (Engl.).....	3 to 6 centigrms.....	$\frac{1}{2}$ to 1 grain.
Extr. stramonii fol. alc.....	2 to 4 centigrms.....	$\frac{1}{2}$ to 2 grain.
Extr. stramonii sem.....	1 to 3 centigrms.....	$\frac{1}{6}$ to $\frac{1}{2}$ grain.
Extr. stramonii fl.....	0.06 to 0.40 C. c.....	1 to 6 minims.
Extr. subul fl.....	1 to 4 C. c.....	15 to 60 minims.
Extr. taraxaci.....	0.30 to 1 Gm.....	5 to 15 grains.
Extr. taraxaci fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. thuyæ fl.....	0.50 to 1 C. c.....	8 to 15 minims.
Extr. toxicodendri fl.....	0.06 to 0.30 C. c.....	1 to 5 minims.
Extr. trifol. prat. fl.....	4 to 8 C. c.....	1 to 2 fl. drachms.

DOSE TABLE—Continued

Remedies	Dose in metric weights or measures.	Dose in apothecaries weights or measures
Extr. trillii fl.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms.
Extr. trit. rep. fl.....	1 to 15 C. c.....	1 to 4 fl. drachms
Extr. tussilag. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. urticae rad. fl.....	0.30 to 1 C. c.....	5 to 15 minims.
Extr. ustilag. maid. fl.....	1 to 4 C. c.....	15 to 60 minims.
Extr. uva-ursi fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. vaccin. crassifol. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. valer.....	0.30 to 1 Gm.....	5 to 15 grains.
Extr. valer. fl.....	2 to 4 C. c.....	30 to 60 minims.
Extr. veratr. vir. fl.....	0.10 to 0.50 C. c.....	2 to 8 minims.
Extr. verbena fl.....	1 to 4 C. c.....	15 to 60 minims.
Extr. viburni opuli fl.....	4 to 8 C. c.....	1 to 2 fl. drachms.
Extr. viburni [prunifol.] fl.....	4 to 8 C. c.....	1 to 2 fl. drachms.
Extr. xanthoxyli cort. fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. xanthoxyli fruct. fl.....	1 to 2 C. c.....	15 to 30 minims.
Extr. zingiberis fl.....	0.50 to 2 C. c.....	8 to 30 minims.
Fel bovis purif.....	20 to 40 centigrams	3 to 6 grains.
Ferri arsen.....	3 to 30 milligrams	$\frac{1}{2}$ to $\frac{1}{2}$ grain.
Ferri benzoas.....	5 to 30 centigrams	1 to 5 grains.
Ferri bromid.....	5 to 30 centigrams	1 to 5 grains.
Ferri carb. sacch.....	0.25 to 1 Gm.....	4 to 15 grains.
Ferri chlorid.....	5 to 20 centigrams	1 to 3 grains.
Ferri citr.....	30 to 60 centigrams	5 to 10 grains.
Ferri et ammon. citr.....	30 to 60 centigrams	5 to 10 grains.
Ferri et ammon. sulph.....	30 to 60 centigrams	5 to 10 grains.
Ferri et ammon. tartr.....	0.30 to 1 Gm.....	5 to 15 grains.
Ferri et cinchonid. citr.....	30 to 60 centigrams	5 to 10 grains.
Ferri et pot. tartr.....	1 to 4 Gm.....	15 to 60 grains.
Ferri et quin. citr.....	30 to 60 centigrams	5 to 10 grains.
Ferri et strychn. citr.....	5 to 30 centigrams	1 to 5 grains.
Ferri ferrocyanid.....	20 to 30 centigrams	3 to 5 grains.
Ferri hypophosphis.....	30 to 60 centigrams	5 to 10 grains.
Ferri iodidum.....	5 to 30 centigrams	1 to 5 grains.
Ferri iodidum sacch.....	10 to 60 centigrams	2 to 10 grains.
Ferri lactas.....	5 to 20 centigrams	1 to 3 grains.
Ferri oxalas.....	5 to 20 centigrams	1 to 3 grains.
Ferri oxid. magnet.....	30 to 60 centigrams	5 to 10 grains.
Ferri oxid. hydrat.....	15 to 60 Gm.....	$\frac{1}{2}$ to 2 ounces.
Ferri phosphas.....	5 to 30 centigrams	1 to 5 grains.
Ferri pyrophosphas.....	5 to 30 centigrams	1 to 5 grains.
Ferri subcarb.....	0.30 to 2 Gm.....	5 to 30 grains.
Ferri sulphas.....	5 to 20 centigrams	1 to 3 grains.
Ferri sulphas exsiccant.....	3 to 10 centigrams	$\frac{1}{2}$ to $\frac{1}{2}$ grains
Ferri valer.....	5 to 20 centigrams	1 to 3 grains.
Ferrum dialys.....	0.06 to 1 C. c.....	1 to 15 minims.
Ferrum reduct.....	6 to 30 centigrams	1 to 5 grains.
Guarana.....	0.50 to 2 Gm.....	8 to 30 grains.
Hydrarg. chlorid. corros.....	1 to 6 milligrams	$\frac{1}{64}$ to $\frac{1}{16}$ grain.
Hydrarg. chlorid. mite.....	1 to 50 centigrams	$\frac{1}{4}$ to 8 grains.
Hydrarg. cyanid.....	4 to 30 milligrams	$\frac{1}{16}$ to $\frac{1}{2}$ grain.
Hydrarg. iodid. flav.....	1 to 6 centigrams	$\frac{1}{10}$ to 1 grain.
Hydrarg. iodid. rubr.....	4 to 30 milligrams	$\frac{1}{16}$ to $\frac{1}{2}$ grain.
Hydrarg. iodid. vir.....	1 to 6 centigrams	$\frac{1}{10}$ to 1 grain.
Hydrarg. oxid. flav.....	4 to 30 milligrams	$\frac{1}{16}$ to $\frac{1}{2}$ grain.
Hydrarg. oxid. nigr.....	6 to 60 milligrams	$\frac{1}{10}$ to 1 grain.

DOSE TABLE.—Continued.

Remedies.	Dose in metric weights or measures	Dose in apothecaries' weights or measures
Hydrarg. oxid. rubr.....	4 to 30 milligrams	$\frac{1}{16}$ to $\frac{1}{2}$ grain
Hydrarg. subsulphas flav.....	15 to 60 milligrams	$\frac{1}{4}$ to 1 grain
Hydrarg. c. creta.....	15 to 50 centigrams	$\frac{3}{8}$ to 3 grains
Hyoscyamina, and salts.....	0.50 to 2 milligrams	$\frac{1}{16}$ to $\frac{1}{8}$ grain
Infusum brayeræ.....	60 to 250 C. c.....	2 to 8 fl. ounces
Infusum digitalis.....	8 to 15 C. c.....	2 to 4 fl. drachms
Infusum sennæ comp.....	30 to 60 C. c.....	1 to 2 fl. ounces
Iodoformum.....	5 to 20 centigrams	1 to 3 grains
Iodum.....	1 to 3 centigrams	$\frac{1}{16}$ to $\frac{1}{8}$ grain
Ipecacuanha } expect.....	1 to 6 centigrams	$\frac{1}{16}$ to 1 grain
Ipecacuanha } emet.....	1 to 2 Gm.....	15 to 30 grains
Jalapa.....	1 to 2 Gm.....	15 to 30 grains
Kamala.....	4 to 8 Gm.....	1 to 2 drachms
Kino.....	0.50 to 2 Gm.....	8 to 30 grains
Lactucarium.....	0.50 to 1 Gm.....	8 to 15 grains
Liq. ammon. acet.....	8 to 30 C. c.....	2 to 8 fl. drachms
Liq. acidi arseniosi.....	0.10 to 0.50 C. c.....	2 to 7 minims
Liq. arsen. et hydr. iod.....	0.10 to 0.50 C. c.....	2 to 7 minims
Liq. ferri chloridi.....	0.10 to 0.60 C. c.....	2 to 10 minims
Liq. ferri dialys.....	0.06 to 1 C. c.....	1 to 15 minims
Liq. ferri nitrat.....	0.50 to 1 C. c.....	8 to 15 minims
Liq. pepsini.....	8 to 15 C. c.....	2 to 4 fl. drachms
Liquor potassæ.....	0.30 to 2 C. c.....	5 to 30 minims
Liquor potassii arsenit.....	0.15 to 0.50 C. c.....	3 to 7 minims
Liquor potassii citrat.....	8 to 15 C. c.....	2 to 4 fl. drachms
Liquor sodæ.....	0.30 to 2 C. c.....	5 to 30 minims
Liquor sodii arseniatis.....	0.15 to 0.50 C. c.....	3 to 7 minims
Lithii benzoas.....	10 to 30 centigrams	2 to 5 grains
Lithii bromid.....	6 to 20 centigrams	1 to 3 grains
Lithii carb.....	10 to 40 centigrams	2 to 6 grains
Lithii citr.....	10 to 30 centigrams	2 to 5 grains
Lithii salicylas.....	10 to 50 centigrams	2 to 8 grains
Lupulinum.....	30 to 60 centigrams	5 to 10 grains
Magnesia.....	1 to 4 Gm.....	15 to 60 grains
Magnesi carb.....	1 to 4 Gm.....	15 to 60 grains
Magnesi citr. gran.....	8 to 30 Gm.....	2 to 8 drachms
Magnesi sulphas.....	8 to 30 Gm.....	2 to 8 drachms
Magnesi sulphis.....	0.30 to 2 Gm.....	8 to 30 grains
Mangani sulphas.....	10 to 60 centigrams	2 to 10 grains
Manna.....	30 to 60 Gm.....	1 to 2 ounces
Massa copaibæ.....	0.30 to 2 Gm.....	5 to 30 grains
Massa ferri carb.....	0.30 to 1 Gm.....	5 to 15 grains
Massa hydrarg.....	0.06 to 1 Gm.....	1 to 15 grains
Mist. ammoniaci.....	15 to 30 C. c.....	4 to 8 fl. drachms
Mist. anafœtidæ.....	15 to 30 C. c.....	4 to 8 fl. drachms
Mist. chloroformi.....	15 to 30 C. c.....	4 to 8 fl. drachms
Mist. cretæ.....	30 to 60 C. c.....	1 to 2 fl. ounces
Mist. ferri comp.....	15 to 60 C. c.....	$\frac{1}{2}$ to 2 fl. ounces
Mist. ferri et amm. acet.....	15 to 30 C. c.....	$\frac{1}{2}$ to 1 fl. ounce
Mist. glycyrrh. comp.....	4 to 15 C. c.....	1 to 4 fl. drachms
Mist. magnes. et anafœt.....	4 to 15 C. c.....	1 to 4 fl. drachms
Mist. potassii citr.....	15 to 60 C. c.....	$\frac{1}{2}$ to 2 fl. ounces
Mist. rhei et sodæ.....	15 to 30 C. c.....	$\frac{1}{2}$ to 1 fl. ounce
Morphina, and its salts.....	4 to 30 milligrams	$\frac{1}{16}$ to $\frac{1}{8}$ grain
Moschus.....	0.10 to 1 Gm.....	2 to 15 grains

DOSE TABLE.—Continued.

Remedies	Dose in metric weights or measures	Dose in apothecaries' weights or measures.
Narceina	1 to 10 centigrams	$\frac{1}{4}$ to 2 grains.
Nitroglycerinum	1 to 4 milligrams	$\frac{1}{32}$ to $\frac{1}{8}$ grain.
Nux vomica	5 to 30 centigrams	1 to 5 grains.
Oleoresina aspidii	1 to 4 Gm	15 to 60 grains.
Oleoresina capsici	1 to 3 milligrams	$\frac{1}{4}$ to $\frac{1}{2}$ grain.
Oleoresina cubeba	0.30 to 2 Gm	5 to 30 grains.
Oleoresina lupulini	10 to 30 centigrams	2 to 5 grains.
Oleoresina piperis	6 to 20 centigrams	1 to 3 grains.
Oleoresina zingiberis	6 to 20 centigrams	1 to 3 grains.
Oleum copaiba	0.50 to 1 C. c.	8 to 15 minims.
Oleum cubeba	1 to 2 C. c.	15 to 30 minims.
Oleum eriger	0.30 to 1 C. c.	5 to 15 minims.
Oleum eucalypti	0.50 to 2 C. c.	10 to 30 minims.
Oleum phosphoratum	5 to 20 centigrams	1 to 3 grains.
Oleum sabinæ	0.50 to 0.20 C. c.	1 to 3 minims.
Oleum terebinth	0.30 to 2 C. c.	5 to 30 minims.
Oleum tigllii	1 to 10 centigrams	$\frac{1}{4}$ to $\frac{1}{2}$ grains.
Opium (14 $\frac{1}{2}$ morphine)	1 to 10 centigrams	$\frac{1}{6}$ to $\frac{1}{2}$ grains.
Pepsinum purum	1 to 15 Gm	15 grains to $\frac{1}{2}$ ounce.
Pepsinum saccharatum	2 to 30 Gm	30 grains to 1 ounce.
Phosphorus	0.50 to 3 milligrams	$\frac{1}{32}$ to $\frac{1}{8}$ grain.
Physostigmine salicylas	0.50 to 1 milligram	$\frac{1}{32}$ to $\frac{1}{16}$ grain.
Physostigmine sulphas	0.50 to 1 milligram	$\frac{1}{32}$ to $\frac{1}{16}$ grain.
Picrotoxinum	1 to 8 milligrams	$\frac{1}{16}$ to $\frac{1}{4}$ grain.
Pilocarpina (and salts)	1 to 30 milligrams	$\frac{1}{32}$ to $\frac{1}{2}$ grain.
Pil. aloes	1 to 3 pills	1 to 3 pills.
Pil. aloes et asafet.	2 to 5 pills	2 to 5 pills.
Pil. aloes et ferri	1 to 3 pills	1 to 3 pills.
Pil. aloes et mast	1 to 3 pills	1 to 3 pills.
Pil. aloes et myrrhæ	2 to 5 pills	2 to 5 pills.
Pil. antim. comp	1 to 3 pills	1 to 3 pills.
Pil. asafetida	1 to 6 pills	1 to 6 pills.
Pil. cathart. comp	1 to 4 pills	1 to 1 pills.
Pil. ferri comp	2 to 5 pills	2 to 5 pills.
Pil. ferri iodidi	1 to 4 pills	1 to 4 pills.
Pil. galbani comp	1 to 5 pills	1 to 5 pills.
Pil. opii	1 to 2 pills	1 to 2 pills.
Pil. phosphori	1 to 4 pills	1 to 4 pills.
Pil. rhei	2 to 5 pills	2 to 5 pills.
Pil. rhei comp	2 to 5 pills	2 to 5 pills.
Piperinum	5 to 50 centigrams	1 to 8 grains.
Plumbi acetat.	3 to 20 centigrams	$\frac{1}{2}$ to 3 grains.
Plumbi iodidum	3 to 20 centigrams	$\frac{1}{2}$ to 3 grains.
Potassii acetat.	1 to 4 Gm	15 to 60 grains.
Potassii bicarb.	0.50 to 4 Gm	8 to 60 grains.
Potassii bitart.	4 to 8 Gm	1 to 2 grains.
Potassii bromid	0.50 to 4 Gm	8 to 60 grains.
Potassii carb.	0.50 to 2 Gm	8 to 30 grains.
Potassii chloras	0.50 to 2 Gm	8 to 30 grains.
Potassii citras	1 to 4 Gm	15 to 60 grains.
Potassii cyanid	4 to 8 milligrams	$\frac{1}{8}$ to $\frac{1}{2}$ grain.
Potassii et sodii tart.	15 to 30 Gm	$\frac{1}{2}$ to 1 ounce.
Potassii hypophosphis	0.30 to 1 Gm	5 to 15 grains.
Potassii iodid	0.10 to 1 Gm	2 to 15 grains.
Potassii intras	0.50 to 1 Gm	8 to 15 grains.

DOSE TABLE.—Continued.

Remedies.	Dose in metric weights or measures.	Dose in apothecaries' weights or measures
Potassii sulphas.....	4 to 15 Gm.....	1 to 4 drachms
Potassii sulphidum.....	5 to 60 centigrams	1 to 10 grains
Potassii sulphis.....	1 to 2 Gm.....	15 to 30 grains
Potassii tartras.....	5 to 30 Gm.....	1 to 8 drachms
Pulv. antimonalis.....	15 to 60 centigrams	3 to 10 grains
Pulv. aromat.....	0.50 to 2 Gm.....	8 to 30 grains
Pulv. cretæ comp.....	0.50 to 2 Gm.....	8 to 30 grains
Pulv. glycyrrh. comp.....	2 to 4 Gm.....	30 to 60 grains
Pulv. ipecac. et opii.....	0.30 to 1 Gm.....	5 to 15 grains
Pulv. jalapæ comp.....	2 to 4 Gm.....	30 to 60 grains
Pulv. morphinæ comp.....	0.50 to 1 Gm.....	8 to 15 grains
Pulv. rhei comp.....	2 to 4 Gm.....	30 to 60 grains
Quinidina (and salts).....	0.05 to 2 Gm.....	1 to 30 grains
Quinina (and salts).....	0.05 to 2 Gm.....	1 to 30 grains
Quinnæ arsenias.....	1 to 5 centigrams	$\frac{1}{16}$ to 1 grain
Resina copaibæ.....	10 to 60 centigrams	2 to 10 grains
Resina jalapæ.....	10 to 30 centigrams	2 to 5 grains
Resina podophylli.....	8 to 30 milligrams	$\frac{1}{4}$ to $\frac{1}{2}$ grain
Resina scammonii.....	10 to 60 centigrams	2 to 10 grains
Rheum.....	0.10 to 2 Gm.....	2 to 30 grains
Salicinum.....	0.50 to 2 Gm.....	8 to 30 grains
Santonica.....	0.50 to 4 Gm.....	8 to 60 grains
Santonium.....	6 to 30 centigrams	1 to 5 grains
Sapo.....	0.30 to 2 Gm.....	5 to 30 grains
Scammonium.....	0.20 to 1 Gm.....	3 to 15 grains
Senega.....	0.50 to 4 Gm.....	8 to 60 grains
Sodii acetat.....	1 to 4 Gm.....	15 to 60 grains
Sodii arsenas.....	1 to 6 milligrams	$\frac{1}{16}$ to $\frac{1}{8}$ grain
Sodii benzoas.....	0.30 to 1 Gm.....	5 to 15 grains
Sodii bicarb.....	0.50 to 2 Gm.....	8 to 30 grains
Sodii bisulphis.....	0.50 to 2 Gm.....	8 to 30 grains
Sodii boras.....	0.50 to 2 Gm.....	8 to 30 grains
Sodii bromid.....	0.50 to 2 Gm.....	8 to 30 grains
Sodii carb.....	0.50 to 2 Gm.....	8 to 30 grains
Sodii carb. exsicc.....	0.30 to 1 Gm.....	5 to 15 grains
Sodii chloras.....	0.30 to 2 Gm.....	5 to 30 grains
Sodii hypophosphis.....	0.50 to 1 Gm.....	8 to 15 grains
Sodii hypsulphis.....	0.50 to 2 Gm.....	8 to 30 grains
Sodii iodidum.....	0.30 to 1 Gm.....	5 to 15 grains
Sodii phosphas.....	0.10 to 1 Gm.....	2 to 15 grains
Sodii salicylas.....	0.30 to 2 Gm.....	5 to 30 grains
Sodii santoninas.....	15 to 60 centigrams	2 to 10 grains
Sodii sulphas.....	4 to 30 Gm.....	1 to 2 grains
Sodii sulphis.....	0.50 to 2 Gm.....	8 to 30 grains
Spir. æther.....	2 to 4 C. c.....	30 to 60 minims
Spir. æther. nitrosi.....	2 to 8 C. c.....	$\frac{1}{2}$ to 2 fl. drachms
Spir. ammoniæ.....	0.50 to 2 C. c.....	8 to 30 minims
Spir. ammoniæ arom.....	1 to 4 C. c.....	15 to 60 minims
Spir. camphoræ.....	0.50 to 2 C. c.....	8 to 30 minims
Spir. chloroformi.....	1 to 4 C. c.....	15 to 60 minims
Spir. lavend. comp.....	2 to 4 C. c.....	30 to 60 minims
Spir. menth. pip.....	2 to 4 C. c.....	30 to 60 minims
strychnina (and salts).....	1 to 5 milligrams	$\frac{1}{16}$ to $\frac{1}{8}$ grain
Sulphur.....	2 to 15 Gm.....	$\frac{1}{2}$ to 1 drachms
Syrupus acidi hydroiodici.....		
Syrupus albi.....	5 to 15 C. c.....	1 to 4 fl. drachms

DOSE TABLE. — *Continued.*

Remedies.	Dose in metric weights or measures.	Dose in apothecaries' weights or measures.
Syrupus calcii lactophos.	5 to 10 C.c.	1 to 2 fl. drachms.
Syrupus calcis	1 to 2 C.c.	15 to 30 minims.
Syrupus ferri bromidi	1 to 4 C.c.	15 to 60 minims.
Syrupus ferri iodidi	1 to 4 C.c.	15 to 60 minims.
Syrupus ferri oxidi	5 C.c.	1 fl. drachm.
Syrupus ferri hypophosph.	5 C.c.	1 fl. drachm.
Syr. fer. quin et str. phos.	5 C.c.	1 fl. drachm.
Syrupus hypophosphit.	5 C.c.	1 fl. drachm.
Syrupus hypophosph.c.fer	5 C.c.	1 fl. drachm.
Syrupus ipecac.	2 to 15 C.c.	$\frac{1}{2}$ to 4 fl. drachms.
Syrupus kramerie	2 to 15 C.c.	$\frac{1}{2}$ to 4 fl. drachms.
Syrupus lactucarii	5 to 10 C.c.	1 to 3 fl. drachms.
Syrupus rhei	4 to 15 C.c.	1 to 4 fl. drachms.
Syrupus rhei arom.	4 to 15 C.c.	1 to 4 fl. drachms.
Syrupus rose	4 to 8 C.c.	1 to 2 fl. drachms.
Syrupus rubi	4 to 8 C.c.	1 to 2 fl. drachms.
Syrupus sarsap. comp.	4 to 15 C.c.	1 to 4 fl. drachms.
Syrupus scille	2 to 4 C.c.	$\frac{1}{2}$ to 1 fl. drachm.
Syrupus scille comp.	1 to 4 C.c.	15 to 60 minims.
Syrupus senega	4 to 8 C.c.	1 to 2 fl. drachms.
Syrupus senue	5 to 15 C.c.	1 to 4 fl. drachms.
Tinct. acconiti fol.	0.50 to 1 C.c.	8 to 16 minims.
Tinct. acconiti rad.	0.06 to 0.30 C.c.	1 to 5 minims.
Tinct. accon. rad. Fleming's	0.04 to 0.15 C.c.	$\frac{2}{3}$ to 2½ minims.
Tinct. aloes (1880)	2 to 8 C.c.	$\frac{1}{2}$ to 2 fl. drachms.
Tinct. aloes et myrrha	4 to 8 C.c.	1 to 2 fl. drachms.
Tinct. arnica flor.	0.50 to 2 C.c.	8 to 30 minims.
Tinct. arnica rad.	1 to 2 C.c.	15 to 30 minims.
Tinct. asafetide	2 to 4 C.c.	30 to 60 minims.
Tinct. belladonna	0.30 to 1 C.c.	8 to 15 minims.
Tinct. bryonie	1 to 2 C.c.	15 to 30 minims.
Tinct. calendula	1 to 2 C.c.	15 to 30 minims.
Tinct. calumbæ	3 to 15 C.c.	1 to 4 fl. drachms.
Tinct. cannabis ind.	1 to 2 C.c.	15 to 30 minims.
Tinct. cantharid.	0.50 to 1 C.c.	8 to 15 minims.
Tinct. capsici	0.50 to 1 C.c.	8 to 15 minims.
Tinct. catechu comp.	2 to 8 C.c.	$\frac{1}{2}$ to 2 fl. drachms.
Tinct. chirate	1 to 4 C.c.	15 to 60 minims.
Tinct. cimicifuga	2 to 4 C.c.	30 to 60 minims.
Tinct. cinchona	2 to 8 C.c.	$\frac{1}{2}$ to 2 fl. drachms.
Tinct. cinchona comp.	2 to 8 C.c.	$\frac{1}{2}$ to 2 fl. drachms.
Tinct. colchici rad.	0.30 to 1 C.c.	5 to 15 minims.
Tinct. colchici sem.	0.30 to 1 C.c.	5 to 15 minims.
Tinct. conii	0.30 to 2 C.c.	5 to 30 minims.
Tinct. croci	4 to 8 C.c.	1 to 2 fl. drachms.
Tinct. cubebæ	4 to 8 C.c.	1 to 2 fl. drachms.
Tinct. digitalis	0.30 to 1 C.c.	6 to 15 minims.
Tinct. ferri acet.	1 to 2 C.c.	15 to 30 minims.
Tinct. ferri chloridi	1 to 2 C.c.	15 to 30 minims.
Tinct. fer. chloridi ather.	1 to 2 C.c.	15 to 30 minims.
Tinct. ferri pomati	2 to 4 C.c.	20 to 60 minims.
Tinct. gallæ	2 to 8 C.c.	$\frac{1}{2}$ to 2 fl. drachms.
Tinct. gelsemii	0.50 to 1 C.c.	8 to 15 minims.
Tinct. guaiac.	2 to 4 C.c.	30 to 60 minims.
Tinct. guaiaci ammon.	2 to 4 C.c.	30 to 60 minims.
Tinct. hellebori	0.50 to 1 C.c.	10 to 15 minims.

DOSE TABLE.—Continued.

Remedies.	Dose in metric weights or measures.	Dose in apothecaries' weights or measures.
Tinct. humul.....	4 to 10 C. c.....	1 to 2½ fl. drachms.
Tinct. hydrastis.....	2 to 6 C. c.....	30 to 90 minims.
Tinct. hyoscyami fol.....	1 to 4 C. c.....	15 to 60 minims.
Tinct. hyoscyami sem.....	1 to 2 C. c.....	15 to 30 minims.
Tinct. ignatia.....	0.30 to 1 C. c.....	5 to 15 minims.
Tinct. iodi.....	0.30 to 1 C. c.....	5 to 15 minims.
Tinct. ipecac. et opil.....	0.30 to 1 C. c.....	5 to 15 minims.
Tinct. jalapæ.....	2 to 8 C. c.....	½ to 2 fl. drachms.
Tinct. kino.....	2 to 8 C. c.....	½ to 2 fl. drachms.
Tinct. krameria.....	2 to 8 C. c.....	½ to 2 fl. drachms.
Tinct. lavend. comp.....	2 to 8 C. c.....	½ to 2 fl. drachms.
Tinct. lobelia.....	1 to 3 C. c.....	15 to 45 minims.
Tinct. lupulini.....	2 to 8 C. c.....	½ to 2 fl. drachms.
Tinct. matico.....	2 to 8 C. c.....	½ to 2 fl. drachms.
Tinct. moschi.....	1 to 4 C. c.....	15 to 60 minims.
Tinct. nuc. vomice.....	0.50 to 3 C. c.....	8 to 45 minims.
Tinct. opil.....	0.30 to 1 C. c.....	5 to 15 minims.
Tinct. opii camph.....	0.50 to 5 C. c.....	8 to 75 minims.
Tinct. phytolacæ.....	0.50 to 4 C. c.....	8 to 60 minims.
Tinct. physostigmatis.....	0.30 to 1 C. c.....	5 to 15 minims.
Tinct. pyrethri.....	0.50 to 2 C. c.....	8 to 30 minims.
Tinct. quassia.....	2 to 8 C. c.....	½ to 2 fl. drachms.
Tinct. rhei.....	4 to 30 C. c.....	1 to 8 fl. drachms.
Tinct. rhei arom.....	2 to 5 C. c.....	30 to 75 minims.
Tinct. rhei dulc.....	4 to 15 C. c.....	1 to 4 fl. drachms.
Tinct. sanguinaria.....	1 to 4 C. c.....	15 to 60 minims.
Tinct. scilla.....	0.50 to 4 C. c.....	8 to 60 minims.
Tinct. serpentaria.....	2 to 8 C. c.....	½ to 2 fl. drachms.
Tinct. stramon. fol.....	0.50 to 1 C. c.....	8 to 15 minims.
Tinct. stramon sem.....	0.30 to 1 C. c.....	6 to 15 minims.
Tinct. sumbul.....	0.50 to 2 C. c.....	8 to 30 minims.
Tinct. valer.....	2 to 8 C. c.....	½ to 2 fl. drachms.
Tinct. valer. ammon.....	2 to 8 C. c.....	½ to 2 fl. drachms.
Tinct. veratr. vir.....	0.20 to 0.60 C. c.....	3 to 10 minims.
Tinct. zingiberis.....	1 to 4 C. c.....	15 to 60 minims.
Tritur. elaterini.....	8 to 30 milligrams	¼ to ½ grain.
Veratrina.....	1 to 6 milligrams	½ to 1 grain.
Vin. aloes.....	4 to 8 C. c.....	1 to 2 fl. drachms.
Vin. antim. { expect et. alt. { emet.	0.06 to 0.50 C. c..... 2 to 5 C. c.....	1 to 8 minims. 30 to 75 minims.
Vin. colch. rad.....	0.50 to 3 C. c.....	8 to 45 minims.
Vin. colch. sem.....	0.30 to 2 C. c.....	5 to 30 minims.
Vin. ergotæ.....	4 to 12 C. c.....	1 to 3 fl. drachms.
Vin. ferri amar.....	5 C. c.....	1 fl. drachm.
Vin. ferri citrat.....	5 C. c.....	1 fl. drachm.
Vin. ipecac. { expect..... { emet.....	0.30 to 1 C. c..... 10 to 25 C. c.....	5 to 15 minims. 3 to 6 fl. drachms.
Vin. opil.....	0.30 to 1 C. c.....	5 to 15 minims.
Vin. rhei.....	4 to 8 C. c.....	1 to 2 fl. drachms.
Zinci acet.....	1 to 12 centigrams	1 to 2 grains.
Zinci bromid.....	3 to 10 centigrams	½ to 2 grains.
Zinci iudid.....	3 to 15 centigrams	½ to 3 grains.
Zinci oxid.....	6 to 60 centigrams	1 to 10 grains.
Zinci phosphid.....	5 to 10 milligrams	⅓ to ½ grain.
Zinci sulphas emet.....	1 to 2 Gm.....	15 to 30 grains.
Zinci valerianas.....	6 to 40 centigrams	1 to 6 grains.

TABLE FOR CALCULATING THE PERIOD OF
UTERO-GESTATION.

NINE CALENDAR MONTHS.			TEN LUNAR MONTHS.	
From	To	Days.	To	Days.
January 1....	September 30...	273	October 7.....	280
February 1....	October 31.....	273	November 7.....	280
March 1.....	November 30....	275	December 5.....	280
April 1.....	December 31....	275	January 5.....	280
May 1.....	January 31.....	276	February 4.....	280
June 1.....	February 28....	273	March 7.....	280
July 1.....	March 31.....	274	April 6.....	280
August 1.....	April 30.....	273	May 7.....	280
September 1..	May 31.....	273	June 7.....	280
October 1.....	June 30.....	273	July 7.....	280
November 1..	July 31.....	273	August 7.....	280
December 1..	August 31.....	274	September 6....	280

The above Obstetric "Ready Reckoner" consists of two columns, one of Calendar, the other of Lunar months, and may be read as follows:—a Patient has ceased to menstruate on the 1st of July; her confinement may be expected at soonest about 31st of March (*the end of nine Calendar months*); or at latest, on the 6th of April (*the end of ten Lunar months*). Another has ceased to menstruate on the 20th January; her confinement may be expected on the 30th September, plus 20 days (*the end of nine Calendar months*); at soonest, on the 7th October, plus 20 days (*the end of ten Lunar months*); at latest

A LIST OF THE MOST IMPORTANT NEW REMEDIES,
AND A FEW OLD ONES WHICH HAVE RECENTLY
BEEN REVIVED.

BY HENRY MORRIS, M.D.

ABRUS PRÆCATORIUS—(*Aletris eye*) useful in intractable forms of eye disease, as granular conjunctivitis, inveterate pinnitis, corneal ulcers, etc., but requires caution in its use. It sets up a severe conjunctivitis, which may be purulent, and which on subsiding leaves the conjunctiva in a healthier condition. Take 32 well pulverized seed triturate and then macerate them for 24 hours in 500 Gm. of cold water, then add 500 Gm. of hot water, and filter the liquid immediately on cooling. Bathe the eye 3 times daily, for 3 days, when sufficient inflammation will be induced.

ADONIDINE (recently obtained by Cervello from *Adonis vernalis*) similar in its effects to digitalin, but much more energetic, as gr. 2 is sufficient to arrest the cardiac action of a frog. It appears to have no stimulative action.

AMBROSIA TRIFIDA (*Ragweed*) recommended as a styptic in various hemorrhages.

ARECA CATECHU (*Betelnut*). The seed are recommended as anthelmintic.

ASPIDOSPERMA QUEBRACHO (*Quebracho bark*) acts principally upon the respiratory motor apparatus, rendering the breathing deeper and slower. It is very beneficial in various forms of dyspnoea, as that occurring in emphysema, asthma, bronchial catarrh, etc., and has also been given as an astringent in diarrhoea. It does not prove of as much benefit in asthma due to valvular disease, as in other forms of the affection. Dose of the fluid extract ℥j xx-45.

BERBERIS AQUIFOLIUM (*Oregon-grape root*) is recommended as an alterative and tonic in scrofula, secondary and tertiary syphilis, and chronic cutaneous affections, especially of the squamous variety, as psoriasis, etc. Dose of the fluid extract, ℥jxx-45.

BETEL NUT, see *Areca Catechu*.

CAFFEINA (official) causes cerebral excitement, wakefulness, and hallucinations; at first increases, but afterward depresses the cardiac action and arterial pressure; and produces diuresis in from 12 to 24 hours. It is used in nervous headache (gr. j-ij) in cardiac dropsy (gr. iv gradually increased to gr. xv or even gr. xxx, thrice daily) and to antagonize opium narcosis. It is said to be of great benefit in cases of fatty heart when digitalis does not act or is contra-indicated. It is of no use in albuminuria, when not dependent on cardiac disease.

CASCARA AMARGA (the bark of) recommended in secondary syphilis, as a tonic and alterative, in doses of ℥jxx-45 of the fluid extract.

CASCARA SAGRADA, see *Rhamnus Purshiana*.

CEREUS BONPLANDII and **CEREUS GRANDIFLORIS** have been recommended in palpitation of the heart, especially functional and cardiac dyspnoea. The fluid extract of either may be given in doses of grt x-xxx.

CHAULMUGRA OIL, see *Camponotus clogata*.

CHRYSAROBINUM (*Chrysarobin* official) often improperly called *Chrysophanic acid*, causes, when applied externally, especially about the head and face, irritation and inflammation of the skin (sometimes, a papular eruption) which is not always limited to the seat of application. It is an excellent application in psoriasis and the various forms of ringworm, but should not be used over a large surface at one time. The official ointment contains 10 per cent. of the drug.

CONVALLARIA MAJALIS (*Lily of the Valley*) is said to stimulate the central cardiac inhibitory apparatus, thus slowing the heart and finally arresting it in systole; the respiration is at first increased in frequency, then retarded and deepened; the blood pressure is increased by stimulation of the vaso-motor nerves and contraction of the peripheral vessels; the urine is increased in quantity and reflex action is lowered.

It is recommended to make the cardiac contractions stronger, slower, and more regular; to deepen and render easier the respiration in dyspnea, whether cardiac or emphysematous; as a diuretic, especially in cardiac dropsies; and to lower reflex action. It should not be used in gastric or intestinal catarrh; in acute affections of the liver, spleen, or kidneys; in inflammation of the uterus or ovaries; in pregnancy; in hemorrhoids; or in fatty degeneration of the heart. The *fluid extract* may be given in doses of gtt. xv-xx every 3 or 4 hours, until the desired effects are produced.

CORN ERGOT or **CORN-SMUT**, see *Ustilago*.

CORN SILK, see *Stigmata Maydis*.

COTO BARK (containing *Cotoneum*, *Paracotoneum*, etc.) is highly recommended in various forms of obstinate diarrhoea, cholera infantum, etc. In the diarrhoea following typhoid fever or accompanying phthisis it is often curative, or at least palliative in its effects, and diminishes also the colliquative sweating of the latter disease. It occasionally causes nausea, vomiting and epigastric pain, from the presence of an aromatic resin which it contains; but these effects may be obviated by the use of *cotoneum* (dose gr. $\frac{1}{4}$ iss; t. d.) or *paracotoneum* (dose gr. iv-v; t. d.). Dose of the *tincture* 10 per cent. of the bark) f5j-ij; of the *fluid extract* gtt. i-x or xv.

DUBOISIA MYOPÆOIDES (the leaves of): contains an alkaloid *duboisin* (or *duboisine*). Its effects are similar to those of atropine, from which it differs in causing more rapid dilatation of the pupil, total paralysis of accommodation, and in being less irritating. It may be used topically or by subcutaneous injection (gr. $\frac{1}{16}$).

ERGOTININE is said to be the true active principle of ergot, and as such is said to be used successfully in cases to which ergot is applicable.

ERIODYCTION CALIFORNICUM (*Yerba Santa*) is recommended to disguise the taste of quinine, and as an expectorant in chronic bronchitis, etc. Dose of the fluid extract, f5ss-j.

EUCALYPTOL obtained from the essential oil of *Eucalyptus globulus* is useful in bronchitis and pertussis, and has lately been strongly recommended as an inhalation in infectious diseases, as typhoid fever, influenza, diphtheria, etc. In the latter affection, it is said, the false membrane is easily coughed up, and sometimes casts of the bronchi even, if the patient is kept in a moist atmosphere, which is charged with the vapor of eucalyptol by means of an atomizer. Inhalations are also recommended in croup, bronchitis, asthma, etc.

EUONYMUS ATROPURPUREUS (*Wahoo*) is a powerful hepatic stimulant, and hydragogue cathartic. It is said to act through the solar and hypogastric plexus of the sympathetic. It acts upon the kidneys by increasing the blood pressure, and in over-doses causes albuminuria. It is useful in torpid conditions of the liver and bowels. Dose of the extract (official) gr. iij-v.

FRANCISCEA UNIFLORA (*Mimosa*) has been recommended for chronic rheumatism, chronic joint pains, syphilis, scrofula, etc. Dose of the fluid extract, Mv-f3j, several times a day.

FRANKENIA GRANDIFOLIA (*Yerba Buena*) is recommended as a local application in all chronic catarrhs of mucous membranes, especially in chronic nasal catarrh, and as an injection in gonorrhoea and leucorrhoea, in the strength of *fluid extract*, f3ij to water, f3iv. It has been used internally in doses of ℥x-xx.

FUCHSINE is said to be of service in chronic albuminuria with oedema. *Dose*, gr. j-ij, thrice daily.

GRINDELIA (officinal;—the leaves and flowering tops of the *G. obusta*) slows the cardiac and respiratory movements, and in large doses possesses decided hypnotic effects, during which reflex action, motion, and sensation, are depressed. It is highly recommended in asthma, especially in the spasmodic variety, and will frequently check hay asthma and hay fever. It is also recommended in chronic pyelitis or cystitis, etc. *Dose of the fluid extract*, ℥x-15j.

GRINDELIA SQUARROSA is recommended in chronic malaria, and in enlargement of the spleen, etc., in which affections it is said to prove very beneficial. *Dose of the fluid extract*, ℥x-15ss.

GUACHAMACA produces effects similar to those caused by curare, but differs in not disturbing the respiration, circulation nor cardiac action. The hypodermic injection of gr. ½, given to a young man suffering with cramps, produced no effect for three-quarters of an hour, when the patient suddenly fell into a deep sleep, lasting for 4 hours. It is believed to be a good motor-depressant and hypnotic.

GYNOCARDIA ODORATA. The seed of this plant contain an oil known as *Chaulmugra oil*, which has been highly recommended in leprosy, especially in the macular and anæsthetic varieties. It has also been used in syphilis, rheumatism, scrofula, phthisis, etc., both internally and as a local application. *Dose of the oil*, gr. v, of the powdered seed, gr. v, in pill.

HOANG-NAN, see *Strychnos gaultheriana*.

ICHTHYOL (derived by distillation from a bituminous mineral found in small quantities in the Tyrol) contains 10 per cent. of sulphur, and is recommended for eczema, applied locally. It is also strongly recommended as a local application in acute and chronic rheumatism, at the same time that the salicylic acid treatment is being given internally. It has been used locally in gout. Occasionally it causes a superficial inflammation of the skin.

JAMAICA DOGWOOD, see *Piscidia erythrina*.

JEQUIRITY, see *Abrus precatorius*.

KAIRINA, in doses of gr. v-viii every hour, is recommended as the "most certain, most powerful, and most rapid" antipyretic we possess, and is given in all conditions of hyperpyrexia.

LILY OF THE VALLEY, see *Convallaria Majalis*.

LIPPIA MEXICANA has been recommended as a sedative and expectorant, in various bronchial affections.

MANACA, see *Franciscia uniflora*.

MANGIFERA INDICA (*Mango*) has been used in the treatment of diarrhoea, dysentery, inflammation of the uterus, etc., and as an injection in gonorrhoea, leucorrhoea. It is said to check profuse menstruation and also the hemorrhage which follows a miscarriage. *Dose of the fluid extract*, ℥x-15i.

MENTHOL (*Peppermint camphor*) has antiseptic properties similar to thymol. It has been principally used as a local application in the treatment of neuralgia of various kinds, as neuralgia of the 5th nerve, cervico-brachial, intercostal, and lumbal-abdominal neuralgia, or in sciatica, in which affections it often gives wonderful relief. A solution of 5 to alcohol f3ss should be painted over the painful part. It has also been used in odontalgia, applied to the cavity of the carious tooth.

MUSCARINA (an active principle obtained from *Agaricus muscaria*) causes profuse perspiration and salivation, and increases the hepatic, pancreatic and intestinal secretions. It is a cardiac motor-depressant, and produces dyspnea by contracting the pulmonary vessels. *Dose*, gr. $\frac{1}{2}$ ss, hypodermically. It has been recommended in the sweating of phthisis, in the dose of $\mathcal{M}$ of a "one per cent. solution of liquid extract."

MYRTUS CHEKAN has been recommended as an expectorant in chronic "winter cough," bronchitis, etc. It has also been used in catarrh of the bladder "and other affections of the mucous membranes." *Dose* of the fluid extract, f5j iv.

NITROGLYCERINUM (*Nitroglycerin*—*Trinitroglycerin*—*Glonoïn*) is used in solution of the strength of 1 per cent., the dose of which is, gtt. j-vj. Three minutes after taking gtt. j, it is said, the cardiac action is increased and the sounds much stronger, the pulse rises from 8 to 10 beats per minute, and the arterial tension is lowered. These effects last three-quarters of an hour. When gtt. ij are taken, the effects are headache and flushed face, in addition to the above. Photophobia and painful throbbing of the head are produced by gtt. vj. The urine is said to be unaffected. In toxic doses, death is caused by paralysis of the respiratory muscles. This remedy is highly recommended in angina pectoris from any cause. It seems to be of more benefit in preventing the paroxysms, than in checking them when present. It is recommended also in the commencement of an attack of asthma (especially if due to emphysema) and in neuralgia.

NICKEL ; SALTS OF. These are said to be excellent and powerful disinfectants. It has been found that $\frac{1}{10}$ of a 1 per cent. solution of the chloride, will kill bacteria termo.

OREGON-GRAPE ROOT see *Berberis aquifolium*.

PARACOTO BARK and PARACOTOIN, see *Coto Bark*.

PARALDEHYDE (a polymeric modification of aldehyde) acts first on the hemispheres, then on the medulla and cord. In large doses it paralyzes the respiratory centre. It diminishes reflex action. It is a hypnotic, producing a tranquil sleep, without unpleasant after effects. During sleep the cardiac action is diminished in frequency and increased in force. It is recommended in the insomnia of various mental disorders, and also in sleeplessness from other causes, as from severe mental occupation, etc. *Dose*, 3j-4j.

PELLETERINA (or *Pelchidlin*) the active principle of the Punica granatum or pomegranate, is said to be one of the most efficient tonic-cides we possess. *Dose*, gr. iv-xi.

PILOCARPINA (the active principle of *Pilocarpus pennatifolius* or *Jaborandi*) is similar in its effects to Jaborandi, and may be used for the same purposes, in doses of, gr. $\frac{1}{2}$ ss, hypodermically.

PISCIDIA ERYTHRINA (*Jamaica Dogwood*) is said to be a narcotic, to diminish reflex action; to dilate the pupil; to cause salivation and diaphoresis; to decrease the frequency of the pulse, and elevate arterial tension, and finally, to cause death by asphyxia. It is recommended as a substitute for opium to produce sleep and allay pain. *Dose* of the fluid extract, f3ss-j.

QUEBRACHO, see *Aspidosperma quebracho*.

RAGWEED, see *Ambrosia trifida*.

RESORCIN made by fusing certain gums with potassium hydrate, is a powerful antiseptic. Given internally, it at first increases, then depresses the circulation, causes increased secretion from the salivary and sudoriparous glands, and depresses the temperature. It has been used successfully as a topical application to ulcers and chancre, and as an injection in purulent vaginitis. Internally, it is highly recom-

mended for puerperal fever, erysipelas (in which disease it has been used hypodermically), anthrax, intermittent fever, and in hyperpyrexias in doses of gr. xx-3j.

RHAMNUS PURSHIANA (*Cascara Sagrada*—*Chiltern Bark*) is a good and efficient cathartic, highly recommended in habitual constipation. *Dose of the fluid extract*, ℥x-℥ss.

SAPO VIRIDIS (*Green Soap*) is used in various skin diseases, especially of the scaly variety. When properly applied, it will generally remove infiltrated patches of eczema rubrum. It is much used to remove scales and crusts in psoriasis, seborrhoea, etc.; and will cure tinea versicolor if rubbed in once or twice a day thoroughly, and allowed to remain. The part should not be washed until the fifth day, during which time the treatment should be kept up.

SARRACENIA FLAVA (*Trumpet plant*—*pitcher-plant*) has been used in dyspepsia, acute, chronic and infantile diarrhoea and chronic nasal catarrh, and as an injection in leucorrhoea and gonorrhoea. *Dose of the fluid extract*, ℥x-℥ss.

SODII NITRIS (*Sodium Nitrate*) is recommended as being as active, and reliable in angina pectoris, as either amyl nitrite or nitroglycerin, and having the advantage of not causing headache, giddiness, or other untoward symptoms, when given in medicinal doses. *Dose*, gr. ij-v. in water.

STIGMATA MAYDIS (*Corn Silk*) is recommended as a diuretic in those cases of heart disease and albuminuria which require such remedies. It is said to be a valuable agent in the treatment of cystitis, etc. *Dose of the fluid extract*, ℥ss.

STRYCHNOS GAULTHERIANA (*Hoang-nan*), in common with other members of the same family, is a tetanizing agent. It is recommended in leprosy, syphilis, scrofula, various forms of ulcers, etc. It is given in pill, combined with alum and realgar. *Dose*, gr. 1/4 ss.

TRICHLOPHENOL is recommended as an antiseptic twenty-five times more powerful than carbolic acid. It does not irritate the tissues, and when applied in substance to wounds or ulcers, it has but a mild caustic effect. It is said to be a powerful deodorizer, and has been recommended as the best antiseptic for hospital gangrene, or any extensive gangrene whatever. Topically, it is said to have been used with beneficial results to phagedenic chancres (2.5 per cent. solution in glycerin).

THYMOL (a crystalline substance found in the volatile oil of the *Thymus vulgaris*) is a powerful antiseptic, analogous to carbolic acid in its effects. Internally it is an antipyretic (not, however, as powerful as salicylic acid) and causes diaphoresis, and sometimes diarrhoea, nausea and vomiting. It is recommended in inflammations and ulcerations of the mouth, and as an inhalation to diminish the expectoration of phthisis. Crystallized thymol is recommended internally in smallpox, in doses of gr. ss, and is used in solution as a gargle in diphtheria.

USTILAGO (*Corn Smut*—*Corn Ergot*) resembles ergot in its effects and uses, and may be given "in similar preparations and doses."

VIBURNUM (the bark of *Viburnum Prunifolium*—*Sloe* or *Black Haw*) is said to be a sedative and tonic to the nervous centres governing the uterine functions. It is recommended as an anti-abortion, and in all cases of uterine engorgement, ovarian irritation, etc. In irregular menstruation with dysmenorrhoea and excessive loss of blood, it appears to be of very great service. The *fluid extract* is officinal. *Dose*, ℥ss-i.

WAHOO, see *Euonymus atropurpureus*.

YERBA REUMA, see *Frankenia grandifolia*.

YERBA SANTA, see *Erioduction Californicum*.

SYLVESTER'S METHOD

FOR PRODUCING ARTIFICIAL RESPIRATION.

From Dulles' Accidents and Emergencies.

Lay the person to be treated upon the back, then kneeling behind the head, seize each arm near the elbow and move them round horizontally, first away from the body, then over the head till they meet above it (see Fig. 1),

when they should be given a steady pull for a few seconds. This fills the lungs with air by drawing the ribs up and so enlarging the cavity of the chest.




FIG. 1.

The second manœuvre consists in returning the arms to their former position alongside the chest (see Fig. 2) and making strong pressure against the lower ribs, so as to drive the air out of the chest and effect expiration. This should occupy but a second of time, and be repeated about sixteen times per minute for some time,

and not abandoned until it has been certainly ascertained that the heart has ceased beating.




FIG. 2.

AL

E

T

T

T

T

T

T

T

T

T

T

T

T

T

T

T

T

T

T

T

T

T


T

T

T

T

T



VISITING LIST.

25 Patients per Week.

1885.

JANUARY.

Su. M. Tu. W. Th. F. S. A.

Mr Fuller, and others
Newburyport

cc
c. 20th pd

JANUARY

SU. M. TU. W. TH. F. S.

1 2 3 4 5 6 7 8 9 10 11 12

AMOUNT.

\$ 40

Mr Fuller

c c

Mr J. C. Ross 200 Lumber
for Bullen.

@ 200 ft 6 5/8 ft.

JANUARY.

Su.	M.	Tu.	W.	Th.	F.	S.	Am.
11	12	13	14	15	16	17	

E. Rees Spring (mill. Delco

C 1/4 #2.

JANUARY

Su. M. Tu. W. Th. F. S.

1 2 3 4 5 6 7 8 9 10 11 12

AMOUNT

1/2 Mrs. Lewis

Mrs. [unclear] 75/100

1/2 Pd

JANUARY

Su. M. Tu. W. Th. F. S.

6 27 28 29 30 31

See notes, page 100, letter A5

FEBRUARY

Su.	M.	Tu.	W.	Th	F	S	Month No.
1	2	3	4	5	6	7	

FEBRUARY

Su. M. Tu. W. Th. F. S.

8 9 10 11 12 13 14

FEBRUARY.

Sat.	M.	Tu.	W.	Th.	F.	Sat.	AMOUNT
1	2	3	4	5	6	7	

CALENDAR

Su. M. Tu. W. Th. F. S.
12 13 14 15 16 17 18

MARCH

Su. M. Tu. W. Th. F. S.

ADOLEN

MARCH.

Su.	M.	Tu.	W.	Th.	F.	S.
8	9	10	11	12	13	14

MARCH.

Su.	M.	Tu.	W.	Th.	F.	S.	AMOUNT.
15	16	17	18	19	20	21	

MARCH

Su.	M.	Tu.	W.	Th.	F.	S.
1	2	3	4	5	6	7

MAR & APRIL

Su. M. Tu. W. Th. F. S.

AUGUST

1 2 3 4 5 6 7

APRIL

Su.	M.	Tu.	W.	Th.	F.	S.	Av.
1	6	7	8	9	10	11	

W. H. Lee

Q

En

APRIL

Su.	M.	Tu.	W.	Th.	F.	S.	AMOUNT
12	13	14	15	16	17	18	

Mayes Small

C. C. C. \$30

APRIL.

Su. M. Tu. W. Th. F. S.

10 11 12 13 14 15

APRIL & MAY.

Sa. M. Tu. W. Th. F. S.

26 27 28 29 30 1 2

AMOUNT

MAY.

Su. M. Tu. W. Th. F. S.

1 2 3 4 5 6 7 8 9

Rev. Mr. Gilbert Cochran & c.

MAY.

Su. M. Tu. W. Th. F. S.

10 11 12 13 14 15 16

AMOUNT

Mr E. Albert	a c	\$ 2.75
Miss Small	Thurs 3.00	13.11 8.30
Mr Lewis Howell	2	\$ 5.10

MAY

Su.	M.	Tu.	W.	Th.	F.	S.	Am.
17	18	19	20	21	22	23	

MAY.

Su. M. Tu. W. Th. F. S.

24 25 26 27 28 29 30

AMOUNT.

MAY & JUNE

Su. M. Tu. W. Th. F. S.

31 1 2 3 4 5 6

JUNE.

Su. M. Tu. W. Th. F. S.

AMOUNT

7 8 9 10 11 12

Miss Hilbert	825	Recess	1	Charles	24
Salome P. Hudson			4		
Miss Brader	121	Laurel		1	21
Mr		at P. Hudson		1	

JUNE

Su. M. Tu. W. Th. F. S. A.
 11 12 13 14 15 16 17 18 19 20

Mrs. T. T. T. T. T.

5 6 7 8 9 10 11 12

Mrs. T. T. T. T. T.

13

JUNE

S. M. Tu. W. Th. F.

AGENTS

21 22 23 24 25 26

Madison Philadelphia

/ / / / / / /

Mrs. M. M. M. M. M. M. M.

2

Mr. G. G. G. G. G. G. G.

10

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

JUNE

Su. M. Tu. W. Th. F. S. A.

Mr.	at Putnam	1	to be at H. A. M.
Mrs. Brader		.	@
Mrs. Threlkeld		1	1 / 1 1 /

JULY.

Su. M. Tu. W. Th. F. S.

AMOUNT

1 2 3 4

Madame Toulouan / / / / /

Mr. Morris, West. Phila. " \$10.00.

Mrs. V. Paul " \$10.00
10

JULY

Su. M. Tu. W. Th. F. S.

5 6 7 8 9 10 11

Aug

Madame T. M. L. 1 1 1 1 2 * X /
 New Hall 2
 Mr. J. L. H. L. 2

JULY.

Su. M. Tu. W. Th. F. S.
12 13 14 15 16 17 18

AMOUNT.

1 st The Trade are	1	1	1	1	1	1	
2 nd Mr. Moore	c						85 th and
3 rd S. Morris, & Co.							91 st and
4 th Mr. Gilbert.							and
5 th S. & Co. of New York							@

JULY.

Su. M. Tu. W. Th. F. S.

19 20 21 22 23 24 25

Mrs. Buckley
Mrs. Tinsdale

0 C 325 Savers
/ / / / / / Camp

JULY & AUG.

Su. M. Tu. W. Th. F. S.

26 27 28 29 30 31 1

AMOUNT.

July 1 - 1881

AUGUST.

Su.	M.	Tu.	W.	Th.	F.	S.	V.
2	3	4	5	6	7	8	

AUGUST.

Su. M. Tu. W. Th. F. S.

9 10 11 12 13 14 15

August

AUGUST.

Su.	M.	Tu.	W.	Th.	F.	S.
16	17	18	19	20	21	22

AUGUST

Su.	M.	Tu.	W.	Th.	F.	S.	AMOUNT
23	24	25	26	27	28	29	

AUG. & SEPT.

Su.	M.	Tu.	W.	Th.	F.	S.	Am.
30	31	1	2	3	4	5	

SEPTEMBER

Su.	M.	Tu.	W.	Th.	F.	S.	AMOUNT
6	7	8	9	10	11	12	

Dr. F. W. C.
m.

Received of

SEPTEMBER

Su.	M.	Tu.	W.	Th.	F.	S.	AN
13	14	15	16	17	18	19	

Miss. T. Ballan

or
1

SEPTEMBER.

Su. M. Tu. W. Th. F. S.

29 21 22 23 24 25 26

AMOUNT

SEPT. & OCT

Su. M. Tu. W. Th. F. S.

31 28 29 30 1 2

T.

OCTOBER.

Sa. M. Tu. W. Th. F. S.

1 2 3 4 5 6 7 8 9 10

AMOUNT

Dr Given

in 2000

100

1

100

100

11 12 13 14 15 16 17

11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100										
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100

OCTOBER.

Su.	M.	Tu.	W.	Th.	F.	S.	AMOUNT.
18	19	20	21	22	23	24	

William Wallace

c

OCTOBER.

Su.	M.	Tu	W.	Th.	F.	S.
25	26	27	28	29	30	31

William Wallace

1871

NOVEMBER.

Su. M. Tu. W. Th. F. S.

AMOUNT

1 2 3 4 5 6 7

H. Wallace

c c 1 2

c 8.272

NOVEMBER.

Su.	M.	Tu.	W.	Th.	F.	S.	Av.
8	9	10	11	12	13	14	

[Faint handwritten notes, possibly "Nov 1st" and "Nov 2nd", are visible in the top left corner.]
[Faint handwritten notes, possibly "Nov 3rd" and "Nov 4th", are visible in the top right corner.]
[Faint handwritten notes, possibly "Nov 5th" and "Nov 6th", are visible in the middle right area.]
[Faint handwritten notes, possibly "Nov 7th" and "Nov 8th", are visible in the bottom right area.]

NOVEMBER.

Su. M. Tu. W. Th. F. S. AMO. N.

10 16 17 18 19 20 21

C. Collorey

c c 2 10 11 12

c 1 10 11 12

c 1 10 11 12

1 1

2 1

d

NOVEMBER

Su. M. Tu. W. Th. F. S.

22 23 24 25 26 27 28

Am

Handwritten entries in a ledger format:
 4th Nov.
 5th Nov.
 6th Nov.
 7th Nov.
 8th Nov.
 9th Nov.
 10th Nov.
 11th Nov.
 12th Nov.
 13th Nov.
 14th Nov.
 15th Nov.
 16th Nov.
 17th Nov.
 18th Nov.
 19th Nov.
 20th Nov.
 21st Nov.
 22nd Nov.
 23rd Nov.
 24th Nov.
 25th Nov.
 26th Nov.
 27th Nov.
 28th Nov.
 29th Nov.
 30th Nov.
 1st Dec.
 2nd Dec.
 3rd Dec.
 4th Dec.
 5th Dec.
 6th Dec.
 7th Dec.
 8th Dec.
 9th Dec.
 10th Dec.
 11th Dec.
 12th Dec.
 13th Dec.
 14th Dec.
 15th Dec.
 16th Dec.
 17th Dec.
 18th Dec.
 19th Dec.
 20th Dec.
 21st Dec.
 22nd Dec.
 23rd Dec.
 24th Dec.
 25th Dec.
 26th Dec.
 27th Dec.
 28th Dec.
 29th Dec.
 30th Dec.
 31st Dec.

Vertical handwritten note on the right margin:
 1st Dec. 1830

NOV. & DEC.

Su.	M.	Tu.	W.	Th.	F.	S.	AMOUNT.
20	30	1	2	3	4	5	

Worcester

1

Price

1

25.00

C

\$

Worcester H. Y. Co.

C

\$ 1.00

Price

2

DECEMBER

SU.	M.	TU.	W.	TH.	F.	S.	W.
6	7	8	9	10	11	12	

DECEMBER.

Su. M. Tu. W. Th. F. S.

AMOUNT

13 14 15 16 17 18 19

DECEMBER

Su.	M.	Tu.	W.	Th.	F.	S.
20	21	22	23	24	25	26

1	1					
	1					
	1					

DECEMBER.

Sa. M. Tu. W. Th. F. S.

AMOUNT

27 28 29 30 31

MEMORANDA.

JANUARY.

Carried over. 3000.00
Income

\$ 400.00

428

1st Rent - Convent

\$ 20.00

2nd Rent

10.00

7th College cheque

333.14

25th Dean Brothers & Co

31.60

21st Rent

10.00

7th. Mr. Cunningham

20.00

College cheque

333.00

March 20. John E. Rees

20.00

Mr. King. (Jan 12th)

5.00

April 8. Convent. 1000.00

5.00

10th Dean & Co

8.00

Mr. G. Webb

25.00

Mr. Simon

5.00

College cheque (May 23rd)

743.00

Dean Brothers & Co

10.00

24th Dean. (July 7th)

10.00

7/10 Dean Brothers & Co

25.00

Lippincott & Co

3.00

7/11. Convent. 1000.00

5.00

7/19 Fuller cheque

400.00

7/30 Mayor Smith

30.00

Convent. 1000.00

5.00

College 7/11 - 1000.00

50.00

Rent 24th & 25th

5.00

2200.00

MEMORANDA.

JANUARY.

Cash disburse

2nd. To Kresling's Book bound	
Supping	5.00
" Hurling hat-	1.00
8. Nansen	100.00
" Club	4.00
10. Anna Coleman	8.00
" Deha paper	4.00
" Book etc	3.00
15. Therman	35.00
20. Anna L.	4.50
15. Old. & new	35.00
20. Board. & food	10.00
" Bottle	1.00
" Bottle	10.00
20. New York	6.00
" Bottle	5.00
20th March 18th. Expenses Hugen	3.00
April 8th. Trip Home	38.00
May 14th. Hugen	10.00
May 20th. Hugen	2.00
" 12th. Hugen	5.00
June 20th. Hugen	5.00

MEMORANDA

~~FEBRUARY~~

I came

October. 2. near Hull

\$

15th 52

" 4. Hays Feb 104.

10. 67

" 13 A. A. Borevic

10 m

b) $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$, $r = \sqrt{x^2 + y^2 + z^2}$

100

Sept. Hays T. C. C. C. C. C.

20

Nov 4 U. S. Dept. of Agriculture

4.

533.4

Nov. 6. Crossed with the Hollister.

100

1871

215

18. Mr. [illegible] 1861

22

" 21. Hany: 5: - 84.

10

" fi 1 A . . . de

2.

" 1891 1 21

5

1892

5

3. Hilary.

10 5-

66

6. $\frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}$ 25-

28

1 page. Elements

31

5.

MEMORANDA.

FEBRUARY.

Oct.	Expenditure	\$
	W. - books	1. 00
12	New York	7. 00
13	Edm. 86 8.50	13. 00
		2. 00
14		34. 15
"		2. 00
		10. 00
		12. 00
		10. 00
		1. 00
		12. 00
		58. 00
		7. 00
23.	(May, Nov Aug. Sept. Oct)	15. 00
Nov 5th	Munroe &	50. 00
"	" "	10. 00
" 9	B. ...	8. 00
" "		23. 00
"		12. 00
"		10. 00
"		12. 00
in		10. 00
		

MEMORANDA.

MARCH.

MEMORANDA.

MARCH.

MEMORANDA

APRIL.

MEMORANDA.

APRIL.

MEMORANDA.

MAY.

MEMORANDA.

MAY.

MEMORANDA.

JUNE.

MEMORANDA.

JUNE.

MEMORANDA.

JULY.

received sent 66

7 t. 11. 1/2	Smith	8	30
4 1/2	Fuller.	10d.	Commonwealth No. 4.
10	14 Dec.	11	10 Pd.
10	11 Dec.	11	10 Pd.
10	11 Dec.	11	10 Pd.
10	11 Dec.	11	10 Pd.

MEMORANDA.

JULY

From C. C. C. C. C.

Morris W. Phil	Pd. 16/11/8	\$ 10 00
Alf E Morris Chestnut Hill	19/10/86	15 00
Oct 14th Mrs Gunnison Erie		<u>20 00</u>
Hon J L Thibodeau Oct 27.		200 00
William Wallace		
E. C.		10 00
... ..		10 00
... ..		\$ 30 00
... ..		\$ 30 00
... ..		\$ 15 00
... ..		2 00
... ..		5 00
... ..		5 00
... ..		10 00
... ..		10 00

MEMORANDA.

AUGUST.

MEMORANDA.

AUGUST.

MEMORANDA.

SEPTEMBER.

MEMORANDA.

SEPTEMBER.

MEMORANDA.

OCTOBER.

MEMORANDA.

OCTOBER.

MEMORANDA.

NOVEMBER.

MEMORANDA.

NOVEMBER

MEMORANDA.

DECEMBER.

MEMORANDA.

DECEMBER.

ADDRESSES OF PATIENTS AND OTHERS.

Name.

Address.

ADDRESSES OF PATIENTS AND OTHERS.

Name.

Address.

ADDRESSES OF PATIENTS AND OTHERS.

Name.

Address.

ADDRESSES OF PATIENTS AND OTHERS.

Name.

Address.

ADDRESSES OF PATIENTS AND OTHERS.

Name.

Address.

ERS

ADDRESSES OF PATIENTS AND OTHERS.

Name.

Address.

no.

NURSES' ADDRESSES.

Name and Address.

NURSES' ADDRESSES.

Name and Address.

BILLS AND ACCOUNTS ASKED FOR.

Delivered.

Name.

Amount.

BILLS AND ACCOUNTS ASKED FOR.

Delivered.

Name.

Amount.

BILLS AND ACCOUNTS ASKED FOR.

Delivered.

Name.

Amount.

BILLS AND ACCOUNTS ASKED FOR.

Delivered.

Name.

Amount.

MEMORANDA OF WANTS.

MEMORANDA OF WANTS.

MEMORANDA OF WANTS

MEMORANDA OF WANTS.

OBSTETRIC ENGAGEMENTS.

Name and Address.

OBSTETRIC ENGAGEMENTS.

*Time of Expected
Confinement.*

*Date of
Attendance.*

Remarks.

OBSTETRIC ENGAGEMENTS.

Name and Address.

OBSTETRIC ENGAGEMENTS.

*Time of Expected
Confinement.*

*Date of
Attendance.*

Remarks

VACCINATION ENGAGEMENTS.

Name and Address.

*Date of
Vaccination*

*Taken
or not.*

VACCINATION ENGAGEMENTS.

Remarks

VACCINATION ENGAGEMENTS.

Name and Address.

*Date of
Vaccination.*

*Taken
or not.*

VACCINATION ENGAGEMENTS.

Remarks.

JANUARY.

Date.	Name of Child.	Sex.	Color.	Residence.
-------	----------------	------	--------	------------

FEBRUARY

Date	Name of Child.	Sex.	Color	Residence
------	----------------	------	-------	-----------

OBSTETRIC CASES.

Parents' Name.

Occupation.

OBSTETRIC CASES.

Parents' Name.

Occupation.

MARCH.

Date.	Name of Child.	Sex.	Color.	Residence.
-------	----------------	------	--------	------------

APRIL.

Date.	Name of Child.	Sex.	Color.	Residence.
-------	----------------	------	--------	------------

OBSTETRIC CASES.

Parents' Name.

Occupation.

OBSTETRIC CASES

Parents' Name.

Occupation.

MAY.

Date,	Name of Child,	Sex,	Color,	Residence,
-------	----------------	------	--------	------------

JUNE.

Date.	Name of Child.	Sex.	Color.	Residence.
-------	----------------	------	--------	------------

OBSTETRIC CASES

Parents' Name.

Occupation

OBSTETRIC CASES.

Parents' Name.

Occupation

JULY.

<i>Date.</i>	<i>Name of Child.</i>	<i>Sex.</i>	<i>Color.</i>	<i>Residence.</i>
--------------	-----------------------	-------------	---------------	-------------------

AUGUST.

Date.	Name of Child.	Sex.	Color.	Residence.
-------	----------------	------	--------	------------

OBSTETRIC CASES

Parents' Name,

Occupation

OBSTETRIC CASES.

Parents' Name,

Occupation.

SEPTEMBER.

Date.	Name of Child.	Sex.	Color.	Residence.
-------	----------------	------	--------	------------

OCTOBER.

Date	Name of Child	Sex	Color	Residence
------	---------------	-----	-------	-----------

OBSTETRIC CASES.

Parents' Name.

Occupation

OBSTETRIC CASES.

Parents' Name.

Occupation.

NOVEMBER.

[illegible]

DECEMBER.

Date	Name of Child.	Sex.	Color.	Residence.
------	----------------	------	--------	------------

OBSTETRIC CASES.

Parents' Name.

Occupation.

OBSTETRIC CASES.

Parents' Name.

Occupation.

RECORD OF

Name.

Color.

Sex.

Age.

DEATHS.

*Married
Single.*

Date of Death.

Cause of Death.

GENERAL MEMORANDA.

Temperature of the air, water, &c.

GENERAL MEMORANDA.

Jan 1880

GENERAL MEMORANDA.

GENERAL MEMORANDA.

GENERAL MEMORANDA

GENERAL MEMORANDA.

GENERAL MEMORANDA.

GENERAL MEMORANDA.

GENERAL MEMORANDA.

GENERAL MEMORANDA.

GENERAL MEMORANDA.

[Faint handwritten notes]

32

2. 2. 2. 2. 2.

4/650
 400
 150
 300
 1000
 1000
 1000
 1000

CATALOGUE No. 7.



A SELECT LIST
OF
MEDICAL BOOKS.

INCLUDING A FULL LIST OF

The ?Quiz-Compend?

AND MANY OF

THE MOST PROMINENT

Students' Manuals and Text-Books

PUBLISHED BY

P. BLAKISTON, SON & CO.,

Medical Booksellers, Importers and Publishers,

No. 1012 WALNUT STREET,

PHILADELPHIA.

For sale by all Booksellers, or any book will be sent by mail, postpaid, upon receipt of price. Catalogues of books on all branches of Medicine, Dentistry, Pharmacy, etc., supplied upon application.

THE ?QUIZ-COMPENDS?

A NEW SERIES OF COMPENDS FOR STUDENTS.

For Use in the Quiz Class and when
Preparing for Examinations.

Price of Each. Bound in Cloth. \$1.00 Interleaved, \$1.25.

Based on the most popular text-books, and on the lectures of prominent professors, they form a most complete set of manuals, containing information nowhere else collected in such a condensed, practical shape. The authors have had large experience as quiz masters and attachés of colleges, with exceptional opportunities for noting the most recent advances and methods. The arrangement of the subjects, illustrations, types, etc., are all of the most improved form, and the size of the books is such that they may be easily carried in the pocket.

No. 1. ANATOMY. (Illustrated.)

THIRD REVISED EDITION.

A Compend of Human Anatomy. By SAMUEL O. L. POTTER, M.A., M.D., U. S. Army. With 63 Illustrations.

"The work is reliable and complete, and just what the student needs in reviewing the subject for his examinations."—*The Physician and Surgeon's Investigator*, Buffalo, N. Y.

"To those desiring to post themselves hurriedly for examination, this little book will be useful in refreshing the memory."—*New Orleans Medical and Surgical Journal*.

"The arrangement is well calculated to facilitate accurate memorizing, and the illustrations are clear and good."—*North Carolina Medical Journal*.

Nos. 2 and 3. PRACTICE.

A Compend of the Practice of Medicine, especially adapted to the use of Students. By DAN'L E. HUGHES, M.D., Demonstrator of Clinical Medicine in Jefferson Medical College, Philadelphia. In two parts.

PART I.—Continued, Eruptive, and Periodical Fevers, Diseases of the Stomach, Intestines, Peritoneum, Biliary Passages, Liver, Kidneys, etc., and General Diseases, etc.

PART II.—Diseases of the Respiratory System, Circulatory System, and Nervous System; Diseases of the Blood, etc.

. These little books can be regarded as a full set of notes upon the Practice of Medicine, containing the

THE ? QUIZ-COMPENDS ?

Synonyms, Definitions, Causes, Symptoms, Prognosis, Diagnosis, Treatment, etc., of each disease, and including a number of new prescriptions. They have been compiled from the lectures of prominent Professors, and reference has been made to the latest writings of Professors FLINT, DA COSTA, REYNOLDS, BARTHOLOW, ROBERTS and others.

"It is brief and concise, and at the same time possesses an accuracy not generally found in compends."—*Jas. M. French, M.D., Ass't to the Prof. of Practice, Medical College of Ohio, Cincinnati.*

"The book seems very concise, yet very comprehensive. An unusually superior book."—*Dr. E. T. Ryan, Demonstrator of Clinical Medicine, University of Pennsylvania.*

"I have used it considerably in connection with my branches in the Quiz-class of the University of La."—*J. H. Bemiss, New Orleans.*

"Dr. Hughes has prepared a very useful little book, and I shall take pleasure in advising my class to use it."—*Dr. George W. Hall, Professor of Practice, St. Louis College of Physicians and Surgeons.*

No. 4. PHYSIOLOGY.

A Compend of Human Physiology, adapted to the use of Students. By ALBERT P. BRUBAKER, M.D., Demonstrator of Physiology in Jefferson Medical College, Philadelphia.

"Dr. Brubaker deserves the hearty thanks of medical students for his *Compend of Physiology*. He has arranged the fundamental and practical principles of the science in a peculiarly inviting and accessible manner. I have already introduced the work to my class."—*Maurice N. Miller, M.D., Instructor in Practical Physiology, formerly Demonstrator of Physiology, University City of New York.*

"'Quiz-Compend' No. 4 is fully up to the high standard established by its predecessors of the same series."—*Medical Bulletin Philadelphia.*

"I can recommend it as a valuable aid to the student."—*C. N. Hinwood, M.D., Professor of Physiology, Cooper Medical College, San Francisco.*

"This is a well-written little book."—*London Lancet.*

No. 5. OBSTETRICS.

A Compend of Obstetrics. For Physicians and Students. By HENRY G. LANDIS, M.D., Professor of Obstetrics and Diseases of Women, in Starling Medical College, Columbus. Illustrated.

"We have no doubt that many students will find in it a most valuable aid in preparing for examination."—*The American Journal of Obstetrics.*

"It is complete, accurate and scientific. The very best book of its kind I have seen."—*J. S. Knox, M.D., Lecturer on Obstetrics, Rush Medical College, Chicago.*

Price of each Book, Cloth, \$1.00. Interleaved for Notes, \$1.25.

THE ? QUIZ-COMPENDS ?.

"I have been teaching in this department for many years, and am free to say that this will be the best assistant I ever had. It is accurate and comprehensive, but brief and pointed."—*Prof. P. D. Yost, St. Louis.*

No. 6. MATERIA MEDICA. Revised Ed.

A Compend on Materia Medica and Therapeutics, with especial reference to the Physiological Actions of Drugs. For the use of Medical, Dental, and Pharmaceutical Students and Practitioners. Based on the New Revision (Sixth) of the U. S. Pharmacopœia, and including many unofficial remedies. By SAMUEL O. L. POTTER, M.A., M.D., U. S. Army.

"I have examined the little volume carefully, and find it just such a book as I require in my private Quiz, and shall certainly recommend it to my classes. Your Compends are all popular here in Washington."—*John E. Brackett, M.D., Professor of Materia Medica and Therapeutics, Howard Medical College, Washington.*

"Part of a series of small but valuable text-books. . . . While the work is, owing to its therapeutic contents, more useful to the medical student, the pharmaceutical student may derive much useful information from it."—*N. Y. Pharmaceutical Record.*

No. 7. CHEMISTRY. Revised Ed.

A Compend of Chemistry. By G. MASON WARD, M.D., Demonstrator of Chemistry in Jefferson Medical College, Philadelphia. Including Table of Elements and various Analytical Tables.

"Brief, but excellent. . . . It will doubtless prove an admirable aid to the student, by fixing these facts in his memory. It is worthy the study of both medical and pharmaceutical students in this branch."—*Pharmaceutical Record, New York.*

No. 8. VISCERAL ANATOMY.

A Compend of Visceral Anatomy. By SAMUEL O. L. POTTER, M.A., M.D., U. S. Army. With 40 Illustrations.

*** This is the only Compend that contains full descriptions of the viscera, and will, together with No. 1 of this series, form the only complete Compend of Anatomy published.

No. 9. SURGERY. Illustrated.

A Compend of Surgery; including Fractures, Wounds, Dislocations, Sprains, Amputations and other operations, Inflammation, Suppuration, Ulcers, Syphilis, Tumors, Shock, etc. Diseases of the Spine, Ear, Eye, Bladder, Testicles, Anus, and other Surgical Diseases. By ORVILLE HORWITZ, A.M., M.D., with 43 Illustrations.

Price of Each, Cloth, \$1.00. Interleaved for Notes, \$1.25.

THE ? QUIZ-COMPENDS ?

No. 10. ORGANIC CHEMISTRY.

JUST PUBLISHED.

A Compend of Organic Chemistry, including Medical Chemistry, Urine Analysis, and the Analysis of Water, and Food, etc. By HENRY LEFFMANN, M.D., Professor of Clinical Chemistry and Hygiene in the Philadelphia Polyclinic; Professor of Chemistry, Pennsylvania College of Dental Surgery; Member of the N. Y. Medico-Legal Society. Cloth. \$1.00

Interleaved, for the addition of Notes. \$1.25.

Nature of Organic Bodies. Transformations under various conditions. Organic Synthesis. Homologous and Isomeric Bodies. Empirical and Rational formulæ. Classification of organic bodies. Hydrocarbon. Derivatives of Hydrocarbons. Alcohols and Ethers. Benzenes and Turpenes. Fat Acids, Oils and Fats, Sugars, Glucosides. Cyanogen Compounds. Amines and Amides. Alkaloids. Ptomaines. Animal Chemistry. Nutrition and Assimilation. Food, Water and Air. Urinary Analysis. Index.

The Essentials of Pathology.

BY D. TOD GILLIAM, M.D.,

Professor of Physiology in Starting Medical College, Columbus, O

With 47 Illustrations. 12mo. Cloth. Price \$2.00.

*.*The object of this book is to unfold to the beginner the fundamentals of pathology in a plain, practical way, and by bringing them within easy comprehension to increase his interest in the study of the subject. Though it will not altogether supplant larger works, it will be found to impart clear-cut conceptions of the generally accepted doctrines of the day, and to prevent confusion in the mind of the student.

A POCKET-BOOK OF

PHYSICAL DIAGNOSIS

OF THE

Diseases of the Heart and Lungs.

A MANUAL FOR STUDENTS AND PHYSICIANS

BY DR. EDWARD T. BRUEN,

Demonstrator of Clinical Medicine in the University of Pennsylvania. Assistant Physician to the University Hospital, etc.

Second Edition. Revised. With new Illustrations. 12mo. \$1.50

*.*The subject is treated in a plain, practical manner, avoiding questions of historical or theoretical interest, and without laying special claim to originality of matter, the author has made a book that presents the somewhat difficult points of Physical Diagnosis clearly and distinctly.

STUDENTS' MANUALS.

- TYSON, ON THE URINE.** A Practical Guide to the Examination of Urine. For Physicians and Students. By JAMES TYSON, M.D., Professor of Pathology and Morbid Anatomy, University of Pennsylvania. With Colored Plates and Wood Engravings. Fourth Edition. 12mo, cloth, \$1.50
- HEATH'S MINOR SURGERY.** A Manual of Minor Surgery and Bandaging. By CHRISTOPHER HEATH, M.D., Surgeon to University College Hospital, London. 6th Edition. 115 Ill. 12mo, cloth, \$2.00
- REESE. A MANUAL OF MEDICAL JURISPRUDENCE** and Toxicology, for Students and Physicians. Small 8vo, 550 pages.
- VIRCHOW'S POST-MORTEM.** Post-Mortem Examinations. A Description and Explanation of the Methods of Performing them. By PROF. RUDOLPH VIRCHOW, of Berlin. Translated by DR. T. B. SMITH. 2d Ed. 4 Lithographic Plates. 12mo, cloth, \$1.25
- DULLES' ACCIDENTS AND EMERGENCIES.** What To Do First in Accidents and Emergencies. A Manual Explaining the Treatment of Surgical and other Accidents, Poisoning, etc. By CHARLES W. DULLES, M.D., Surgeon Out-door Department, Presbyterian Hospital, Philadelphia. Colored Plate and other Illustrations. 32mo, cloth, .75
- BEALE, ON SLIGHT AILMENTS.** Their Nature and Treatment. By LIONEL S. BEALE, M.D., F.R.S. Second Edition, Revised, Enlarged and Illustrated. 283 pages. 8vo.
Paper covers, 75 cents; cloth, \$1.25
- ALLINGHAM, ON THE RECTUM.** Fistula, Hemorrhoids, Painful Ulcer, Stricture, Prolapsus, and other Diseases of the Rectum; Their Diagnosis and Treatment. By WM. ALLINGHAM, M.D. Fourth Revised and Enlarged Edition. Illustrated. 8vo.
Paper covers, 75 cents; cloth, \$1.25
- THOMPSON, ON THE URINARY ORGANS.** On Diseases of the Urinary Organs. By SIR HENRY THOMPSON, M.D., F.R.C.S. Seventh Edition. 84 Illustrations. 8vo. Paper covers, 75 cents; cloth, \$1.25

STUDENTS' MANUALS.

MARSHALL AND SMITH, ON THE URINE.

The Chemical Analysis of the Urine. By JOHN MARSHALL, M.D., Chemical Laboratory, University of Pennsylvania, and PROF. E. F. SMITH. Illus. Cloth, \$1.00

MEARS' PRACTICAL SURGERY.

Surgical Dressings, Bandaging, Ligation, Amputation, etc. By J. EWING MEARS, M.D., Demonstrator of Surgery in Jefferson Med. College. 227 Illus. 2d Ed. *In Press*

HOLDEN'S ANATOMY.

A Manual of the Dissection of the Human Body. Fifth Edition, Revised and Enlarged, with over 170 Illustrations.

BLOXAM'S LABORATORY TEACHINGS

Progressive Exercises in Practical Chemistry. By PROF. C. L. BLOXAM. 80 Illustrations. 12mo, cloth, \$1.75

TYSON, ON THE CELL DOCTRINE: its His-

tory and Present State. By PROF. JAMES TYSON, M.D. Second Edition. Illustrated. 12mo, cloth, \$2.00

MEADOWS' MIDWIFERY.

A Manual for Students. By ALFRED MEADOWS, M.D. From Fourth London Edition. 145 Illustrations. 8vo, cloth, \$2.00

WYTHE'S DOSE AND SYMPTOM BOOK.

Containing the Doses and Uses of all the principal Articles of the Materia Medica, etc. Eleventh Edition. 32mo, cloth, \$1.00; pocket-book style, \$1.25

PHYSICIAN'S PRESCRIPTION BOOK.

Containing Lists of Terms, Phrases, Contractions and Abbreviations used in Prescriptions, Explanatory Notes, Grammatical Construction of Prescriptions, etc., etc. By PROF. JONATHAN PEREIRA, M.D. Sixteenth Edition. 32mo, cloth, \$1.00; pocket-book style, \$1.25

POCKET LEXICONS.

CLEAVELAND'S POCKET MEDICAL LEXI-

CON. A Medical Lexicon, containing correct Pronunciation and Definition of Terms used in Medicine and the Collateral Sciences. Thirtieth Edition. Very small pocket size. Red Edges.

Cloth, 75 cents; pocket-book style, \$1.00

LONGLEY'S POCKET DICTIONARY.

The Student's Medical Lexicon, giving Definition and Pronunciation of all Terms used in Medicine, with an Appendix giving Poisons and Their Antidotes, Abbreviations used in Prescriptions, Metric Scale of Doses, etc. 24mo, cloth, \$1.00; pocket-book style, \$1.25

ROBERTS' PRACTICE.

FIFTH EDITION.

Recommended as a Text-book at University of Pennsylvania, Long Island College Hospital, Yale and Harvard Colleges, Bishop's College, Montreal, University of Michigan, and over twenty other Medical Schools.

A HANDBOOK OF THE THEORY AND PRACTICE OF MEDICINE. By FREDERICK T. ROBERTS, M.D., M.R.C.P., Professor of Clinical Medicine and Therapeutics in University College Hospital, London. Fifth Edition. Octavo.

CLOTH, \$5.00; LEATHER, \$6.00.

. This new edition has been subjected to a careful revision. Many chapters have been rewritten. Important alterations and additions have been made throughout, and new illustrations introduced.

"A clear, yet concise, scientific and practical work. It is a capital compendium of the classified knowledge of the subject."—*Prof. J. Adams Allen, Rush Medical College, Chicago.*

"I have become thoroughly convinced of its great value, and have cordially recommended it to my class in Yale College."—*Prof. David P. Smith.*

"I have examined it with some care, and think it a good book, and shall take pleasure in mentioning it among the works which may properly be put in the hands of students."—*A. B. Palmer, Prof. of the Practice of Medicine, University of Michigan.*

"It is unsurpassed by any work that has fallen into our hands, as a compendium for students preparing for examination. It is thoroughly practical, and fully up to the times."—*The Clinic.*

"Our opinion of it is one of almost unqualified praise. The style is clear, and the amount of useful and, indeed, indispensable information which it contains is marvelous."—*Boston Medical and Surgical Journal.*

BIDDLE'S MATERIA MEDICA.

NINTH REVISED EDITION.

Recommended as a Text-book at Yale College, University of Michigan, College of Physicians and Surgeons, Baltimore, Baltimore Medical College, Louisville Medical College, and a number of other Colleges throughout the U. S.

BIDDLE'S MATERIA MEDICA. For the Use of Students and Physicians. By the late PROF. JOHN B. BIDDLE, M.D., Professor of Materia Medica in Jefferson Medical College, Philadelphia. The Ninth Edition, thoroughly revised, and in many parts rewritten, by his son, CLEMENT BIDDLE, M.D., Past Assistant Surgeon, U. S. Navy, assisted by HENRY MORRIS, M.D.

CLOTH, \$4.00; LEATHER, \$4.75.

"I shall unhesitatingly recommend it (the 6th Edition) to my students at the DEERFIELD HOSPITAL MEDICAL COLLEGE. *Prof. A. A. Smith, New York, June, 1883.*

"The standard 'Materia Medica' with a large number of medical students is Biddle's."—*Buffalo Medical and Surgical Journal.*

"The larger works usually recommended as text-books in our medical schools are too voluminous for convenient use. This work will be found to contain in a condensed form all that is most valuable, and will supply students with a reliable guide."—*Chicago Med. J.*

. This Ninth Edition contains all the additions and changes in the U. S. Pharmacopœia, Sixth Revision

STANDARD TEXT-BOOKS.

BLOXAM'S CHEMISTRY. Inorganic and Organic, with Experiments. Fifth Edition. Revised and Illustrated.

8vo, cloth, \$1.75; leather, \$4.75

CARPENTER ON THE MICROSCOPE and Its Revelations. Sixth Edition, Enlarged. With 500 Illustrations and Colored Plates, handsomely printed.

Demi 8vo, cloth, \$3.50

DRUITT'S SURGERY. A Manual of Modern Surgery. Eleventh London Edition. 360 Illustrations. Demi 8vo, cloth, \$5.00

FLOWER, DIAGRAMS OF THE NERVES of the Human Body, Origin, Divisions, Connections, etc. 4to. Cloth, \$4.50

GALLABIN'S MIDWIFERY. A Manual for Students. Illustrated.

In Preparation

GLISAN'S MODERN MIDWIFERY. A Text-book. 120 Illustrations.

8vo, cloth, \$4.00; leather, \$8.00

HOLDEN'S ANATOMY and Manual of Dissections of the Human Body. Fourth Edition. Illus. New Ed. *In Press*

HOLDEN'S OSTEOLOGY. A Description of the Bones, with Colored Delineations of the Attachments of the Muscles. Sixth Edition. 61 Lithographic Plates and many Wood Engravings.

Royal 8vo, cloth, \$1.00

HEATH'S PRACTICAL ANATOMY and Manual of Dissections. Fifth Edition. 24 Colored Plates and nearly 300 other Illustrations.

Demi 8vo, cloth, \$2.00

HEADLAND, THE ACTION OF MEDICINE in the System. Ninth American Edition.

8vo, cloth, \$1.00

KIRKE'S PHYSIOLOGY. A Handbook for Students. Tenth Edition. 420 Illustrations.

Demi 8vo, cloth, \$1.00

MANN'S PSYCHOLOGICAL MEDICINE and Allied Nervous Diseases; including the Medico-Legal Aspects of Insanity. With Illustrations.

8vo, cloth, \$5.00; leather, \$8.00

MACNAMARA ON THE EYE. A Manual for Students and Physicians. Illustrated.

Demi 8vo, cloth, \$1.50

MEIGS AND PEPPER ON CHILDREN. A Practical Treatise on Diseases of Children. Seventh Edition, Revised.

8vo, cloth, \$6.00; leather, \$7.00

PARKES' PRACTICAL HYGIENE. Sixth Revised and Enlarged Edition. Illustrated.

8vo, cloth, \$1.00

RIGBY'S OBSTETRIC MEMORANDA.

8vo, cloth, \$1.00

SANDERSON & FOSTER'S PHYSIOLOGICAL LABORATORY. A Handbook for the Laboratory. Over 350 Illustrations.

8vo, cloth, \$5.00; leather, \$6.00

WILSON'S HUMAN ANATOMY. General and Special Tenth Edition. 26 Colored Plates and 424 Illustrations.

\$6.00

WYTHE'S MICROSCOPIST. A Manual of Microscopy and Compend of the Microscopic Sciences. Fourth Edition. 252 Illustrations.

8vo, cloth, \$3.00; leather, \$4.00

AITKEN, THE SCIENCE AND PRACTICE OF MEDICINE. A New (Seventh) Edition. 2 Vols

8vo, cloth, \$12.00; leather, \$14.00

ACTON, ON THE REPRODUCTIVE ORGANS. Their Functions, Disorders and Treatment. 7th Edition. Cloth, \$2.00

FENNER, ON VISION. Its Optical Defects and the Adaptation of Spectacles. 2d Edition, Enlarged. Illus. 8vo, cloth, \$3.50

FOTHERGILL, ON THE HEART. Its Diseases and their Treatment. Second Edition.

8vo, cloth, \$3.50

HARLEY ON THE LIVER. Diagnosis and Treatment. Colored Plates and other Illustrations. 8vo, cloth, \$5.00; sheep, \$9.00

INDEX OF DISEASES; WITH TREATMENT AND FORMULÆ.

By THOS. HAWKES TANNER, M.D.

REVISED AND ENLARGED BY DR. BROADBENT.

Octavo, Cloth. Price \$3.00.

The worth of a work of this kind, by so eminent a professor as Dr. Tanner, cannot be over-estimated. As an aid to physicians and druggists, both in the country and city, it must be invaluable. It contains a full list of all diseases, arranged in alphabetical order, with list of formulæ, and appendix giving points of interest regarding health resorts, mineral waters, and information about cooking and preparing food, etc., for the invalid and convalescent. The page headings are so indexed that the reader is enabled to find at once the disease wanted; its synonyms, classification, varieties, description, etc., with the course of treatment recommended by the best authorities, and is referred, by number, to the several prescriptions that have proved most efficacious. These prescriptions are also arranged so that they can be easily referred to, with directions how to use them, when to use them, and what diseases they are generally used in treating. The directions for cooking foods and preparing poultices, lotions, etc., are very full. The work will be found specially useful to students and young physicians.

RICHTER'S CHEMISTRY,

A TEXT-BOOK of INORGANIC CHEMISTRY for STUDENTS

By PROF. VICTOR von RICHTER,

University of Breslau,

AUTHORIZED TRANSLATION FROM THE THIRD GERMAN EDITION.

By EDGAR F. SMITH, M.A., Ph.D.,

*Professor of Chemistry in Wittenberg College, Springfield, Ohio;
formerly in the Laboratories of the University of Pennsylvania;
Member of the Chemical Society of Berlin.*

12mo. 89 Wood-cuts and Col. Lithographic Plate of Spectra. \$2.00

In the chemical text-books of the present day, one of the striking features and difficulties we have to contend with is the separate presentation of the theories and facts of the science. These are usually taught apart, as if entirely independent of each other, and those experienced in teaching the subject know only too well the trouble encountered in attempting to get the student properly interested in the science and in bringing him to a clear comprehension of the same. In this work of PROF. VON RICHTER, which has been received abroad with such hearty welcome, two editions having been rapidly disposed of, theory and fact are brought close together, and their intimate relation clearly shown. From careful observation of experiments and their results, the student is led to a correct understanding of the interesting principles of chemistry. The descriptions of the various inorganic substances are full, and embody the results of the latest discoveries.

In preparation, "ORGANIC CHEMISTRY," By the same author. Translated.

BYFORD, DISEASES OF WOMEN.

NEW REVISED EDITION.

The Practice of Medicine and Surgery, as applied to the Diseases of Women. By W. H. BYFORD, A.M., M.D., Professor of Gynæcology in Rush Medical College; of Obstetrics in the Woman's Medical College; Surgeon to the Woman's Hospital; President of the American Gynæcological Society, etc. Third Edition. Revised and Enlarged; much of it Rewritten; with over 160 Illustrations. Octavo.

PRICE, CLOTH, \$5.00; LEATHER, \$6.00.

"The treatise is as complete a one as the present state of our science will admit of being written. We commend it to the diligent study of every practitioner and student, as a work calculated to inculcate sound principles and lead to enlightened practice."—*New York Medical Record*.

"The author is an experienced writer, an able teacher in his department, and has embodied in the present work the results of a wide field of practical observation. We have not had time to read its pages critically, but freely commend it to all our readers, as one of the most valuable practical works issued from the American press."—*Chicago Medical Examiner*.

DAY ON CHILDREN. Second Edition.

The Diseases of Children. A Practical and Systematic Treatise for Practitioners and Students. By WILLIAM H. DAY, M.D., Author of "Headaches; Their Causes, Nature and Treatment;" Member of the Royal College of Physicians of London; Physician to the Samaritan Hospital for Women and Children, etc. Second Edition. Rewritten and very much Enlarged. 8vo. 752 pp.

PRICE. CLOTH, \$5.00; SHEEP, \$6.00.

"A careful examination of this book leads us to characterize it as a plain, straightforward treatise on the subject upon which it treats . . . giving sound practical advice."—*Philad'a Med. Times*.

"On the whole, we must confess we are pleased with this book and can heartily recommend it—a recommendation which it does not appear to need, as it has already reached its second edition."—*American Journal of Medical Science*.

MACKENZIE, THE THROAT AND NOSE.

By MORELL MACKENZIE, M.D., Senior Physician to the Hospital for Diseases of the Chest and Throat; Lecturer on Diseases of the Throat at the London Hospital, etc.

VOL. I. Including the PHARYNX, LARYNX, TRACHEA, etc. 112 Illustrations. Cloth, \$4.00; Leather, \$5.00.

VOL. II. DISEASES OF THE OESOPHAGUS, NASAL CAVITIES AND NECK. Cloth, \$3.00; Leather, \$4.00.

The two volumes at one time. Cloth, \$6.00; Leather, \$7.50.

YEO'S PHYSIOLOGY.

A MANUAL FOR STUDENTS. JUST READY.

300 CAREFULLY PRINTED ILLUSTRATIONS.

FULL GLOSSARY AND INDEX.

By **GERALD F. YEO, M.D., F.R.C.S.**, Professor of Physiology in King's College, London. Small Octavo. 750 pages. Over 300 carefully printed Illustrations.

PRICE, CLOTH. \$4.00; LEATHER, \$5.00.

"By his excellent manual, Prof. Yeo has supplied a want which must have been felt by every teacher of physiology. In conclusion, we heartily congratulate Prof. Yeo on his work, which we can recommend to all those who wish to find within a moderate compass a readable and pleasantly written exposition of all the essential facts of physiology as the science now stands."—*The Director General of the Medical Service.*

"The work will take a high rank among the smaller text-books of Physiology."—*Prof. H. P. Bowditch, Harvard Med. School, Boston.*

"The brief examination I have given it was so favorable that I placed it in the list of text-books recommended in the circular of the University Medical College."—*Prof. Lewis A. Stimpson, M.D., 37 East 23d Street, New York.*

"For students' use it is one of the very best text-books in Physiology."—*Prof. L. B. How, Dartmouth Med. College, Hanover, N. H.*

RINDFLEISCH.

THE ELEMENTS OF PATHOLOGY.

TRANSLATED BY **WM. H. MERCUR, M.D.**

REVISED AND EDITED BY **PROF. JAS. TYSON,**
Of the University of Pennsylvania.

OCTAVO. CLOTH. NEARLY READY.

*It is the object of Prof. Rindfleisch to present in this volume of moderate size the fundamental principles of Pathology. A large number of the general processes which underlie disease, a knowledge of which is essential to the practical physician, are plainly presented. They include, among others, inflammation, tumor formation, fever, derangements of nutrition, including atrophy, derangements of the movement of the blood, of blood formation and blood purification, hyperaesthesia, anaesthesia, convulsions, paralysis, etc. The well-known reputation of the author, his thorough familiarity with, and his method of treating the subject, make this most recent work peculiarly useful to the student, as well as to the practicing physician who wishes to brush up his pathology.

THE
MEDICAL NEWS
VISITING-LIST.
1886.



PHILADELPHIA:
LEA BROTHERS & CO.
1885.


Entered according to Act of Congress, in the year 1885, by
LEA BROTHERS & CO.,
in the Office of the Librarian of Congress. All rights reserved.

WESTCOTT & THOMSON,
Stereotypers and Electrotypers, Philada.

WILLIAM J. DORNAN,
Printer, Philada.

CONTENTS.

	PAGE
PREFACE	5
HOW TO KEEP ACCOUNTS	7
HOW TO KEEP THE VISITING-LIST	7
TABLE TO FIND DAY OF CONFINEMENT	8
SIGNS OF PREGNANCY	8
SIGNS OF DENTITION	8
WEIGHTS AND MEASURES (ORDINARY AND METRIC)	9-10
COMPARATIVE SCALES	11
EXAMINATION OF URINE	12-13
DISINFECTANTS	13
TABLE OF ERUPTIVE FEVERS	14
SOME REMEDIES NOT IN GENERAL USE	15-18
INCOMPATIBLES	18
ARTIFICIAL RESPIRATION	19
POISONS AND ANTIDOTES	20-21
TABLE OF DOSES	22-26
THERAPEUTIC TABLE	27-43
LIGATION OF ARTERIES	43-48
DAILY RECORD OF PRACTICE	50-155
CLINICAL RECORD	156-183
CONSULTATION PRACTICE	184-189
OBSTETRIC ENGAGEMENTS	190-195
OBSTETRIC PRACTICE	196-207
VACCINATIONS	208-211
DEATHS	212-215
ADDRESSES OF PATIENTS	216-219
ADDRESSES OF NURSES	220-221
CASH ACCOUNT	222-236
GENERAL MEMORANDA	237-240



PREFACE.

THIS Visiting List has been arranged so as to include a number of desirable features, some of which are now to be found scattered about in a variety of similar books, others of which are here presented for the first time. A visiting list should not be expected to contain a treatise on the practice of medicine. It should, however, furnish brief memoranda which may be useful in an emergency, or in the moments, which come to the best of men, when one cannot recall a fact with which he may be perfectly familiar. A visiting list should also be so arranged as to permit a physician to keep a clear record of his engagements and of his professional work, from which he may transfer to a ledger a permanent record which will have a legal status in case of a dispute. Both these objects have been aimed at in this Visiting List, the purchasers of which are strongly urged to complete their records by the use of a suitable ledger, since the imperfect accounts of visiting lists have been decided to have no legal value as evidence in the collection of debts.



To the
Faint text visible on the right edge of the page, likely from the adjacent page or a marginal note.

THERMOMETRIC SCALES.

To convert Degrees of the Centigrade Scale into those of Fahrenheit: Multiply the number of the former by 9 and divide by 5; if the temperature was above the freezing-point (0°), add 32° ; if it was below the freezing-point, subtract the result from 32° .

To convert Degrees of the Fahrenheit Scale into those of Centigrade: If the temperature be above the freezing point (32°), subtract 32 ; if it be below the freezing point, subtract the number from 32 ; then multiply the remainder by 5 and divide by 9.

In taking temperatures, the thermometer should always be used in the same situation (axilla, mouth, etc.), and left in for the same length of time. It is important also, in keeping a record or making a report, to state the situation of the thermometer.

ORDINARY WEIGHTS AND MEASURES.

Troy, or Apothecaries', Weight.

20 grains (gr.)	= 1 scruple (℥)	= 20 grains.
3 scruples	= 1 drachm (℥)	= 60 "
8 drachms	= 1 ounce (℥)	= 480 "
12 ounces	= 1 pound (lb.)	= 5760 "

Wine, or Apothecaries', Measure.

60 minims (℥)	= 1 fluidrachm.
8 fluidrachms (fldr.)	= 1 fluidounce.
16 fluidounces (floz.)	= 1 pint.
2 pints (O)	= 1 quart.
4 quarts	= 1 gallon (Cong.) = 231 cubic inches.

Linear Measure.

12 lines	= 1 inch.
12 inches	= 1 foot.

THE METRIC SYSTEM.

This system gets its name from the meter, which is the unit of linear measure. It is equal to 39.370432 inches, or a little over three and one-quarter feet.

The gram is the unit of weight, and is equal to 15,434 grains (Troy).

The liter is the unit of solid or fluid measure, and is equal to 1.0567 quarts.

The names of the higher weights and measures are formed by prefixing to the name of the unit the Greek terms deka, hecto, kilo, and myria. The names of the lower weights and measures are formed by prefixing to the name of the unit the Latin terms deci, centi, milli.

TABLE OF LINEAR MEASURES.

1 millimeter (m.m.)	= 1 : 1000 of a meter	= 0.03937 inch.
1 centimeter (c.m.)	= 1 : 100 "	= 0.3937 "
1 decimeter (d.cm.)	= 1 : 10 "	= 3.937 "
1 meter (m.)		= 39.37 "
1 dekameter (d.km.)		= 32 feet 9.7 inches.

METRIC WEIGHTS AND MEASURES,

Length:

1 millimeter		= .0393704 inches.
10 millimeters (m.m.)	= 1 centimeter	= .393704 "
10 centimeters (c.m.)	= 1 decimeter	= 3.937043 "
10 decimeters (d.c.m.)	= 1 Meter	= 39.370432 "
10 meters (m.)	= 1 dekameter	= 32 feet 9.7 inches.
10 dekameters (dkm.)	= 1 hektometer	= 328 " 1. "
10 hektometers (h.m.)	= 1 kilometer	= 3280 " 10.4 "
10 kilometers (k.m.)	= 1 myriameter (myr.m.)	= 6.2137 miles.

Volume:

1 milliliter (m.l.)	= 1 cubic centimeter	= 16.23 minims.
10 milliliters	= 1 centiliter	= 2.71 fluidrachms.
10 centiliters (c.l.)	= 1 deciliter	= 3.38 fluidounces.
10 deciliters (dc.l.)	= 1 Liter (= 1 cubic decimeter)	= 33.81 fluidounces = 1.0567 quarts.
10 liters (l.)	= 1 dekaliter	= 2.6417 gallons.
10 dekaliters (dk.l.)	= 1 hektoliter	= 26.417 "
10 hektoliters (h.l.)	= 1 kiloliter or stere	= 264.17 "


Weight:

1 milligram (m.gm.)	= .001 gram	= .015 grain Troy.
10 "	= 1 centigram	= .154 " "
10 centigrams (c.gm.)	= 1 decigram	= 1.543 " "
10 decigrams (dc.gm.)	= 1 Gram	= 15.432 " "
1 gram (gm.)	weight of 1 cubic centimeter of water at 19° F.	
10 grams	= 1 dekagram	= 25 34.37 grains Troy.
10 dekagrams (dk.gm.)	= 1 hektogram	= 35 103.235 " "
10 hektograms (h.gm.)	= 1 kilogram (k.gm.)	= 2 lb. 83 72 " "

A teaspoon holds about 5 cubic centimeters, and a tablespoon about 20 cubic centimeters. A Troy ounce equals about 30 grams, and an eight-ounce bottle holds about 250 cubic centimeters.

In prescribing, the French usually confine themselves to ordering grams and multiples of the gram for weights, and cubic centimeters and multiples of the cubic centimeter in measures.

COMPARATIVE SCALES, showing at a glance the exact equivalent of ordinary weights and measures in those of the Metric System, and *vice versa*.



The equivalents of fractions, whether large or small, may be found with great accuracy by these scales. For instance, $\frac{1}{2}$ grain = $\frac{1}{2}$ of the metric equivalent of 7 grains, and 1.06 grain 1.4000 of the metric equivalent of 20 grains. This method is, of course, reversible.

EXAMINATION OF THE URINE.

THE most important matters to be investigated in an examination of the urine are—1, the amount passed in a given time; 2, the specific gravity; 3, the reaction; 4, the proportion of urates present; 5, the proportion of phosphates present; 6, the absence or presence of albumen, and the quantity if present; 7, the absence or presence of sugar, and the quantity if present; and, 8, the character of the microscopic sediment.

The examination may be conducted as follows:

1. The amount may be measured by collecting the whole amount passed in a given 24 hours.

2. The specific gravity may be ascertained with a "urinometer."

3. The reaction should be ascertained, immediately or soon after the passage of the sample, by means of slips of blue litmus-paper, which is turned red by acids, or by reddened litmus-paper, which is made blue by alkalis, or by turmeric paper, which is made brown by alkalis.

4. The proportion of urates may be very well estimated by the amount of sediment and turbidity in an *acid* urine. A high specific gravity in an acid urine usually indicates the presence of a proportionate excess of urates. A turbidity which is cleared up by warming the urine also indicates an excess of urates.

5. The proportion of phosphates may be ascertained by adding to some urine, in a test-tube, half its bulk of a solution of caustic potash (1:8), and gently warming the mixture. If the deposit which forms is deeper than a third of an inch in a test-tube three-quarters of an inch wide with two inches of the mixture in it, the phosphates are in excess. If much less, they are below normal. Turbidity of the urine, which is cleared up by the addition of a little acetic acid, is to be set down to an excess of phosphates—unless the urine is stale, when it may be due to alkaline decomposition.

6. The presence of albumen is detected by the appearance of a whitish cloudiness or a curdy deposit upon boiling a little clear urine, after slightly acidulating it with a *few drops* of acetic acid. Another good test is to pour a little urine into a test-tube, then to draw up some nitric acid into a long, pointed glass tube, close the upper end with a forefinger, thrust the point to the bottom of the urine, and then withdraw the finger gradually, so as to permit the acid to flow out slowly at the bottom, under the urine, which will be raised up as the acid flows out. If there be albumen present, it will be disclosed by the formation of a layer of coagulation between the acid and the urine. A light, cloudy appearance a little higher than the line where the acid and the urine are in contact is to be attributed to a precipitation of urates by the acid, and is not a sign of albumen, which coagulates just at the line of contact of acid and urine.

7. The presence of sugar, in quantity sufficient to have any pathological significance, is easily ascertained by adding to some urine, in a test-tube half its bulk of solution of caustic potash (1:8), and boiling the mixture thoroughly. Any noteworthy amount of sugar will cause a distinct deepening of the color, even to making it dark brown or black if the proportion of sugar be very large. A slight change (to a rum color) does not

indicate the presence of sugar, but occurs when there is none or only a physiological quantity present. After the boiling, some muriatic acid may be added, when, if the amount of sugar be large, the odor of burnt sugar may be perceived. A high specific gravity in a clear, syrupy-looking urine is apt to go with the presence of sugar in it.

8. The character of the sediment as revealed by the microscope is of more significance than any other recognizable characteristic of the urine in disease. To conduct this part of the examination requires a good microscope and the ability to recognize the various sedimentary bodies to be found in health or disease.

DISINFECTANTS.

Clothing, Bedding, etc.—If these are not of too much value, the best and simplest method of disinfection is to burn them. If they are worth saving, they may be subjected to prolonged baking or boiling—that is to say, to dry or moist heat—which will effectually destroy any germs which they may hold. Or they may be exposed to the fumes of sulphur, or soaked in a solution of corrosive sublimate and permanganate of potassium, of each 1 ounce in 8 gallons of water.

Bed-pans, Chamber-pots, Water-closets, etc.—These should be washed with boiling-hot water and rinsed with a solution of corrosive sublimate and permanganate of potassium, of each 2 drachms in a gallon of water. A quantity of this solution should always be left in each vessel. According to the most recent investigations, this is the most efficient disinfectant for fecal discharges which we have.

Rooms.—Thorough cleanliness, plenty of fresh air, and sunlight are the most important purifiers of inhabited rooms. To these may sometimes be added the exposure of chloride of lime in saucers. When a room can be given over to thorough disinfection, this may be done by removing all curtains and carpets (which must be disinfected in the manner described for clothing), washing all the solid parts with hot water and soft soap, drying thoroughly, then shutting all doors and windows and filling the room with the fumes of burning sulphur. Fumigation with sulphur is effected by placing tubs of water in the room and standing in the midst of the water a brick, and upon this an earthenware saucer (like that for a flower-pot) in which some sulphur has been placed. From 1 to 3 pounds of sulphur should be allowed for each 1000 cubic feet of air-space. The room should be kept closed for 24 hours, and then thoroughly aired.

Water-closets, Drains, and Sinks.—To disinfect these it is important to cleanse them as well as possible, and the best way to do this is to use a strong solution of lye. There are a number of preparations which may be used, one of which is well known as "Banner Lye." A strong solution of this may be poured into the water-closet pan and allowed to remain there for some time, or into the drain-pipes and sinks, and managed so as to remain in contact with the polluted parts as long as possible. After this the solution of corrosive sublimate and permanganate of potassium recommended for bed-pans may be used.

Privies.—The contents of privies should be frequently and well strewn with chloride of lime.

Carbolic acid, which is often used for disinfection, is a very weak and unreliable agent, except when used in strong concentration. As sometimes used, by being exposed in saucers in sick rooms, it has no value whatsoever.

TABLE OF ERUPTIVE FEVERS.

Disease.	Manner of Invasion.	Characteristics of Eruption.	Course of Disease.
Scarlatina (Scarlet fever).	Usually sudden; chilliness; high fever; rapid pulse; flushed face; convulsions around sore throat; "strawberry tongue."	Appears very early, often on second day. First on neck and chest. May become universal. May be only about joints and very trifling. Bright scarlet; punctate. Followed by desquamation; skin burning hot; sometimes itches.	Progresses steadily and rapidly. Temperature rises to 104°-106°. Defervescence gradual. Conia a dangerous sign. Albumen in urine common, and often true desquamative nephritis.
Rötheln (German measles).	Usually <i>unaccompanied</i> . Slight fever or none; mild catarrhal symptoms; vomiting; slight sore throat; enlargement of cervical lymphatic glands almost always present.	Appears early. Often first symptom noted, first on face and neck, then on trunk. Rosy papules, scattered or clustered. Fades in one part and appears in another. Lasts about three days.	Usually mild. Disappears gradually.
Varicella (Chicken pox).	Usually <i>unaccompanied</i> . Slight fever or none; slight malaise or none.	Appears early. Usually on chest and shoulders, then on body and limbs and face. Distinct, clear, shot-sized vesicles; sometimes with distinct areolae and even umbilicated; appear in successive crops, usually few in number.	Usually mild. Vesicles disappear by drying up. May form pus and leave pits.
Variola (Smallpox).	Usually sudden; chill; high fever; rapid pulse; violent pain in head and back; debility; mucous membranes often involved.	Appears on third or fourth day; first on face, and spreads over whole body and limbs in a day or two. Erythroid like papules; then vesicles; then umbilicated pustules, about eighth day developing adherent crusts; well marked areolae. Intense itching. Peculiar odor.	Symptoms sometimes moderate when eruption first appears, and grow worse as pustules mature. Often severe constitutional disturbance. Great restlessness and desire to scratch the eruption. Defervescence gradual and slow. May be interrupted or exacerbate. Crusts fall off, and often leave deep pits or pocks.
Morbilli (Measles).	Generally sudden; chilliness or even convulsions; moderate fever; lachrymation; catarrh of respiratory passages; bronchitis.	Appears about fourth day; first on face, and spreads to body and limbs. May be limited; usually almost universal. Consists of rosy papules, arranged in parts of circles, crescentic; appears sometimes in successive crops.	Usually mild and regular. Temperature may be high from complications. Lasts about a week. Recovery rapid.

SOME REMEDIES NOT IN GENERAL USE.

Antipyrine.—An artificial hydrocarbon obtained from coal-tar, soluble in three parts of cold and one-half part of warm water. Useful as an antipyretic, especially in typhoid fever, although it does not modify the course of the disease. Does not produce deafness or dizziness. Best excipient is syrup of tolu. Dose, usually 20-30 grains at first, followed by 10-grain doses hourly until temperature has fallen decidedly.

Berberis aquifolium.—Alterative and tonic in strumous and syphilitic disorders and in scaly skin diseases. Dose of the fluid extract, $\frac{1}{2}$ 1 fluidrachm.

Cascara sagrada.—A laxative much praised in the treatment of chronic constipation. May be prescribed in solid or fluid extract. Dose of the former, 1-2 grains; of the latter, 10-60 minims.

Chrysarobin (Chrysophanic acid) is used in the treatment of skin diseases, especially stubborn cases of psoriasis and tinea capitis. It is used in ointment (strength 10 per cent.) and in solution in chloroform (7 grains to the fluidounce), to be well rubbed in. It makes a very indelible stain if it comes into contact with clothing.

Cocaine is the alkaloid of the *Erythroxylon coca*. It is a valuable local anæsthetic, first employed upon the eye, in which it also produces dilatation of the pupil, lachrymation, and slight paralysis of the accommodation. It is very useful as a local anæsthetic to allay the pain of burns, to dull sensation in operations upon the skin and the mucous membranes. It is commonly used in the form of the hydrochlorate, in 2-4 per cent. solution in water. For very rapid action it may be used as strong as 20 per cent.

Convallaria majalis (Lily-of-the-Valley), recently revived as a remedy in heart disease. Said to be useful in cases where digitalis is not well borne. Said to be contraindicated in hepatic disease and in any acute inflammatory disorder. Dose of the fluid extract, 15-30 minims.

Coto, Cotoin, Paracotoin.—Astringent and tonic in obstinate diarrhoea, as that of children, and after typhoid fever or during phthisis. Dose of coto or cotoin, $\frac{1}{2}$ 1 grain; of paracotoin, 2-5 grains; of the fluid extract, 5-15 minims.

Duboisia, Duboisina.—The sulphate or hydrochlorate of this alkaloid is sometimes used to dilate the pupil and to paralyze the muscles of accommodation instead of atropia, because its effects are more prompt and less lasting than those of atropia. It is used in the same strength as atropia, from 2-4 grains to the fluidounce of water. It has the disadvantage of often producing disagreeable though transient cerebral disturbance.

Eucalyptus, Oil of. is a very valuable stimulating expectorant in doses of from 2-5 drops. It has also been used as an antiseptic by external application and by inhalation. For internal administration it can be put in capsules.

Euonymin is the active principle of the *Euonymus atropurpureus*, and is a valuable cholagogue. The dose is 2 grains, to be followed by an aperient the next morning.

Grindelia robusta.—Used in the form of fluid extract, dose 15 minims to a fluidrachm. It has been highly recommended for the treatment of dyspnea dependent on heart disease, as well as asthma, hay asthma, and the asthmatic attacks of bronchitis, whooping cough, and emphysema.

Hamamelis (Witch-Hazel). Used in the form of the fluid extract, dose 1-2 fluidrachms. It is used as a hemostatic and to give tone to the muscular walls of the veins. It has been used in a variety of internal hemorrhages and to cure varicose veins and varicose ulcers of the leg.

Homatropine Hydrobromate is a useful and safe substitute for atropia to dilate the pupil (for which instillation of a few drops of a solution of 1-2 grains in a fluidounce of water is sufficient), and to paralyze the muscles of accommodation (for which a strength of 24 grains to the fluidounce of water is required).

Hydrogen, Peroxide of. is a transparent oily liquid, commonly used in watery solution of 10-15 per cent. It has a high oxidizing power, and when injected into septic cavities is decomposed, the extra atom of oxygen attacking the septic material and disinfecting it. It has also been recommended for the treatment of diphtheria.

Iridin.—An extract of the blue flag. A useful cholagogue, like euonymin. Dose, 4 grains.

Kairine is an artificial hydrocarbon compound. It is used as an antipyretic in doses of from 3-30 grains, best given in capsules or wafer paper. It produces profuse sweating.

Nitro-glycerine has been found to be very useful in relieving the pain of neuralgia and angina pectoris, as well as headache, vertigo, and epilepsy. It is usually administered in a 1 per cent. solution (water or alcohol), the dose beginning with 1 drop, and being increased up to 10 or more drops thrice daily, as it can be tolerated. It may be administered in pill form, made up with cacao-butter, and it has also been made into granules or tablets.

Nitrite of Amyl has effects similar to those of nitro-glycerine, but less lasting. It is administered by inhalation, a few drops being poured on a thin cloth and held before the nose, and withdrawn when the face becomes distinctly flushed or relief has been afforded. This drug is not so dangerous as was supposed when it was first used. But it should not be used recklessly. It is believed to be of value in averting attacks of epilepsy and to relieve the syncope of chloroform-poisoning.

Oleates.—Under this name certain combinations of metals and alkalis with oleic acid have recently been strongly urged upon the profession as superior to a number of preparations in common use. Of these but few are really better than the old preparations. The oleate of zinc, for example, is the finest and smoothest preparation of zinc for external application which has yet been made. The oleate of copper and the oleate of mercury are also excellent preparations for similar employment. Some of the oleates of the alkaloids are apparently good combinations, and the most recent one, an oleate of cocaine, prepared by Dr. Squibb, is claimed to be superior to the watery solution of this drug for application to the skin. There is no evidence that oleates are better than the old preparations with lard when absorption of the drug is desirable, although they are handsomer preparations and keep better.

Paraldehyde has been much used recently as a hypnotic, especially in conditions of high excitement, as mania, and found to produce sound sleep without tendency to coma, it being possible to arouse the patient at once, while he falls asleep again when let alone. It is best administered in combination with syrup of orange or simple elixir; so that a tablespoonful of the mixture shall contain from 25 minims to a fluidrachm of paraldehyde.

Potassium Permanganate has been found useful in the treatment of amenorrhœa. It is to be given in pill form, combined with some non-oxidizable vehicle, such as kaolin (ointment). The dose is from 1-3 grains thrice daily, to be given before the time for menstruating.

Physostigma (Calabar Bean) is used in the form of a tincture, of which the dose is 12 minims, and an extract, of which the dose is from $\frac{1}{4}$ to $\frac{1}{2}$ grain. It is useful in tetanus, strychnia-poisoning, paralysis of the insane, paraplegia, and locomotor ataxia. It has also been used in constipation dependent upon atony of the muscular coat of the bowels. Sulphate of eserine (the active principle) is used, in 1-1 per cent. watery solution, by instillation into the eye, for glaucoma, staphyloma, and to neutralize the effect of mydriatics. It often causes sharp supraorbital and nasal pain for half an hour after the application.

Pilocarpine, the active principle of jaborandi, is used in the form of the hydrochlorate, in doses of from $\frac{1}{2}$ to $\frac{1}{4}$ of a grain. It produces great activity of the salivary and sweat glands, as well as increased secretion from the mucous surfaces and of urine from the kidneys. It is used chiefly in dropsy and uræmia to supplement, as well as to assist, the kidneys. It should be given with care in fatty heart and where there is valvular disease. The fluid extract of jaborandi is sometimes used, but it is not so exact a remedy as pilocarpine, which can be very well administered hypodermically.

Resorcin is an artificial hydrocarbon, used as an antiseptic and antipyretic. It has been found to be of value in the treatment of diphtheria, typhoid fever, and in the fever of phthisis. The dose is from 5-30 grains. It may be given in syrup of orange, well diluted. It produces profuse perspiration and its antipyretic action does not last long.

Santonin, the active principle of *santonica*, is used as an anthelmintic, in doses of from 2-10 grains, depending upon the age of the patient. In the last *U. S. Pharmacopæia* the santoninate of sodium was substituted for santonin. But the latter is much preferable to the former, as it is not so readily absorbed, and is therefore less likely to produce toxic effects.

Viburnum prunifolium (Black Haw).—The fluid extract, of which the dose is 30-60 minims, is said to be very valuable in the prevention of threatened abortion and in menorrhagia. It is, however, by no means a certain remedy.

LIST OF THE MOST IMPORTANT INCOMPATIBLES.

- Acacia (gum), with alcohol, iron, lead-water, and mineral acids.
- Acids (mineral), with alkalis and relatively weak salts of other acids, such as bromides, chlorides, and iodides.
- Alkalis, with acids and with relatively weak salts.
- Arsenic, with tannic acid, salts and oxide of iron, and lime and magnesia.
- Bitter infusions and tinctures, with salts of iron and lead.
- Bromides, with acids, acid salts, or alkalis.
- Calomel, with alkalis, lime-water, salts of iron and lead, and iodide of potassium.
- Camphor (spirits of), with water.
- Carbonates, with acids and acid salts.
- Chlorides, with silver salts, lead salts, and alkalis.
- Chloroform (except in minute quantity), with water.
- Corrosive sublimate, with alkalis, lime-water, salts of iron and lead, iodide of potassium, albumen, gelatine, and vegetable astringents. (It may, however, be advantageously combined with the tincture of the chloride of iron and the liq. acidi arseniosi of the new Pharmacopæia, or, in spite of the chemical reaction, with iodide of potassium, as in the famous Gibert's syrup.)
- Digitalis, with iron and preparations containing tannic acid.
- Iron (salts), with anything containing tannic acid. Tincture of the chloride of iron, with alkalis, carbonates, mucilages, and preparations containing tannic acid.
- Mucilages, with acids, iron salts, and alcohol.
- Potassium (iodide of), with all strong acids and acid salts. (See Corrosive Sublimate.)
- Spirits of nitrous ether, with sulphate of iron, tincture of guaiacum, and most carbonates.
- Tinctures of gums or resins, with water.
- Vegetable preparations containing tannic acid, with salts of iron and lead.

ARTIFICIAL RESPIRATION (Sylvester's Method).

FREE the body from any clothing that binds the neck, chest, or waist; turn it over upon the face for a moment, thrusting a finger into the mouth and sweeping it round, to bring away anything that may have gotten in or accumulated there. Then lay the body flat on the back, with something a few inches high under the shoulders, so as to cause the neck to be stretched out and the chin to be carried from the chest. Draw the tongue well forward out of the mouth and let it be held by an assistant. (If there be no one to do this, a pencil or small stick may be thrust across the mouth on top of the tongue and back of the last teeth, to keep the mouth open and the tongue out of the throat.) Place yourself on your knees behind the head, seize both arms near the elbows and sweep them round horizontally, away from the body and over the head till they meet above it; give a good, strong pull, and keep it up for a




FIG. 1.

few seconds. After this return the arms to their former position alongside the chest, and make strong pressure against the lower ribs, so as to drive the air out of the chest and effect an act of expiration. This need occupy but a second of time.

This plan, regularly carried out, will make about 16 complete acts of respiration in a minute. It should be kept up for a long time, and not aban-



FIG. 2.

doned until the heart has ceased to beat. It should be remembered that cessation of the pulse at the wrists amounts to nothing as a sign of death; and life is present when only a most acute ear can detect the sound of the heart. In a moderately thin person deep pressure with the finger-ends just below the lower end of the breastbone may sometimes reveal pulsation in the aorta when it cannot be found anywhere else.

POISONS AND ANTIDOTES.

THE following table contains suggestions for the proper treatment of every form of poisoning which is likely to occur. Though brief, they are all that will be needed to prompt the physician's memory:

POISON.	TREATMENT.
<i>Nature unknown</i>	{ Provoke repeated vomiting; Give bland liquids; Stimulate, if necessary.
<i>Acids</i> — Sulphuric, } Nitric, } Muratic, } Oxalic, }	{ Give an alkali; Provoke vomiting; Give bland fluids; Secure rest; Stimulate, if necessary.
<i>Prussic Acid</i>	{ Give dilute hartshorn; chlorine-water; so- lution of a salt of iron; Produce vomiting and stimulate.
<i>Carbolic Acid</i>	{ Give Epsom salts, dilute sulphuric acid, and oil; Produce vomiting and stimulate.
<i>Alkalies</i> — Hartshorn, } Soda, } Potash, } Lye, }	{ Give an acid or a fixed oil: Provoke vomiting; Give bland liquids; Secure rest; Stimulate, if necessary.
<i>Arsenic</i> — Paris green, } Scheele's green, } Fowler's solution, }	{ Provoke vomiting; Give hydrated oxide of iron or dialyzed iron and salt; Give dose of castor oil; Secure rest; Stimulate, if necessary.
<i>Acetate of Lead</i>	{ Give Epsom salts or dilute sulphuric acid: Provoke vomiting; Give bland liquids; Give dose of castor oil.
<i>Mercury</i> , <i>Corrosive Sublimate</i> , } <i>Antimony</i> , } <i>Tartar Emetic</i> , }	{ Provoke vomiting; Give some infusion containing tannic acid; Give raw eggs and milk; Give dose of castor oil; Stimulate, if necessary.
<i>Copper Salts</i>	{ Give albumen; Provoke vomiting; Give bland fluids.
<i>Phosphorus</i>	{ Provoke vomiting; Give five-grain doses sulphate of copper; Give dose of magnesia, but no oil.
<i>Nitrate of Silver</i> } (lunar caustic), }	{ Give strong salt and water; } repeat many Provoke vomiting; } times.
<i>Iodine</i>	{ Provoke vomiting; Give starch and water; Give bland fluids.

POISON.	TREATMENT.
Opium— Morphine, Laudanum, Paregoric, etc., Chloral,	{ Provoke vomiting, repeatedly; Give strong coffee, or tea, without milk; Keep up the breathing, preferably with battery.
Strychnine.....	{ Provoke vomiting, once or twice; Give a purgative; Secure absolute quiet.
Aconite	{ Provoke vomiting; Stimulate well.
Jamestown Weed, Hemlock, Nightshade (belladonna), Toadstool, Tobacco, etc., etc.,	{ Provoke vomiting and give a purge; Stimulate well.
Alcohol.....	{ Provoke vomiting; Give hartshorn and water.
Decayed Meat or Vegetables.....	{ Provoke vomiting; Give a purgative; Give powdered charcoal.
Poisonous Gases— Carbonic acid or oxide, Sulphuretted hydrogen,	{ Fresh air; Artificial respiration; Stimulation.

To provoke vomiting. warm water may be used with or without ground mustard (a tablespoonful to a pint of water), or ipecac a teaspoonful of the powder or a tablespoonful or so of the syrup, or a finger may be thrust down the throat. It is best to give large quantities (that is, a pint at a time) of warm water whenever vomiting is to be excited.

Bland liquids are milk, raw eggs, some sort of oil, gruel, etc.

Stimulants are tea, coffee, whiskey, wine, etc., or hartshorn and water. Of this a teaspoonful in a teacupful of water will be enough for a dose. In making tea or coffee one must not wait to do it as if for the table, but mix hot water and the leaves or grounds, squeeze them well, stir together, and give the whole—leaves, grounds, everything.

Alkaline antidotes which are most likely to be at hand are hartshorn and water a tablespoonful in two teacupfuls of water, soap and water, lime, whiting, soda, chalk, tooth powder, plaster, magnesia, white-wash, and even wood-ashes.

Acid antidotes most commonly accessible are vinegar and lemon-juice.

In giving an antidote it is well to remember that it is not always necessary to wait for it to dissolve, but that it may be stirred up in any fluid at hand (except oil), and swallowed immediately.

TABLE OF DOSES

OF REMEDIES MOST FREQUENTLY ADMINISTERED.

For hypodermic use the dose should be half that used by the mouth.

For use by rectum the dose should be twice that by the mouth.

Doses for Children.—Dr. Young's rule: Add 12 to the age, and divide by the age to give the denominator of a fraction; the numerator of which is 1. Example: For a child two years old, $\frac{12+2}{2} = \frac{1}{2}$ th; the dose should be $\frac{1}{2}$ th that for an adult. In giving powerful medicines and opium still smaller doses must be used for children.

REMEDY.	DOSE.	REMEDY.	DOSE.
Abstract. belladonnae.....	$\frac{1}{2}$ grain.	Ant. et pot. tart.; diaph... emetic	$\frac{1}{2}$ to $\frac{1}{4}$ gr. 1 to 2 grs.
cannab. ind.....	1 grain.	Antimonii oxysulphuret... Antipyrin	$\frac{1}{2}$ to 2 grs. 5 to 30 grs.
coni.....	1 to 3 grains.	Apomorph. hydrochlor.....	$\frac{1}{32}$ to $\frac{1}{8}$ gr.
gelsemii.....	1 grain.	Aqua ammonia.....	5 to 30 mins.
hyoscyami.....	2 grains.	amygd. amar.....	2 fl. drachms.
nuc. vom.....	1 grain.	camphoræ.....	1 to 4 fl. drs.
podophylli.....	4 grains.	chlori.....	1 to 4 fl. drs.
Acet. opi.....	5 minims.	creasoti.....	1 to 4 fl. drs.
scilla.....	10 to 30 mins.	laurocerasi.....	6 to 30 mins.
Acid. acet. dil.....	60 minims.	Argenti nitras.....	$\frac{1}{2}$ to $\frac{1}{4}$ gr.
arsenios.....	$\frac{1}{64}$ to $\frac{1}{2}$ grain.	Arsenii iodidum	$\frac{1}{64}$ to $\frac{1}{16}$ gr.
benzoic.....	5 to 15 grs.	Asafetida.....	5 to 20 grs.
boric.....	5 to 10 grs.	Atropina.....	$\frac{1}{128}$ to $\frac{1}{16}$ gr.
carbolic.....	1 to 3 grs.	Atropinae sulph.....	$\frac{1}{128}$ to $\frac{1}{16}$ gr.
gallie.....	3 to 15 grs.	Auri et sodii chlorid.....	$\frac{1}{32}$ to $\frac{1}{16}$ gr.
hydrobrom. 34 %.....	10 to 15 grs.	Bismuthi citras.....	3 to 15 grs.
hydrobrom. dil.....	15 to 60 mins.	et ammon. citr.....	1 to 15 grs.
hydrochlor.....	3 to 10 mins.	subnitr.....	3 to 15 grs.
hydrochlor. dil.....	10 to 30 mins.	valer.....	1 to 3 grs.
hydrocyan. dil.....	2 to 6 mins.	Caffeina.....	1 to 5 grs.
lactic.....	15 to 60 grs.	Calcii hypophosphis.....	3 to 15 grs.
nitric.....	3 to 10 mins.	iodidum.....	1 to 3 grs.
nitr. dil.....	10 to 30 mins.	phosphas.....	15 to 30 grs.
nitro-hydrochlor.....	3 to 10 mins.	Calx sulphurata.....	$\frac{1}{4}$ to 1 gr.
nitro-hydrochlor. dil.....	5 to 20 mins.	Cambogium.....	1 to 4 grs.
phosphoric (50 %).....	3 to 15 grs.	Camphora.....	3 to 10 grs.
phosphoric dil.....	10 to 20 mins.	monobrom.....	2 to 5 grs.
salicylic.....	5 to 15 grs.	Cantharis.....	$\frac{1}{2}$ to 2 grs.
sulphuric.....	5 to 10 mins.	Capsicum.....	1 to 3 grs.
sulphuric. dil.....	10 to 20 mins.	Castoreum.....	6 to 15 grs.
sulphuric. arom.....	5 to 30 mins.	Catechu.....	15 to 30 grs.
sulphuros.....	30 to 60 mins.	Cerri nitras.....	1 to 3 grs.
tannic.....	2 to 15 grs.	oxalas.....	1 to 3 grs.
Aconitina (white crystals)	$\frac{1}{4}$ to 2 grs.	Chinoidinum.....	3 to 30 grs.
Aloe.....	2 to 5 grs.	Chloral.....	10 to 30 grs.
Aloinum.....	1 to 3 grs.	Chloroformum.....	1 to 5 mins.
Alumen.....	5 to 15 grs.	Cinchona.....	3 to 15 grs.
Ammonii benzoas.....	10 to 20 grs.	Cinchonidina.....	15 to 60 grs.
bromid.....	5 to 30 grs.	Cinchonina.....	1 to 30 grs.
carb.....	3 to 10 grs.	Cinnamomum.....	6 to 30 grs.
chlorid.....	5 to 30 grs.	Cocaine hydrochlorate.....	4 soluti.....
iodid.....	3 to 15 grs.	Codeina.....	$\frac{1}{4}$ to 2 grains.
phosph.....	5 to 20 grs.		
valer.....	3 to 15 grs.		
Amyl nitric.....	2 to 5 mins.		

REMEDY.	DOSE.	REMEDY.	DOSE.
Confectio senna.....	1 to 2 grs.	Extr. cinchona arom. fl.....	30 to 60 mins.
Cardina and its salts.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.	cinchona comp. fl.....	$\frac{1}{2}$ to 1 $\frac{1}{2}$ fl. drs.
Copaiba.....	15 to 60 mins.	cocculi comp.....	1 to 3 mins.
Cot. ma.....	$\frac{1}{2}$ to 1 gr.	colch. rad.....	$\frac{1}{2}$ to 1 $\frac{1}{2}$ grs.
Craeastum.....	1 to 3 mins.	colch. rad. fl.....	3 to 15 mins.
Cratae prepar.....	15 to 75 grs.	colch. sem. fl.....	1 $\frac{1}{2}$ to 10 mins.
Croton chloral.....	1 to 10 grs.	colocynth.....	1 to 5 grs.
Cuba.....	15 to 60 grs.	colocynth. comp.....	1 to 5 grs.
Cupri acetat.....	$\frac{1}{2}$ to 6 grs.	coni fol. fl.....	3 to 15 mins.
sulphas.....	$\frac{1}{2}$ to 10 grs.	convallaria rad. fl.....	15 to 30 mins.
Cuprum ammon.....	$\frac{1}{2}$ to 1 gr.	corn. flor. fl.....	30 to 60 mins.
Curare.....	$\frac{1}{32}$ to $\frac{1}{2}$ gr. ?	coto fl.....	5 to 15 mins.
Decoct. aloes comp.....	$\frac{1}{2}$ to 2 fl. oys	cubeba fl.....	15 to 30 mins.
susp. comp.....	2 to 6 fl. oys	cypripedium fl.....	15 to 60 mins.
Digitalium.....	$\frac{1}{4}$ to $\frac{1}{2}$ gr.	damiana fl.....	$\frac{1}{2}$ to 2 fl. drs.
Digitalis.....	$\frac{1}{2}$ to 3 grs.	digitalis.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.
Duboisina, and its salts.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.	digitalis fl.....	1 to 6 mins.
Elaterium; U.S.P., 1880.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.	drosera fl.....	5 to 10 mins.
Elaterium; U.S.P., 1870.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.	dulcamara.....	5 to 15 grs.
Ergota.....	15 to 60 grs.	dulcamara fl.....	1 to 2 fl. drs.
Ergotinum.....	2 to 8 grs.	ergotæ.....	1 to 8 grs.
Eserina and its salts.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.	ergotæ fl.....	15 to 60 mins.
Euonymin.....	2 to 4 grs.	erythroxyl fl.....	$\frac{1}{2}$ to 2 fl. drs.
Extr. acon. rad.; U.S.P. 1880.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.	eucalypti fl.....	15 to 60 mins.
aconiti [rad.] fluid.....	1 to 5 mins.	euonymi fl.....	15 to 60 mins.
aloes apion.....	$\frac{1}{2}$ to 3 grs.	eupatorii fl.....	30 to 60 mins.
argemone fl.....	15 to 45 mins.	euphorb. ipec. fl.....	5 to 30 mins.
angelica rad. fl.....	30 to 60 mins.	gallæ fl.....	$\frac{1}{2}$ to 2 fl. drs.
anemil.....	2 to 10 grs.	gelsemii.....	2 to 8 mins.
anthemidis fl.....	30 to 60 mins.	gelsemii fl.....	1 to 8 mins.
apocyni cannab. fl.....	8 to 30 mins.	gent. fl.....	30 to 60 mins.
arece fl.....	45 to 75 mins.	gent. comp. fl.....	30 to 60 mins.
arnicae flor.....	3 to 8 grs.	geranii fl.....	15 to 30 mins.
arnicae fl.....	5 to 15 mins.	gillenæ fl.....	15 to 30 mins.
arnicae rad.....	2 to 5 grs.	gossypii fl.....	15 to 45 mins.
arnicae rad. fl.....	5 to 15 mins.	granati rad. cort. fl.....	$\frac{1}{2}$ to 2 fl. drs.
aromat. fl.....	30 to 60 mins.	grind. rob. fl.....	30 to 60 mins.
aspidospermæ fl.....	15 to 45 mins.	grind. squarr. fl.....	30 to 60 mins.
aurantii cort. fl.....	$\frac{1}{2}$ to 2 $\frac{1}{2}$ fl. drs.	guaiaci ligni fl.....	30 to 60 mins.
azedarach fl.....	15 to 75 mins.	guarana fl.....	15 to 30 mins.
baptisiae fl.....	7 to 30 mins.	hamatoxyl.....	8 to 30 grs.
bellad. fol. fl.....	3 to 6 mins.	hamatoxyl fl.....	30 to 60 mins.
bellad. rad.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.	hamamelidis fl.....	60 to 90 mins.
bellad. rad. fl.....	1 to 3 mins.	helleb. nigris.....	$\frac{1}{2}$ to 3 grs.
brayeræ fl.....	2 to 4 fl. drs.	helleb. nigris. fl.....	5 to 15 mins.
bryoniae fl.....	15 to 60 mins.	heloniae fl.....	8 to 30 mins.
buchu fl.....	$\frac{1}{2}$ to 2 $\frac{1}{2}$ fl. drs.	hepatice fl.....	30 to 60 mins.
calami fl.....	15 to 60 mins.	humuli.....	3 to 15 grs.
cand. fl.....	15 to 60 mins.	humuli fl.....	30 to 60 mins.
calumbæ.....	3 to 10 grs.	hydrangeæ fl.....	30 to 60 mins.
calumbæ fl.....	15 to 60 mins.	hydrastis.....	3 to 10 grs.
canelle fl.....	15 to 60 mins.	hydrastis fl.....	8 to 30 mins.
cannab. Amer. fl.....	3 to 15 mins.	hyoscyami fol. fl.....	3 to 30 mins.
cannab. ind.....	$\frac{1}{2}$ to 1 gr.	hyoscyami sem. fl.....	2 to 8 mins.
cannab. ind. fl.....	3 to 6 mins.	iguatiæ.....	$\frac{1}{2}$ to 1 $\frac{1}{2}$ grs.
cautharidis fl.....	1 to 3 mins.	iguatiæ fl.....	1 to 6 mins.
capsici fl.....	1 to 3 mins.	ipecac. fl.....	3 to 60 mins.
cardam. comp. fl.....	15 to 45 mins.	iris versicol.....	3 to 6 grs.
ascarille fl.....	$\frac{1}{2}$ to 2 $\frac{1}{2}$ fl. drs.	iris versicol. fl.....	15 to 30 mins.
caulophylli fl.....	15 to 30 mins.	japura; U.S.P., 1870.....	5 to 10 grs.
chimaph. fl.....	$\frac{1}{2}$ to 1 $\frac{1}{2}$ fl. drs.	jalape fl.....	15 to 60 mins.
cimicifuge fl.....	8 to 30 mins.	juglandis.....	15 to 30 grs.
cinchona.....	15 to 30 grs.	juglandis fl.....	$\frac{1}{2}$ to 2 fl. drs.
cinchona fl.....	30 to 60 mins.	junip. fl.....	30 to 60 mins.

REMEDY.	DOSE.	REMEDY.	DOSE.
Extr. kino fl.....	15 to 30 mins.	Extr. spigellie et sennæ fl.....	$\frac{1}{2}$ to 2 fl. dra.
kramerie fl.....	5 to 15 grs.	stillingie fl.....	$\frac{1}{2}$ to 2 fl. dra.
kramerie fl.....	30 to 60 mins.	stillingie comp. fl.....	$\frac{1}{2}$ to 2 fl. dra.
lactuce.....	5 to 15 grs.	stramonii fol. alc.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.
lactuce fl.....	15 to 60 mins.	stramonii sem.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.
lactucarii fl.....	8 to 30 mins.	stramonii fl.....	1 to 6 mins.
leptandree.....	3 to 10 grs.	sumbul fl.....	15 to 60 mins.
leptandree fl.....	30 to 60 mins.	taraxaci.....	5 to 15 grs.
lobelie fl.....	5 to 30 c.c.	taraxaci fl.....	$\frac{1}{2}$ to 2 fl. dra.
lobelie sem. fl.....	3 to 15 c.c.	trit. rep. fl.....	1 to 4 fl. dra.
lupulini fl.....	10 to 30 mins.	urticæ rad. fl.....	5 to 15 mins.
lycopi fl.....	5 to 30 mins.	ustilag. maid. fl.....	15 to 60 mins.
malti.....	1 to 2 $\frac{1}{2}$ drs.	uvæ ursi fl.....	30 to 60 mins.
matico fl.....	30 to 60 mins.	valerian.....	5 to 15 grs.
mezerai.....	$\frac{1}{2}$ to 1 gr.	valer fl.....	30 to 60 mins.
mezerai fl.....	3 to 10 mins.	veratr. vir. fl.....	2 to 8 mins.
nectandree fl.....	1 to 4 fl. drs.	viburni [prunifol.] fl.....	1 to 4 fl. dra.
nuc. vom. fl.....	$\frac{1}{2}$ to 1 gr.	xanthoxyli cort. fl.....	15 to 30 mins.
nuc. vom. fl.....	1 to 5 mins.	xanthoxyli fruct. fl.....	15 to 30 mins.
opii.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.	zingiberis fl.....	8 to 30 mins.
papaveris.....	$\frac{1}{2}$ to 2 grs.	Fel bovis purif.....	3 to 6 grs.
papaveris fl.....	15 to 45 mins.	Ferri arsen.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.
pareire fl.....	30 to 60 mins.	benzoas.....	1 to 5 grs.
petroselin fl.....	1 to 2 fl. dra.	bromid.....	1 to 5 grs.
physostigma.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.	carb. sacch.....	4 to 15 grs.
physostigma fl.....	1 to 3 mins.	chlorid.....	1 to 3 grs.
phytolacæ baccar. fl.....	5 to 30 mins.	citr.....	5 to 10 grs.
phytolacæ rad. fl.....	1 to 3 grs.	et ammon. citr.....	5 to 10 grs.
phytolacæ rad. fl.....	5 to 30 mins.	et ammon. sulph.....	5 to 10 grs.
pilocarpi fl.....	15 to 60 mins.	et ammon. tartr.....	5 to 15 grs.
piper. nigr. fl.....	15 to 45 mins.	et cinchonid. citr.....	5 to 10 grs.
pisicidie fl.....	15 to 60 mins.	et pot. tartr.....	5 to 30 grs.
podophylli.....	$\frac{1}{2}$ to 1 $\frac{1}{2}$ grs.	et quin. citr.....	5 to 10 grs.
podophylli fl.....	8 to 30 mins.	et strychn. citr.....	1 to 5 grs.
prun. virg. fl.....	30 to 60 mins.	hypophosphis.....	5 to 10 grs.
pulsatillæ fl.....	2 to 10 mins.	iodidum.....	1 to 5 grs.
quassie.....	1 to 5 grs.	iodidum sacch.....	2 to 10 grs.
quassie fl.....	30 to 60 mins.	lactas.....	1 to 3 grs.
quercus fl.....	30 to 60 mins.	oxalas.....	1 to 3 grs.
rhei.....	5 to 15 grs.	oxid. hydrat.....	$\frac{1}{2}$ to 2 grs.
rhei fl.....	15 to 45 mins.	phosphas.....	1 to 5 grs.
rhois arom. fl.....	15 to 60 mins.	pyrophosphas.....	1 to 5 grs.
rhois glabr. cort. fl.....	30 to 60 mins.	subcarb.....	5 to 30 grs.
rhois glabr. fruct. fl.....	30 to 60 mins.	sulphas.....	1 to 3 grs.
rhois toxicod. fl.....	1 to 6 mins.	sulphas exsiccant.....	1 to 3 grs.
ricini fol. fl.....	$\frac{1}{2}$ to 2 fl. drs.	valer.....	1 to 3 grs.
rose fl.....	$\frac{1}{2}$ to 2 fl. drs.	Ferum dialysatum.....	5 to 15 mins.
rubi fl.....	15 to 50 mins.	redactum.....	1 to 5 grs.
sabina fl.....	5 to 15 mins.	Guarana.....	10 to 30 grs.
salicis fl.....	$\frac{1}{2}$ to 2 fl. drs.	Hydrarg. chlorid. corros.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.
salvia fl.....	$\frac{1}{2}$ to 2 fl. drs.	chlorid. mite.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.
sanguin. fl.....	5 to 15 mins.	cyanid.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.
santali citr. fl.....	1 to 2 fl. drs.	iodid. flav.....	$\frac{1}{2}$ to 1 gr.
santonica fl.....	15 to 60 mins.	iodid. rubr.....	$\frac{1}{2}$ to 1 gr.
sarsap. fl.....	$\frac{1}{2}$ to 2 fl. drs.	iodid. vir.....	$\frac{1}{2}$ to 1 gr.
sassafras fl.....	$\frac{1}{2}$ to 2 fl. drs.	oxid. flav.....	$\frac{1}{2}$ to $\frac{1}{2}$ gr.
scilla fl.....	5 to 30 mins.	oxid. nigr.....	1 to 1 gr.
scillæ comp. fl.....	5 to 30 mins.	oxid. rubr.....	1 to 1 gr.
scoparii fl.....	$\frac{1}{2}$ to 1 fl. dr.	subsulphas flav.....	$\frac{1}{2}$ to 1 gr.
senega fl.....	8 to 15 mins.	c. creta.....	1 to 8 grs.
senna fl.....	1 to 4 fl. drs.	Hyosciamina, and salts.....	$\frac{1}{2}$ to 1 gr.
serpent. fl.....	30 to 60 mins.	Infusum digitalis.....	2 to 4 fl. dra.
simarubæ fl.....	15 to 30 mins.	senna comp.....	1 to 2 fl. dra.
spigellie fl.....	15 to 60 mins.	Iodoformum.....	1 to 3 grs.

REMEDY.	DOSE.	REMEDY.	DOSE.
Iodum.....	$\frac{1}{2}$ to $\frac{1}{4}$ gr.	Pepsinum purum.....	15 grains.
Ipecacuanha } expect.....	$\frac{1}{4}$ to 1 gr.	saccharatum.....	30 grains.
} emet.....	15 to 30 grs.	Phosphorus.....	$\frac{1}{12}$ to $\frac{1}{6}$ gr.
Iridin.....	2 to 4 grs.	Physostigminæ salicylas.....	$\frac{1}{12}$ to $\frac{1}{6}$ gr.
Jalapa.....	15 to 30 grs.	sulphas.....	$\frac{1}{12}$ to $\frac{1}{6}$ gr.
Kamala.....	1 to 2 drs.	Picrotoxinum.....	$\frac{1}{2}$ to $\frac{1}{4}$ gr.
Kino.....	8 to 30 grs.	Pilocarpina, and salts.....	$\frac{1}{2}$ to $\frac{1}{4}$ gr.
Lactucarium.....	8 to 15 grs.	Pil. aloes.....	1 to 3 pills.
Liq. ammon. acet.....	2 to 4 fl. drs.	aloes et asafoet.....	2 to 5 pills.
acidi arseniosi.....	2 to 7 mins.	aloes et ferri.....	1 to 3 pills.
arsen. et hydr. iod.....	2 to 7 mins.	aloes et mast.....	1 to 5 pills.
ferri chloridi.....	2 to 10 mins.	aloes et myrrhæ.....	2 to 5 pills.
ferri dialys.....	10 to 15 mins.	antin. comp.....	1 to 5 pills.
ferri nitrat.....	8 to 15 mins.	asafoetida.....	1 to 6 pills.
pepsini.....	1 to 4 fl. drs.	cathart. comp.....	1 to 4 pills.
potassæ.....	5 to 30 mins.	ferri comp.....	2 to 5 pills.
potassii arsenit.....	3 to 7 mins.	ferri iodidi.....	1 to 4 pills.
potassii citrat.....	2 to 4 fl. drs.	opii.....	1 to 2 pills.
sodæ.....	5 to 30 mins.	phosphori.....	1 to 4 pills.
sodii arseniatas.....	3 to 7 mins.	rhei.....	2 to 5 pills.
Lithii benzoas.....	2 to 5 grs.	rhei comp.....	2 to 5 pills.
bromid.....	1 to 3 grs.	Piperinum.....	1 to 8 grs.
carb.....	2 to 6 grs.	Plumbi acetat.....	$\frac{1}{2}$ to 3 grs.
cit.....	2 to 5 grs.	iodidum.....	$\frac{1}{2}$ to 3 grs.
salicylas.....	2 to 8 grs.	Potassii acetat.....	15 to 60 grs.
Lupulinum.....	5 to 10 grs.	bicarb.....	8 to 60 grs.
Magnesia.....	15 to 60 grs.	bitart.....	1 to 2 drs.
Magnesiæ carb.....	15 to 60 grs.	bromid.....	8 to 60 grs.
cit. gran.....	2 to 4 drs.	carb.....	8 to 30 grs.
sulphas.....	2 to 8 drs.	chloras.....	5 to 30 grs.
sulphis.....	8 to 30 grs.	citras.....	15 to 60 grs.
Mangani sulphas.....	2 to 10 grs.	cyanid.....	$\frac{1}{12}$ to $\frac{1}{6}$ gr.
Manna.....	1 to 2 ozs.	et sodii tartr.....	$\frac{1}{2}$ to 1 oz.
Massa copaiabæ.....	5 to 30 grs.	hypophosphis.....	5 to 15 grs.
ferri carb.....	5 to 15 grs.	iodid.....	5 to 60 grs.
hydrarg.....	1 to 10 grs.	nitras.....	8 to 15 grs.
Mist. ammoniac.....	4 to 8 fl. drs.	sulphidum.....	1 to 10 grs.
asafoetida.....	2 to 4 fl. drs.	sulphis.....	15 to 30 grs.
chloroformi.....	2 to 4 fl. drs.	tartras.....	1 to 8 drs.
crete.....	1 to 4 fl. drs.	Pulv. antimonialis.....	3 to 10 grs.
ferri comp.....	$\frac{1}{2}$ to 2 fl. ozs.	aromat.....	5 to 30 grs.
ferri et ammi. acet.....	1 to 4 fl. drs.	crete comp.....	8 to 30 grs.
glycyrrh. comp.....	1 to 4 fl. drs.	glycyrrh. comp.....	30 to 60 grs.
magnes. et asafoet.....	1 to 4 fl. drs.	ipecac. et opii.....	5 to 15 grs.
potassii citr.....	$\frac{1}{2}$ to 1 fl. oz.	jalapæ comp.....	30 to 60 grs.
rhei et sodæ.....	$\frac{1}{2}$ to 1 fl. oz.	morphinæ comp.....	8 to 15 grs.
Morphina and its salts.....	$\frac{1}{4}$ to $\frac{1}{2}$ gr.	rhei comp.....	30 to 60 grs.
Narcocina.....	$\frac{1}{4}$ to 2 grs.	Quinidina (and salts).....	1 to 30 grs.
Nitro-glycerinum.....	$\frac{1}{100}$ to $\frac{1}{50}$ gr.	Quinina (and salts).....	1 to 30 grs.
Nux vomica.....	1 to 5 grs.	Quininæ arsenias.....	$\frac{1}{2}$ to 1 gr.
Oleoresina capsici.....	$\frac{1}{2}$ to $\frac{1}{4}$ gr.	Resina copaiabæ.....	2 to 10 grs.
cubebæ.....	5 to 30 grs.	jalapæ.....	2 to 5 grs.
inupulini.....	2 to 5 grs.	podophylli.....	1 to $\frac{1}{2}$ gr.
pipenis.....	1 to 3 grs.	scammonii.....	2 to 10 grs.
zingiberis.....	1 to 3 grs.	Rheum.....	2 to 30 grs.
Oleum copaiabæ.....	8 to 15 mins.	Salicinum.....	8 to 30 grs.
cubebæ.....	15 to 30 mins.	Santonica.....	8 to 60 grs.
eriger.....	5 to 15 mins.	Santonium.....	1 to 5 grs.
eucalypti.....	10 to 30 mins.	Scammonium.....	5 to 15 grs.
phosphoratum.....	1 to 3 mins.	Senna.....	8 to 60 grs.
sabina.....	1 to 3 mins.	Sodii acetat.....	15 to 60 grs.
terebinth.....	5 to 60 mins.	arsenias.....	$\frac{1}{2}$ to $\frac{1}{4}$ gr.
tiglii.....	1 to 4 drops.	benzoas.....	5 to 15 grs.
Opium (14 % morphine).....	$\frac{1}{4}$ to $\frac{1}{2}$ grs.	bicarb.....	8 to 30 grs.

REMEDY.	DOSE.	REMEDY.	DOSE.
Sodii bisulphis.....	8 to 30 grs.	Tinct. conii.....	5 to 30 mins.
boras.....	8 to 30 grs.	cubebæ.....	1 to 2 fl. dra.
bromid.....	8 to 30 grs.	digitalis.....	5 to 15 mins.
carb.....	8 to 30 grs.	ferri acet.....	15 to 30 mins.
hypophosphis.....	8 to 15 grs.	ferri chloridi.....	15 to 60 mins.
hyposulphis.....	8 to 30 grs.	ferri chloridi æther..	15 to 30 mins.
iodidum.....	5 to 15 grs.	gallæ.....	½ to 2 fl. dra.
phosphas.....	2 to 15 grs.	gelsemii.....	5 to 10 mins.
sahcyas.....	5 to 30 grs.	gualaci.....	30 to 60 mins.
santoninas.....	2 to 10 grs.	guaiaci ammon.....	30 to 60 mins.
sulphas.....	1 to 2 drs.	hellebori.....	10 to 15 mins.
sulphis.....	8 to 30 grs.	humuli.....	1 to 4 fl. dra.
Spir. æther.....	30 to 60 mins.	hydrastis.....	30 to 90 mins.
æther. nitrosi.....	½ to 2 fl. drs.	hyoscyami fol.....	15 to 60 mins.
ammoniac.....	8 to 30 mins.	hyoscyami sem.....	15 to 30 mins.
ammoniac arom.....	15 to 60 mins.	ignatie.....	5 to 15 mins.
camphoræ.....	8 to 30 mins.	iodi.....	5 to 15 mins.
chloroformi.....	15 to 60 mins.	ipecac. et opii.....	5 to 15 mins.
lavend. comp.....	30 to 60 mins.	jalapæ.....	½ to 2 fl. dra.
menth. pip.....	30 to 60 mins.	kino.....	½ to 2 fl. drs.
Strychnina (and salts).....	½ to ½ gr.	kramerie.....	½ to 2 fl. dra.
Sulphur.....	½ to 4 drs.	lavend. comp.....	½ to 2 fl. dra.
Syrupus acidi hydroiodici	1 to 4 fl. drs.	lobeliæ.....	15 to 45 mins.
allii.....	1 to 4 fl. drs.	lupulini.....	½ to 2 fl. dra.
calci lactophos.....	1 to 2 fl. drs.	matico.....	½ to 2 fl. dra.
calcis.....	15 to 30 mins.	nuc. vomicæ.....	5 to 30 mins.
ferri bromidi.....	15 to 60 mins.	opii.....	5 to 25 mins.
ferri iodidi.....	15 to 60 mins.	opii camp.....	1 to 4 fl. dra.
ferri oxidi.....	1 fl. drachm.	physostigmatis.....	5 to 15 mins.
ferri hypophosph.....	1 fl. drachm.	quassia.....	½ to 2 fl. dra.
fer. quin et str. phos.....	1 fl. drachm.	rhei.....	1 to 8 fl. drs.
hypophosphit.....	1 fl. drachm.	rhei arom.....	30 to 75 mins.
ipecac.....	½ to 4 fl. drs.	rhei dulc.....	1 to 4 fl. dra.
krameria.....	½ to 4 fl. dra.	sanguinaria.....	15 to 60 mins.
lactucarii.....	1 to 3 fl. drs.	scillæ.....	8 to 60 mins.
rhei.....	1 to 4 fl. drs.	serpentaria.....	½ to 2 fl. dra.
rhei arom.....	1 to 4 fl. drs.	stramon. fol.....	8 to 15 mins.
rose.....	1 to 2 fl. drs.	stramon sem.....	6 to 15 mins.
rubi.....	1 to 2 fl. drs.	sumbul.....	8 to 30 mins.
sarsap. comp.....	1 to 4 fl. drs.	valer.....	½ to 2 fl. dra.
scillæ.....	½ to 1 fl. dr.	valer. ammon.....	1 to 2 fl. dra.
scillæ comp.....	15 to 60 mins.	veratr. vir.....	1 to 10 mins.
senega.....	1 to 2 fl. drs.	zingiberis.....	15 to 60 mins.
sema.....	1 to 4 fl. drs.	Tritur. elaterii.....	½ to ½ gr.
Tinct. aconiti fol.....	5 to 16 mins.	Veratrina.....	1 to ½ gr.
aconiti rad.....	1 to 5 mins.	Vin. aloes.....	1 to 2 fl. dra.
acon. rad. Fleming's.....	½ to 2½ mins.	antim. { expect. & alt.	1 to 8 mins.
aloes (1880).....	½ to 2 fl. drs.	{ emet.....	30 to 75 mins.
aloes et myrrhæ.....	1 to 2 fl. drs.	colch. rad.....	8 to 45 mins.
arnicæ flor.....	8 to 30 mins.	colch. sem.....	5 to 30 mins.
arnicæ rad.....	15 to 30 mins.	ergotæ.....	1 to 4 fl. dra.
asafoetida.....	30 to 60 mins.	ferri amar.....	1 fl. drachm.
belladonna.....	5 to 15 mins.	ferri citrat.....	1 fl. drachm.
bryonia.....	15 to 30 mins.	ipecac. { expect.....	5 to 15 mins.
calendula.....	15 to 30 mins.	{ emet.....	2 to 4 fl. dra.
calumbæ.....	1 to 4 fl. drs.	opii.....	5 to 10 mins.
cannabis ind.....	15 to 30 mins.	rhei.....	1 to 2 fl. dra.
causticid.....	5 to 15 mins.	Zinc. acet.....	1 to 2 grs.
capsici.....	8 to 15 mins.	bromid.....	1 to 2 grs.
catechu comp.....	½ to 2 fl. drs.	iodid.....	½ to 3 grs.
cimicifugæ.....	30 to 60 mins.	oxid.....	1 to 10 grs.
cinchonæ.....	½ to 2 fl. drs.	phosphid.....	½ to ½ gr.
cinchonæ comp.....	½ to 2 fl. drs.	sulphas, emet.....	15 to 30 grs.
colchici rad.....	5 to 15 mins.	valerianas.....	1 to 6 grs.
colchici sem.....	5 to 15 mins.		

THERAPEUTIC TABLE.

ARRANGED FROM TEXT-BOOK OF PHARMACOLOGY, THERAPEUTICS, AND MATERIA MEDICA.

BY T. LAUDER BRUNTON, M. D.

Abscess.

CALCIC SULPHIDE. Small doses frequently repeated, to hasten maturation or healing.
COD-LIVER OIL. In serofulous cases.
IRON. As a tonic.

Acidity.

ACIDS. Before meals.
ALKALIES. After meals.
CREASOTE.
NUX VOMICA. Before meals.
SILVER NITRATE.

Acne.

ARSENIC. In chronic acne.
BISMUTH. As ointment or powder.
HOT FOMENTATIONS.
HOT SPONGING.
IODIDE OF SULPHUR. In chronic cases.
SULPHUR. Internally and externally.
WATER. Hot sponging several times a day.

After-pains.

BELLADONNA. As ointment.
CAMPHOR. Combined with morphia.
MORPHIA AND ATROPIA. Hypodermically, $\frac{1}{2}$ - $\frac{1}{4}$ gr. morphia with $\frac{1}{16}$ gr. of atropia.
POULTICES, warm, to the hypogastrium.

Alcoholism.

AMMONIA. Aromatic spirits of, as substitute for alcohol, to be taken when the craving comes on.
ARSENIC. To lessen vomiting.
CAPSICUM. As substitute for alcohol.
NUX VOMICA. Nervous and general tonic.
OPIUM.

Alopecia.

ARSENIC. Internally.
CANTHARIDES TINCT. One part to eight of castor oil, rubbed in roots of hair morning and night.

Amenorrhœa.

ALOES. Alone or with iron.
ARSENIC. With iron, in anæmia.
BATHS, hot foot and hip
MANGANESE and permanganate of potassium, j-ij. grs.

Anæmia.

ARSENIC. With, or in place of, iron.
IRON. Chalybeate waters.
MANGANESE. Given with iron.
NUX VOMICA. With iron.
PANCREATIN. In feeble digestion.
PEPSIN. In feeble digestion.

Aneurism.

ALIMENT. Low diet; absolute rest.
ERGOT. Adjunct to rest.
POTASSIC IODIDE. In doses of 30 grs.

Angina Pectoris.

ETHER. To diminish pain.
MORPHIA. Hypodermically.
NITRITE OF AMYL. During paroxysms.
NITRO-GLYCERINE. Begin with one drop of 1 per cent. solution.
TURPENTINE. Locally to chest during paroxysms.

Anus, Fissure of.

BELLADONNA. Locally.
CASTOR OIL. To keep motions soft.
IODIFORM. Locally, to heal and relieve pain.
OPIUM AND GALL OINTMENT. Locally.
SILVER NITRATE.
TANNIN. Locally.

Aphthæ.

BORAX. In honey or glycerine, alone or with chlorate of potash.
MERCURY. Hydr. cum creta, to remove indigestion.
POTASSIC CHLORATE. As wash, 10 grs. to the oz., alone or with borax.
RHUBARB. To remove indigestion.
SULPHUROUS ACID. Locally.

Apoplexy.

ACONITE. To lower blood-pressure where pulse is strong.
 COLD WATER. To head when face congested.
 CROTON OIL. One or two drops on back of tongue.
 ELATERIUM. In suppository or enema during attack.

Ascites.

ARSENIC. In old persons with feeble heart.
 DIGITALIS. In cardiac dropsy.
 ELATERIUM. As hydragogue cathartic.
 JABORANDI. In anasarca.

Asthma.

ACONITE. In spasmodic cases, also in asthma consequent on nasal catarrh in children.
 AMYL NITRITE. To check paroxysm.
 ARSENIC. In small doses in cases associated with bronchitis or hay fever.
 BELLADONNA. Internally, in large doses.
 BROMIDES.
 CHLORAL. During paroxysm.
 CHLOROFORM. Inhaled.
 MORPHIA. Hypodermically.
 NUX VOMICA.
 PEPSIN. To prevent attacks.
 POTASSIC IODIDE. In large doses, when asthma due to acute bronchial catarrh.
 POTASSIC NITRATE. Inhalation of fumes.
 QUINIA. In intervals, when attacks are periodical.
 STRAMONIUM. In cigarettes, or 20 grs. of dried leaves mixed with nitrate of potash, and fumes inhaled.
 TOBACCO. Smoking sometimes beneficial.

Balanitis.

ASTRINGENT LOTIONS. Alum; sulphate of zinc.
 MERCURY. $\frac{3}{4}$ gr. chlor. corrosiv. to an oz. lime-water, as lotion.

Bed Sores.

ALUM. With white of egg, as local application.
 CHARCOAL. Dry, or in poultices.
 IODOFORM.

Biliousness.

ALOES. In constipation.
 MERCURIAL CATHARTICS. In moderate doses night and morning.

Biliousness (continued).

MINERAL ACIDS. Nitro-hydrochloric especially useful.
 PODOPHYLLUM. In place of mercury when stools dark.

Bladder, Irritable.

BELLADONNA.
 BENZOIC ACID. In large prostate and alkaline conditions of urine.
 CANTHARIDES. Not to be used in states of acute inflammation.

Bladder, Paralysis of.

CANNABIS INDICA. In retention, from spinal disease.
 CANTHARIDES. In atonic bladder.
 ERGOT. In paralysis of bladder or sphincter.
 STRYCHNIA.

Blepharitis.

BISMUTH. Equal parts of the subnitrate and glycerine locally.
 MERCURY (UNG. HYDRARG. NITRATIS). If too strong, dilute with vaseline or simple ointment.

Boils.

ARSENIC.
 BELLADONNA. Internally or locally.
 COLLODION. Painted over surface in papular stage, and over base, leaving centre free, in pustular stage.
 OPIUM. Locally to remove pain.
 POULTICES.
 SULPHIDES. In small doses to abort or hasten maturation.

Breath, Fetid.

CAMPHOR.
 PERMANGANATE OF POTASH. As wash to mouth.

Bright's Disease.

BATHS. Hot water and air.
 BROOM. As diuretic.
 CAFFEINE.
 ELATERIUM. As cathartic.
 GALLIC ACID. Lessens albumen.
 JABORANDI.
 LEAD. Lessens albumen.
 NITROUS ETHER.
 POTASSIC B.TARTRATE. Cathartic and diuretic.
 POTASSIC IODIDE.
 TURPENTINE. As diuretic.
 WATER. In large draughts.

Bronchiectasis.

CHLORINE. Inhalation, to lessen fetor.
 IODINE. Inhalation.
 PHOSPHATES AND HYPOPHOSPHITES.
 QUINIA.

Bronchitis, Acute.

ACONITE. $\frac{1}{2}$ -1 m. every hour at commencement.

AMMONIUM CARBONATE. Where much expectoration and depression.

IPECACUANHA. When expectoration scanty with dryness in chest.

NITRIC ACID. When expectoration too copious.

QUINIA. To reduce temperature.

TARTAR EMETIC. In dry stage

TURPENTINE. When expectoration is profuse.

Bronchitis, Capillary.

AMMONIUM CARBONATE. Much expectoration and lividity.

CAMPHOR. As expectorant.

IPECACUANHA. As expectorant and emetic.

MUSTARD. In poultices.

POULTICES. Over whole chest.

Bronchitis, Chronic.

AMMONIA. For difficulty in expectoration

AMMONIACUM. Useful for old people.

ASAFCETIDA.

BALSAM OF PERU, AND BALSAM OF TOLU. When expectoration copious.

CAMPHOR.

COD-LIVER OIL.

DIGITALIS. Where heart is feeble.

IODINE. By inhalation.

IRON.

OPIUM. To lessen secretion and cough.

PHOSPHATES.

SENEGA. Where expulsion is feeble.

SERPENTARIA. Like SENEGA.

SQUILL. Where expectoration is thick.

SULPHUR. Where expectoration is copious.

TAR. To lessen secretion and allay cough.

TURPENTINE.

Bronchorrhœa.

ASAFCETIDA.

EUCALYPTUS.

LEAD ACETATE. To lessen secretion.

Bruises.

ACONITE. Liniment.

CAPSICUM. To remove discoloration of bruise

OPIUM. Locally applied.

Bubo.

ICE.

IODINE.

Bubo (continued).

NITRIC ACID. Locally to indolent bubo.

POTASSIO-TARTRATE OF IRON. Locally and internally.

Burns and Scalds.

BORACIC ACID. Saturated watery solution.

CARBOLIC ACID. One to six of olive oil, locally.

COCAINE.

CREASOTE WATER.

LEAD CARBONATE. In white paint.

LIME. Lime-water with linseed oil.

SODA. Bicarbonate.

WARM BATH. Whole body immersed for days in extensive burns or scalds. It relieves pain, although it may not save life.

Calculi, Biliary.

CHLOROFORM. Inhalation in paroxysm.

COUNTER-IRRITANTS. Mustard poultices.

MORPHIA. $\frac{1}{2}$ gr. with $\frac{1}{16}$ gr. atropia subcutaneously in paroxysm.

SODIUM PHOSPHATE. In 20 or 30 gr. doses before meals as prophylactic.

Calculi, Renal and Vesical.

ALKALIES.

AMMONIUM BENZOATE.

ANÆSTHETICS. To relieve pain.

CHLOROFORM.

MORPHIA. Hypodermically.

Cancer.

BELLADONNA. Locally.

CHARCOAL, poultices, and dry.

CITRIC ACID. As lotion.

IODOFORM. Locally.

OPIUM. Locally and internally.

Cancerum Oris.

ARSENIC. Internally.

NITRIC ACID. Locally.

POTASSIC CHLORATE.

Carbuncle.

BELLADONNA EXTRACT. Locally.

COLLODION.

POULTICES.

STRAPPING. Leaving centre free.

Catarrh, Acute Nasal.

ACONITE. Internally at commencement

ACONITE AND BELLADONNA. In sore throat and cold, one drop of tinct. aconite to two of belladonna every hour.

BELLADONNA. 5 m. of tinct., and afterward one or two every day, for profuse watery secretion.

Catarrh, Acute Nasal (*cont'd.*)

CAMPHOR. As inhalation from steaming water.
 IODINE. By inhalation.
 OPIUM. Dover's powder at commencement.
 QUINIA. 10 gr. of quinia at commencement.

Cerebral Congestion.

ACONITE. In acute cases.
 BELLADONNA.
 BROMIDES.
 CATHARTICS.
 HYDROCYANIC ACID.
 VENESECTION.

Chancre.

ODOFORM. Powdered.

Chancroid.

CAUSTICS. Rarely necessary.
 IODOFORM. Very useful.

Cholera Asiatica.

ALCOHOL. Iced brandy, to stop vomiting.
 CAMPHOR.
 SALINE INJECTIONS.

Cholera Infantum.

BISMUTH.
 FOOD.
 MERCURY. Gray powder or calomel.
 OPIUM.
 POTASSIC BROMIDE. In nervous irritation.
 ZINC OXIDE. With bismuth and pepsin.

Chorea.

ARSENIC.
 CHLORAL.
 CONIUM.
 IRON. In anemia.
 STRYCHNIA.

Colic, Intestinal.

ASAFOETIDA.
 CHLOROFORM.
 ESSENTIAL OILS. ANISEED, CARDAMOM, CINNAMON, CLOVES, PEPPERMINT.
 MORPHIA. Hypodermically.
 POULTICES.

Colic, Lead.

ALUM.
 CASTOR OIL.
 IODIDE OF POTASSIUM.
 MAGNESIUM SULPHATE.

Condylomata.

CHROMIC ACID. Locally.
 MERCURY. Wash with chlorine-water or chlorinated soda, and dust with calomel.

Conjunctivitis.

BELLADONNA. Locally.
 BLISTERS. Behind ear.
 BORIC ACID. Solution.
 CASTOR OIL. Drop in eye.
 MERCURY. Citrine ointment.

Constipation.

ALOES.
 BELLADONNA EXTRACT.
 CASCARA SAGRADA.
 COLOCYNTH COMPOUND EXTRACT.
 LIQUORICE POWDER COMPOUND.
 MERCURY.
 NUX VOMICA. 5-10 ℥ in a glass of cold water before breakfast or before dinner.
 PHYSOSTIGMA. 10 ℥ of tincture along with belladonna and nuxvomica.
 PODOPHYLLUM. In biliousness.
 SALINE WATERS. Friedrichshalle, Hunyadi Janos, or Pullna.
 SENNA.
 WATER. Draught in the morning before breakfast.

Convulsions, Infantile.

ASAFOETIDA. A small dose in an enema arrests convulsions from teething.
 BATHS. Warm, with cold effusions to the head.
 BROMIDE OF POTASSIUM.
 CHLORAL. 5 gr., per mouth or rectum.
 CHLOROFORM. To arrest fit.
 VALERIAN. When due to worms.

Cough.

ACONITE. In throat-cough.
 APOMORPHIA. In bronchitis, with deficient secretion.
 ASAFOETIDA. In the after-cough from habit, and in the sympathetic whooping cough of mothers.
 GELSEMIUM. In convulsive and spasmodic cough.
 HYDROCYANIC ACID. For irritable cough.
 HYOSCYAMUS. In tickling night-coughs.
 PRUNUS VIRGINIANA.
 VALERIAN. In hysterical cough.

Croup.

ALUM. Teaspoonful, with honey or syrup, every $\frac{1}{2}$ or $\frac{1}{3}$ hour until vomiting is induced.
 HOT APPLICATIONS TO NECK.
 IPECACUANHA.
 SULPHUROUS ACID. As spray.

Cystitis.

ALKALIES. When urine is acid.
 BELLADONNA. To allay irritability.

Cystitis (continued).

CANTHARIDES. In small doses.
 OPIUM. By enema or suppository,
 to relieve pain.
 TURPENTINE.

Delirium Tremens.

ALCOHOL. Sometimes necessary.
 BEEF TEA.
 BROMIDES. In large doses.
 CAPSICUM.
 DIGITALIS. In large doses has had
 some success.
 FOOD, nutritious, more to be de-
 pended upon than anything else.

Diabetes Mellitus.

ARSENIC.
 CODEIA. Sometimes requires to be
 pushed to the extent of 18 grs. or
 more per diem.
 GLYCERINE. As remedy, and as food
 in place of sugar.
 SKIM-MILK DIET.

Diarrhea.

ACONITE. In high fever and cutting
 abdominal pains.
 BISMUTH.
 CAMPHOR.
 CASTOR OIL AND OPIUM. At com-
 mencement.
 CATECHU.
 CORROSIVE SUBLIMATE. In small
 doses.
 KINO.
 LEAD ACETATE.
 NUX VOMICA. In chronic cases.
 OPIUM.
 ZINC OXIDE.
 ZINC SULPHATE.

Diphtheria.

ALCOHOL.
 ARSENIC.
 CHLORINE-WATER. Internally and
 locally.
 HYDROCHLORIC ACID. Diluted as
 gargle, or strong as caustic.
 IRON PERCHLORIDE.
 LACTIC ACID. As solvent of false
 membrane.
 MERCURY. Calomel or corrosive sub-
 limate internally.
 PAIN. As solvent of false mem-
 brane.
 POTASSIUM CHLORATE. Internally
 and locally.
 STRYCHNIA.
 SULPHUROUS ACID.

Dropsy.

BROOM.
 BRYONIA. Purgative and diuretic.

Dropsy (continued).

COLCHICUM. In hepatic, cardiac,
 and scarlatinal dropsy.
 DIGITALIS. Especially in cardiac
 dropsy; infusion is the best form.
 ELATERIUM.
 IRON.
 JUNIPER.
 NITROUS ETHER.
 POTASSIUM BITARTRATE. With com-
 pound jalap powder.

Dysentery.

ACONITE.
 BELLADONNA.
 CASTOR OIL. In small doses, with
 opium.
 CORROSIVE SUBLIMATE. In small
 doses in chronic cases.
 HAMAMELIS. Much blood in mo-
 tions.
 IPECACUANHA. In 30-gr. doses on
 empty stomach, with complete
 rest.
 LEAD ACETATE. By mouth, enema,
 or suppository.
 NUX VOMICA.
 OPIUM.
 TANNIN.
 ZINC OXIDE.
 ZINC SULPHATE.

Dysmenorrhœa.

APIOL (Oil of Parsley). In neuralgic
 form.
 ARSENIC. When membranous dis-
 charge.
 BELLADONNA. In neuralgic form.
 CANNABIS INDICA. Very useful.
 CROTON CHLORAL. In neuralgic
 form.
 GELSEMIUM.
 IRON.
 NUX VOMICA.
 OPIUM. 3 to 5 ms. of tincture, alone
 or with 3 or 4 gr. of chloral.

Dyspepsia.

ACIDS. Especially nitro-muriatic
 acid.
 ALKALIES.
 ARSENIC.
 ASAFETIDA.
 BELLADONNA.
 BISMUTH.
 CAPSICUM.
 CHARCOAL.
 CHOLAGOGUES.
 FOOD. Nutritious.
 HYDROCYANIC ACID.
 LACTIC ACID.
 LIME-WATER.
 MAGNESIUM SULPHATE.
 NUX VOMICA.
 PANCREATIN.

Dyspepsia (*continued*).

PEPSIN.
SILVER NITRATE.
TARAXACUM.

Dysuria.

ALKALIES.
BELLADONNA.
CAMPHOR.
CANNABIS INDICA.
CANTHARIDES, TINCTURE.
ERGOT.
GELSEMIUM.
NITROUS ETHER.
OPIUM.

Earache.

ALMOND OIL.
ATROPIA SOLUTION. Locally.
GLYCERINE.
OPIUM.
CARBOLIZED OIL.

Eczema.

ALKALIES. Weak solutions of, as constant dressing.
ARSENIC.
BENZOLIN. Locally.
BISMUTH. Locally.
LEAD SALTS. Locally.
LIME-WATER. Alone or with glycerine or oil, locally.
ZINC OXIDE. Ointment.

Emphysema. See *Asthma and Bronchitis*.**Endocarditis.**

ACONITE. At commencement.
BLISTERS.
OPIUM.

Endometritis.

CARBOLIC ACID.
GLYCERINE.
HOT-WATER INJECTIONS.
IODINE.
NITRIC ACID.

Epididymitis.

ACONITE. In small doses frequently repeated.
MERCURY AND MORPHIA. Locally as oleate.
OINTMENT OF MERCURY AND BELLADONNA.
PULSATILLA. In very small doses along with aconite.

Epilepsy.

ARSENIC.
ASAFOETIDA.
BELLADONNA.
BROMIDES OF POTASSIUM, SODIUM, LITHIUM, AND IRON.
HYDROBROMIC ACID.

Epilepsy (*continued*).

NITRITE OF AMYL. By inhalation.
NITRO-GLYCERINE.
QUINIA.
STRYCHNIA.
VALERIAN.
ZINC OXIDE.

Epistaxis.

ALUM. In powder or solution, locally.
DIGITALIS.
GALLIC ACID.
VINEGAR.

Erysipelas.

ACONITE. At commencement.
IRON. Large doses of tincture of chloride frequently.
QUININE
STARCH and other dry powders.
SULPHUROUS ACID.

Fever. See *Intermittent, Typhoid*, etc.

ACIDS. To allay thirst.
ALCOHOL. Small doses frequently.
ANTIPYRINE.
ARSENIC. In malarious fevers.
BELLADONNA. In eruptive fevers.
CAMPHOR. In adynamic fevers.
DIGITALIS.
SYRUP OF HYDRIODIC ACID.
MERCURY. Small doses at commencement of typhoid or scarlet fever.
SALICIN, } In rheumatic fevers or
SALICYLATE OF SODA, } in hyper-
SALICYLIC ACID, } pyrexia.
TURPENTINE. As stimulant.

Fistula.

PEPPER. The confection, as laxative.

Flatulence.

ALKALIES. Before meals.
ASAFOETIDA.
CARMINATIVES.
CHARCOAL.
CREASOTE.
NUX VOMICA.

Gangrene.

BALSAM OF PERU.
BROMINE.
CHARCOAL. As poultice or dry.
NITRIC ACID.

Gastralgia.

ATROPIA.
BISMUTH.
CHLOROFORM. Two or three drops on sugar.
MORPHIA. Subcutaneously or internally, combined with chloroform.

Gastric Ulcer.

ARSENIC.
BISMUTH.
GALLIC ACID.
LEAD ACETATE. To check hemorrhage.
MILK
MORPHIA.
PEPSIN.
SILVER NITRATE.
SILVER OXIDE.

Gastritis.

ARSENIC.
BISMUTH.
NUX VOMICA.
OPIMUM.
SILVER NITRATE.
SILVER OXIDE.

Gleet.

CANTHARIDES.
IRON.
SANDAL-WOOD OIL. Locally and internally.
TANNIN, GLYCERINE OF. Locally.
TURPENTINE.

Goitre.

IODINE. Internally and locally ointment or tincture, and by injection.
MERCURIC BIODIDE. As ointment.

Gonorrhœa.

ALKALINE salts or waters.
ALUM. As an injection.
CANTHARIDES.
CUBEBES. Alone or mixed with copaiba.
SILVER NITRATE. As injection.
ZINC SALTS. As injection.

Gout.

ACONITE.
ALKALIES.
ARSENIC.
BELLADONNA.
COLCHICUM.
LITHIUM.
VERATRIA. As ointment.

Hay Fever.

ACONITE.
ARSENIC.
IODIDE OF POTASSIUM. Internally and locally.

Headache.

ACONITE. When circulation excited.
AMMONIA. Aromatic spirits $\frac{1}{2}$ -2 dr. doses.

Headache (continued).

BELLADONNA. In frontal headache, especially at menstrual period, or from fatigue.
BICARBONATE OF SODA. With biters before meals in frontal headache.
BROMIDE OF POTASSIUM.
BRYONIA. In bilious headache.
CAMPHOR. Internally and externally.
CANNABIS INDICA. In neuralgic headache.
CHLOROFORM, SPIRITS OF. In nervous headache.
DIGITALIN. $\frac{1}{16}$ of a grain twice a day for congestive hemicrania.
GUARANA.
HEAT. As hot-water bag or poultice to nape of neck.
IODIDE OF POTASSIUM. In rheumatic headache, with tenderness of scalp.
MAGNESIA SULPHATE. For frontal headache, with constipation.
MENTHOL. Locally.
MERCURY. In bilious headache.
NUX VOMICA. In nervous or bilious headache.
SPECTACLES. Where the headache depends on astigmatism.
TEA.
VALERIAN. In nervous and hysterical cases.

Heart, Dilated.

DIGITALIS.
MORPHIA.

Heart, Fatty.

ARSENIC.
BELLADONNA.
IRON.
STRYCHNIA.

Heart, Hypertrophied.

ACONITE. To be used with care when valvular disease is present.
CAMPHOR. In palpitation and dyspnea.
DIGITALIS. In small doses.
VERATRUM VIRIDE.

Heart, Palpitation of.

BELLADONNA. Internally, useful in cardiac strain.
BROMIDE OF POTASSIUM. In fluttering heart.
CAMPHOR.
DIGITALIS.
HYOSCYAMUS. In nervous palpitation.
VALERIAN. In nervous cases with dyspnea.

Hematemesis.

ALUM.
GALLIC ACID.
ICE.
LEAD ACETATE.
SULPHURIC ACID.
TANNIC ACID.
TURPENTINE.

Hematuria.

ALKALINE DIURETICS.
ALUM.
BITARTRATE OF POTASH.
DIGITALIS.
GALLIC ACID.
LEAD ACETATE.
TANNIC ACID.
TURPENTINE.

Hemoptysis.

ALUM.
DIGITALIS.
ERGOT.
GALLIC ACID. Very useful.
LEAD ACETATE.
TURPENTINE.

Hemorrhage, Intestinal.

GALLIC ACID.
LEAD ACETATE.
TANNIC ACID.
TURPENTINE.

Hemorrhage, Post-partum.

ALCOHOL.
DIGITALIS.
ERGOT.
HOT WATER. Injection into uterus.
ICE. To abdomen, uterus, or rectum.
IRON PERCHLORIDE, DILUTED. 1:4
injected into uterus.
PRESSURE OVER UTERUS.

Hemorrhoids.

ALKALINE mineral waters useful.
CARBOLIC ACID. Injection into piles.
ERGOT.
GALLS, OINTMENT. With opium.
IODIFORM. In ointment or suppository.
NUX VOMICA. Very useful.
SENNA. Confection or compound liquorice powder.
SULPHUR.

Hepatitis.

AMMONIUM CHLORIDE.
BRYONIA.
NITRO-HYDROCHLORIC ACID.

Herpes Zoster.

ACONITE AND OPIUM. Locally.
MORPHIA.
OIL OF PEPPERMINT. Locally.

Hiccough.

AMYL NITRITE.
BELLADONNA.
CAMPHOR.
CANNABIS INDICA.
CHLORAL.
CHLOROFORM.
ETHER.
LAUREL-WATER.
MORPHIA. Hypodermically.
NUX VOMICA.

Hydrocephalus, Acute.

BLISTERS. To nape of neck.
BROMIDE OF POTASSIUM.
IODIDE OF POTASSIUM.
LEECHES.
MERCURIC CHLORIDE. Small doses.

Hydrocephalus, Chronic.

BLISTERS.
COD-LIVER OIL.
IODIDE OF IRON.
IODIDE OF POTASSIUM.
MERCURY.

Hydrophobia.

CANNABIS INDICA
CHLOROFORM, and hypodermic of morphia $\frac{1}{4}$ grain.

Hydrothorax.

BLISTERS.
BROOM.
DIGITALIS.
ELATERIUM.
JABORANDI.

Hypochondriasis.

ALCOHOL.
ASAFETIDA.
BROMIDE OF POTASSIUM.
HYOSCYAMUS.
MUSK.

Hysteria.

ALOES. In constipation.
AMMONIA, AROMATIC SPIRITS OF
ASAFETIDA.
BROMIDE OF CAMPHOR.
BROMIDE OF POTASSIUM
CAMPHOR.
IGNATIA.
IRON
SPIRITUS ETHERIS NITROSI
VALERIAN.
ZINC VALERIANATE.

Ichthyosis.

BATHS.
COD-LIVER OIL.
ELM-BARK DECOCTION.
GLYCERINE.

Impetigo.

ARSENIC.
COD-LIVER OIL.

Impetigo (continued).

GUTTA-PERCHA. Locally, in solution.
 HYDROCYANIC ACID. To relieve itching.
 LAUREL-WATER. To relieve itching.
 QUININE.
 SULPHUR. Internally.
 ZINC OINTMENT.

Impotence.

ARSENATE OF IRON.
 CANTHARIDES.
 NUX VOMICA.
 PHOSPHORUS.
 ZINC PHOSPHATE.

Influenza.

OPUM. With ipecacuanha.
 POTASSIUM NITRATE. Freely diluted, as limonade.
 QUININE. In later stages.
 SPIRITUS ETHERIS NITROSI.

Insomnia.

ACONITE. 1 m. every quarter of an hour, when skin dry and harsh.
 ALCOHOL.
 CAMPHOR.
 CANNABIS INDICA. Alone, or with hyoscyamines.
 CHLORAL.
 CHLORAL CHLORAL. If heart weak.
 DIGITALIS. Which deranges the cardiovascular system.
 FOOD at bed time.
 GELSEMIUM. In simple wakefulness.
 HYOSCYAMUS. Alone, or with cannabis indica.
 PARALDEHYDE.
 POTASSIUM BROMIDE. In full doses.
 WARMTH. Internally and externally.
 WET COMPRESS.

Intermittent Fever.

ARSENIC.
 ATROPIA. Subcutaneously, to arrest or cut short cold stage.
 CAMPHOR.
 CASCARA. With quinine.
 CHLOROFORM. To prevent or cut short cold stage.
 GELSEMIUM. Pushed until it produces dilated pupils or double vision.
 IODINE TINCTURE. To prevent recurrence of ague.
 OPUM. In full doses, to prevent chill.
 PEPPER. With quinine.
 QUININE.
 QUININE BROMOHYDRATE. Like quinine, and less liable to produce chlorosis.

Intermittent Fever (continued).

SODIUM CHLORIDE. Tablespoonful in glass of hot water at a draught on empty stomach.

Intertrigo.

BISMUTH. Locally.
 FULLER'S EARTH.
 LEAD LOTION.
 LIME-WATER.

Iritis.

ATROPIA. Locally.
 BELLADONNA. Internally and locally.
 DUBOISIA. Substitute for atropia.
 MERCURY. Most serviceable.

Jaundice.

ALKALINE mineral waters in catarrh of duodenum or bile-ducts.
 AMMONIUM CHLORIDE. In scruple doses.
 CALOMEL. Often very useful.
 EUONYMIN.
 IRIDIN.
 MAGNESIUM SULPHATE.
 NITRO-HYDROCHLORIC ACID.
 PODOPHYLLUM.
 QUININE. In malarial cases.

Lactation, Defective.

CASTOR-OIL LEAVES. Poultice or infusion, to breast.
 GOSSYPIUM. Decoction of seeds two or three times every hour.
 JABORANDA.

Lactation, Excessive.

BANDAGE.
 BELLADONNA. Internally and locally.
 CAMPHOR AND GLYCERINE. Locally.
 IODIDE OF POTASSIUM.

Laryngismus Stridulus.

ACONITE.
 BELLADONNA.
 BROMIDES.
 CHLORAL.
 CHLOROFORM. By inhalation.
 COLLAPSONG.
 EMETICS.
 MORPHIA. Hypodermically.
 QUININE.

Laryngitis, Acute.

ACONITE.
 MORPHIA.
 QUININE.
 SCARIFICATION OF LARYNX.
 SULPHUROUS ACID. As inhalation or spray.
 ZINC SULPHATE. Astringent.

Laryngitis, Chronic.

ALUM. As gargle.
 AMMONIUM CHLORIDE. As spray.
 BISMUTH. Locally.
 CARBOLIC ACID. As spray.
 FERRIC CHLORIDE. Locally.
 SILVER NITRATE. Solution locally.
 TANNIN. As gargle or spray.

Leucorrhœa.

ALUM. By injection.
 BALSAM OF PERU. Internally
 BELLADONNA. Locally.
 BORAX. By injection.
 GLYCERINE.
 IODOFORM. Locally.
 IRON. Internally
 POTASSIUM PERMANGANATE. Locally.
 TANNIN. By injection or suppository.

Locomotor Ataxia.

CANNABIS INDICA.
 ELECTRICITY.
 ERGOT.
 PHOSPHORUS.
 POTASSIUM IODIDE.
 STRYCHNIA.

Lumbago.

ACONITE. Internally and locally.
 CHLOROFORM, liniment.
 HOT DOUCHE. Or poultice.
 ICE.
 IODIDE OF POTASSIUM.
 MORPHIA.
 QUININE.

Lupus.

ARSENIC.
 IODIDE OF POTASSIUM.
 IODIDE OF SULPHUR.
 IODOFORM.
 PYROGALLIC ACID.
 SODIUM ETHYLATE.
 ZINC CHLORIDE.

Mania.

CANNABIS INDICA.
 CHLORAL. As narcotic and carminative.
 DIGITALIS.
 HYOSCYAMUS.
 HYOSCYAMINE.
 IRON.
 OPIUM.
 PARALDEHYDE.
 PHYSOSTIGMA.
 WET PACK.

Mastitis.

AMMONIUM CHLORIDE. Locally.
 BELLADONNA. Locally.

Mastitis (continued).

MERCURY AND MORPHIA OLEATE
 Locally.
 SUPPORT, by bandaging or strapping.

Measles.

ACONITE.
 DIGITALIS.
 POTASSIUM BROMIDE. For sleeplessness.
 QUININE.

Melancholia.

ARSENIC.
 BROMIDE OF POTASSIUM.
 CANNABIS INDICA.
 CHLORAL.
 IRON.
 OPIUM.
 VALERIAN.

Meningitis, Cerebral.

ACONITE.
 BELLADONNA.
 BLISTERS. To nape of neck.
 BROMIDE OF POTASSIUM.
 COLD, to head.
 HYOSCYAMUS.
 MERCURY.
 OPIUM. In small doses, alone or with tartar emetic.
 POTASSIUM IODIDE.
 PURGATIVES.
 VENESECTION.

Meningitis, Cerebro-Spinal.

ACONITE.
 BELLADONNA.
 CAUTERY, to back.
 COLD, to spine.
 DIGITALIS.
 ERGOT.
 OPIUM.
 QUININE.
 VENESECTION.

Menorrhagia.

ARSENIC.
 BROMIDE OF POTASSIUM.
 CANNABIS INDICA.
 DIGITALIS.
 ERGOT.
 GALLIC ACID.
 HOT-WATER BAG to dorsal and lumbar vertebrae.
 IRON.
 QUININE.
 SULPHURIC ACID.

Metritis, Acute.

ACONITE.
 HOT WATER. Vaginal injection.
 OPIUM. By suppository or enema.
 POULTICES.
 TURPENTINE STUPES.

Myalgia.

AMMONIUM CHLORIDE.
ARNICA. Internally and locally.
BELLADONNA LINIMENT. Locally and internally.
CHLOROFORM LINIMENT.
FRICTION.
GELSEMIUM. Large doses.
OPIUM.
PACKING.
POTASSIUM IODIDE.
SALICYLATE OF SODA.

Myelitis.

BELLADONNA.
ERGOT.
PHOSPHORUS.
SILVER NITRATE.

Nævus.

CHROMIC ACID.
ELECTROLYSIS.
GALVANO-CAUTERY.
LIGATION.
NITRIC ACID.

Nausea.

BISMUTH.
CLOVES.
HYDROCYANIC ACID.
ICE.
IPECACUANHA. In sickness of pregnancy and chronic alcoholism; 1 ℥ of wine, repeated frequently.
MORPHIA.
PEPPERMINT.

Nephritis, Acute.

ACONITE. At commencement.
BELLADONNA.
CANNABIS INDICA. As diuretic, especially in hæmaturia.
CANTHARIDES. 1 ℥ of tincture every three hours to stop hæmaturia after acute symptoms have subsided.
DIGITALIS. As diuretic.
EUCALYPTUS.
GALLIC ACID.
JABORANDI.
JUNIPER.
POULTICES. Over loins.
TURPENTINE. $\frac{1}{2}$ -1 ℥ doses, every two to four hours.

Nervousness.

ACONITE. 1 ℥ of tincture at bedtime for restlessness.
BROMIDE OF POTASSIUM.
CAFFEINE.
CAMPHOR.
CHLORAL.
IGNATIA.
STRYCHNINE.

Neuralgia.

ACONITIA.
AMMONIUM VALERIANATE.
AMYL NITRITE.
ATROPINE. In liniment or hypodermically.
BELLADONNA.
BLISTERS.
BROMIDE OF POTASSIUM.
CAFFEINE.
CANNABIS INDICA.
CHLORAL AND CAMPHOR.
CHLOROFORM. Locally, and by inhalation, when pain very severe.
COD-LIVER OIL.
CROTON CHLORAL.
DIGITALIS.
ELECTRICITY.
ERGOT.
GELSEMIUM.
IODIDES.
IRON.
MORPHIA. Hypodermically.
MUSTARD POULTICE.
NITRO-GLYCERINE.
NUX VOMICA.
OIL OF CLOVES.
PEPPERMINT.
PULSATILLA.
QUININE.
SALICYLIC ACID.
THERMO-CAUTERY.
VALERIAN.
VIBRATION.
WATER. Hot or cold, locally.

Nightmare.

BROMIDE OF POTASSIUM.
CAMPHOR-WATER.

Nipples, Sore.

BALSAM OF PERU.
BENZOIN.
BORAX.
COLLODION.
LEAD SALTS.
TANNIN.

Nodes.

MERCURY.
POTASSIUM IODIDE.

Nymphomania.

BROMIDE OF POTASSIUM.
CAMPHOR.
LUPULINE.

Obesity.

ALKALINE WATERS.
AMMONIUM BROMIDE.
BANTING'S SYSTEM—living on meat and green vegetables, and avoiding starch, sugars, and fats.

Onychia.

IODOFORM.
 POTASSIUM PERMANGANATE.
 STRAPPING.

Ophthalmia (vide Conjunctivitis).**Orchitis.**

MERCURY AND BELLADONNA. Ointment.
 STRAPPING.

Otitis.

BORACIC ACID. Locally.
 DRY CLEANSING.
 SILVER NITRATE. Solution.
 ZINC SULPHATE.

Ozæna.

ALKALINE WASHES.
 IODINE. As inhalation: washing out nose with solution of common salt, to which a few drops of the tincture of iodine have been added.
 IODOFORM
 POTASSIUM PERMANGANATE. Solution
 TANNIN. GLYCERINE OF.

Parotitis.

ACONITE.
 JABORANDI.
 MERCURY $\frac{1}{4}$ gr. of gray powder three or four times a day.
 POULTICE.

Pediculi.

MERCURY. As ointment or wash.

Pemphigus.

ARSENIC.
 COD-LIVER OIL.
 IODIDE OF POTASSIUM.
 MERCURY.
 PHOSPHORUS.

Pericarditis.

ACONITE.
 BLISTERS. Near heart.
 BRYONIA.
 DIGITALIS. Heart rapid and feeble, with cyanosis and dropsy.
 OPIUM. In grain doses every three to six hours.
 QUININE.
 VERATRUM VIRIDE.

Periostitis.

IODIDE OF POTASSIUM, or AMMONIUM.
 IODINE. Locally.
 POULTICES.

Peritonitis.

ACONITE. At commencement.
 BRYONIA. When exudation.
 LEECHES.
 MERCURY.
 OPIUM. Freely used.
 POULTICES.
 TURPENTINE.

Perspiration.

AROMATIC SULPHURIC ACID.
 ATROPIA.
 BELLADONNA. In liniment, for local sweats.
 JABORANDI.
 OPIUM.
 QUININE
 SPONGING, very hot.
 STRYCHNINE.
 VINEGAR.

Pertussis.

ACONITE.
 ALUM.
 BELLADONNA.
 BROMIDE OF POTASSIUM.
 CHLORAL.
 CHLOROFORM. By inhalation
 COD-LIVER OIL.
 CROTON CHLORAL.
 ERGOT.
 IPECACUANHA.
 MONOBROMIDE OF CAMPHOR.
 OPIUM.
 VALERIAN.

Pharyngitis.

ACONITE.
 ALUM. Gargle.
 CAPSICUM. Gargle.
 FERRIC CHLORIDE. Locally and internally.
 GLYCERINE. Locally.
 POTASSIUM CHLORATE. Locally
 QUININE.
 SILVER NITRATE. Locally.
 STRYCHNINE.
 TANNIN. As powder, or glycerine, locally.
 ZINC SULPHATE. Gargle.

Phlebitis.

HAMAMELIS.
 HOT FOMENTATIONS.
 OPIUM.
 REST, absolute.

Phlegmasia Alba.

AMMONIUM CARBONATE. In full doses, when much prostration.
 BELLADONNA EXTRACT. With mercurial ointment locally.
 HAMAMELIS.
 OPIUM. Internally and locally.

Phthisis.

ALCOHOL. With food or cod-liver oil.
 ANTIPIRYNE. To reduce temperature.
 ARSENIC. To remove commencing consolidation, and when tongue is red and irritable.
 ATRAPIA. To check perspiration.
 BENZOIN. As inhalation to lessen cough and expectoration.
 CHLORAL.
 CHLOROFORM, to check cough.
 COCAINE. Solution 4-20 per cent. to throat and mouth to relieve irritation.
 COD-LIVER OIL.
 DIGITALIS.
 GLYCERINE. In place of cod-liver oil.
 HYPOPHOSPHITES.
 IODINE LINIMENT.
 IRON.
 MERCURIC CHLORIDE. In minute doses for diarrhoea.
 MORPHIA, WITH STARCH OR BIS-MUTH. To larynx.
 OPIUM.
 OXYGEN.
 PHOSPHATE OF LIME.
 PILOCARPINE. To check sweats.
 QUININE.
 SALICYLIC ACID. For foul breath and offensive expectoration.

Pleurisy.

ACONITE.
 BELLADONNA PLASTER.
 BLISTERS.
 BRYONIA. After aconite.
 DIGITALIS.
 IODIDE OF POTASSIUM. To aid absorption.
 LOCAL WET PACK.

Pleurodynia.

BELLADONNA. Plaster or liniment.
 CHLORAL. With camphor locally.
 ETHER. Locally, in spray.
 IRON.
 MUSTARD LEAVES.
 OPIUM.
 POULTICES.
 QUININE.
 STRAPPING.

Pneumonia.

ACONITE. At commencement.
 AMMONIUM CARBONATE. As stimulant toward end.
 BELLADONNA. At commencement.
 BRYONIA. When pleurisy present.
 DIGITALIS.
 OPIUM.

Pneumonia (continued).

POULTICES.
 QUININE.
 SALICYLATE OF SODA. As antipyretic.
 TURPENTINE. As stimulant.

Polypus.

ALUM.
 TANNIN. By insufflation.

Prolapsus Ani.

ALUM. Locally.
 I.C.
 CATERIZATION.
 STRYCHNIA.
 TANNIN. Locally.

Prolapsus Uteri.

ALUM. By vaginal douche.
 GATES. Decoction by injection.
 TANNIN.

Prostate, Enlarged.

ALKALIES. For irritation of bladder, with acid urine.
 AMMONIUM BENZOATE. For cystitis with alkaline urine.
 CONIUM.
 ERGOT.
 IODOFORM. By suppository.

Prurigo and Pruritus.

ARSENIC. Internally.
 BALSAM OF PERU.
 BORACIC ACID.
 CALOMEL. Ointment.
 CHLORAL AND CAMPHOR.
 COD-LIVER OIL. By inunction.
 CORROSIVE SUBLIMATE. Wash.
 GLYCERINE.
 GLYCEROLE OF TAR.
 HYDROCYANIC ACID. Locally.
 IODOFORM. Ointment.
 OPIUM.
 OIL OF CADE.
 TAR OINTMENT.

Psoriasis.

ALKALINE BATH.
 ARSENIC.
 CALOMEL. Ointment.
 CHRYSOPHANIC ACID.
 COD-LIVER OIL.
 GLYCERINE.
 INDIA-RUBBER SOLUTION.
 LEAD IODIDE. Locally.
 PHOSPHORUS. Substitute for arsenic.
 POTASSIUM IODIDE.
 SULPHUR. Internally.
 TAR. Ointment.
 UREUS.
 VASELINE.

Ptyalism.

ACIDS. Small doses, and in gargles.
 BELLADONNA
 CHLORATE OF POTASH. Gargle.
 TANNIN.

Puerperal Convulsions.

BELLADONNA.
 BLEEDING.
 CHLORAL. In full doses.
 CHLOROFORM. By inhalation.
 MORPHIA. Hypodermically.
 VERATRUM VIRIDE. Pushed to nausea.

Puerperal Fever.

ACONITE. At commencement.
 OPIUM. For wakefulness and delirium, very useful.
 QUININE. In large doses.
 TURPENTINE
 VERATRUM VIRIDE.

Puerperal Mania.

ANÆSTHETICS.
 BROMIDE OF POTASSIUM.
 CHLORAL.
 HYOSCYAMUS.
 OPIUM.
 QUININE.

Puerperal Peritonitis.

ACONITE.
 LAXATIVES.
 OPIUM.
 QUININE.
 TURPENTINE.

Purpura Hæmorrhagica.

ERGOT.
 GALLIC ACID.
 IRON.
 NUX VOMICA.
 QUININE.
 TURPENTINE.

Pyæmia.

IRON.
 OPIUM.
 PERMANGANATE OF POTASH.
 QUININE.
 TURPENTINE.

Pyrosis.

ACIDS, MINERAL.
 ALKALIES.
 BISMUTH.
 CAMPHOR.
 CARBOLIC ACID.
 CREASOTE.
 GALLIC ACID.
 GLYCERINE.
 NUX VOMICA.

Rheumatism, Acute.

ACONITE.
 ALKALIES.
 AMMONIUM BROMIDE.
 BLISTERS.
 COLD BATHS.
 DIGITALIS.
 IRON.
 LIME-JUICE.
 OPIUM.
 QUININE.
 SALICYLATE OF SODA. Relieves pain most quickly.

Rheumatism, Chronic.

BELLADONNA.
 BROMIDE OF LITHIUM.
 COD-LIVER OIL.
 GUAIAC.
 IODIDE OF POTASSIUM.
 MERCURY AND MORPHIA. Oleate locally.
 QUININE.
 SULPHUR. Locally and as sulphides, or sulphur waters internally.
 TURKISH BATHS.

Rickets.

CALCIUM PHOSPHATE.
 COD-LIVER OIL.
 COLD SPONGING.
 IODIDE OF IRON.
 QUININE.

Scabies.

BAKING of clothes to destroy ova.
 BALSAM OF PERU. Locally, agreeable and effective.
 MERCURY. White precipitate ointment.
 PETROLEUM.
 SULPHUR. Ointment.

Scarlet Fever. (Treatment chiefly symptomatic.)

ACONITE.
 BELLADONNA.
 CHLORINE-WATER. As gargle.
 COLD COMPRESS to throat.
 DIGITALIS.
 FAT, inunction over the whole body.
 PACKING. Useful and comforting.
 SALICYLATE OF SODA. As antipyretic.
 SPONGING.

Sciatica.

ACUPUNCTURE.
 ATROPIA.
 BELLADONNA.
 BLISTERS.
 CAUTERY. Slight application.
 CHLOROFORM. Liniment.
 ELECTRICITY.
 ETHER. Spray.

Sciatica (continued).

MORPHIA.
POULTICES.
TURPENTINE. In $\frac{1}{4}$ -oz. doses for 3 or 4 nights.

Scrofula.

CALCIUM PHOSPHATE.
COD-LIVER OIL.
IODIDE OF IRON.

Scurvy.

ACIDS.
ALUM. With myrrh, for ulcerated gums.
CITRIC ACID.
LIME-JUICE.
QUINA. With mineral acids.
VINEGAR.

Sea-Sickness.

BROMIDE OF POTASSIUM.
BROMIDE OF SODIUM.
CANNABIS INDICA.
CHAMPAGNE.
CHLORAL.
CHLOROFORM. 2-5 ms. on sugar
COCAINE.
NITRO-GLYCERINE.
NUX VOMICA.

Seborrhoea.

BORAX, with GLYCERINE.

Spermatorrhoea.

ARSENIC.
BELLADONNA.
BROMIDES.
CAMPHOR.
CANTHARIDES.
ELECTRICITY.
NUX VOMICA.
PHOSPHORUS.
TURPENTINE.

Spinal Congestion.

ACONITE.
ERGOT.
NUX VOMICA.

Spinal Irritation.

ACONITE OINTMENT.
ARSENIC.
BELLADONNA.
BLISTERS.
CONIUM.
COUNTER-IRRITATION.
DIGITALIS.
ERGOT.
NUX VOMICA.
OPIUM.
PHOSPHORUS.

Sprains.

AMMONIUM CHLORIDE.
ARNICA.
BANDAGING.
FOMENTATIONS.
MASSAGE.

Stomatitis.

ACIDS, MINERAL.
ALUM.
BORAX.
CHLORATE OF POTASSIUM.
GLYCERINE OF TANNIN.
NITRIC ACID.

Sweating, vide Perspiration.**Sycosis.**

CORROSIVE SUBLIMATE, solution 2-3 grs. to $\frac{1}{3}$ of water.
EPILATION.
SULPHUROUS ACID.

Synovitis.

ACONITE.
ARNICA.
BANDAGE, OR STRAPPING.
BLISTERS.
COD-LIVER OIL.
IODINE. Injection.
MASSAGE.

Syphilis.

ARSENIC.
BITTER TONICS.
COD-LIVER OIL.
IODIDES.
IODOFORM.
IRON.
MERCURY.
POTASSIC IODIDE.

Tabes Mesenterica.

CHLORIDE OF LIME.
COD-LIVER OIL.
FATTY INUNCTION.
PHOSPHORIC ACID.
SARSAPARILLA.

Tetanus.

BROMIDES.
CHLORAL.
GELSEMIUM.
NEUROTOMY.
OPIUM.
PHYSOSTIGMA.
WARM BATHS.

Tic Douloureux.

ACONITIN.
AMMONII CHLORIDUM.
ARSENIC.
CANNABIS INDICA.
MORPHIA.
PHYSOSTIGMA.
VERATRIA.

Tinea Circinata.

COD-LIVER OIL.
 IODINE, TINCTURE. Locally.
 MERCURY. Locally.
 SULPHUROUS ACID. Locally.

Tinea Favosa.

ARSENIC.
 EPILATION.
 HYPOSULPHITES
 MERCURY. Locally.
 SULPHUROUS ACID.

Tinea Tonsurans.

COD-LIVER OIL.
 EPILATION.
 MERCURY OLEATE.
 PARASITICIDE LOTIONS.

Tonsillitis.

ACONITE.
 ALUM.
 BELLADONNA.
 BICARBONATE OF SODA. Dry, locally.
 COLD WET COMPRESS.
 GUAIAC.
 ICE AND WET PACK.
 MERCURY.
 OPIUM.
 POTASSIC IODIDE.
 TANNIC ACID.

Tonsils, Enlarged.

AMMONIUM IODIDE.
 IRON, tincture of chloride.
 MASSAGE.
 SILVER NITRATE.
 TANNIN.

Tonsils, Ulcerated.

BICARBONATE OF SODA. Dry locally.
 IRON, tincture.
 MERCURIC IODIDE.
 POTASSIUM CHLORATE.
 SULPHUROUS ACID.

Toothache.

BICARBONATE OF SODA.
 CARBOLIC ACID.
 CLOVES, OIL.
 CREASOTE.

Torticollis.

ARSENIC.
 ELECTRICITY.
 MASSAGE.
 MERCURY.
 NUX VOMICA.

Trismus.

ANÆSTHETICS.
 BELLADONNA.
 CHLORAL.
 OPIUM.
 PHYSOSTIGMA.

Tympanites.

ACIDS AND ALKALIES.
 ASAFOETIDA.
 CARBOLIC ACID.
 HYOSCYAMUS.
 NUX VOMICA.
 TURPENTINE.
 VEGETABLE CHARCOAL.

Typhlitis.

BELLADONNA.
 ICE-BAG.
 LEECHES.
 MERCURY.
 OPIUM.

Typhoid Fever.

ALCOHOL.
 ANTIPYRETICS.
 ANTIPYRIN.
 ASAFOETIDA.
 BATHS.
 CALOMEL.
 CAMPHOR.
 CARBONATE OF AMMONIA.
 COLD COMPRESS.
 DIGITALIS.
 HYDROCHLORIC ACID.
 HYOSCYAMUS.
 MURIATIC ACID.
 OPIUM.
 QUININE.
 TURPENTINE.

Typhus Fever.

ALCOHOL.
 CAMPHOR.
 DIGITALIS.
 HYOSCYAMUS.
 MINERAL ACIDS.
 OPIUM.
 QUININE.
 STRYCHNIA.

Vomiting.

AMMONIUM SALTS AND AROMATIC
 SPIRITS.
 ARSENIC.
 BELLADONNA.
 BISMUTH.
 BLISTERS.
 BROMIDES.
 CALOMEL.
 CARBOLIC ACID.
 CERIUM OXALATE.
 CHLOROFORM.
 CINNAMON.
 CLOVES.
 CREASOTE.
 COUNTER-IRRITATION.
 HYDROCYANIC ACID.
 ICE.
 IPECACUANHA.
 KOUMISS.
 MORPHIA.

Vomiting (continued).

NUTRIENT ENEMATA.
 NUX VOMICA.
 OPIUM.
 PEPSIN.
 STRYCHNIA.
 ZINC.

Warts.

ACIDS, MINERAL.
 ALKALIES, CAUSTIC.
 CHROMIC ACID.
 SILVER NITRATE.

Worms.

ARTEM. Injection for thread-worms.
 CASTOR OIL.
 EUCALYPTOL. Injection for thread-worms.
 PILIN MAS. For tape-worms.
 IGNATIA.
 IRON. Injection for thread-worms, and as a tonic.

Worms (continued).

KAMALA. For tape-worms.
 KOUSSO. For tape-worms.
 LIME-WATER. Injection for thread-worms.
 PEPO.
 QUINIA.
 QUASSIA. Injection for thread-worms.
 SANTONINE. For round-worms
 SODIUM CHLORIDE. Injection for thread-worms.
 TURPENTINE. For tape-worms

Yellow Fever.

CALOMEL.
 CAMPHOR.
 CHLORODYNE.
 IPECACUANHA.
 LEAD ACETATE.
 NUX VOMICA.
 QUININE.
 STIMULANTS.
 TURPENTINE.

LIGATION OF ARTERIES.

The accompanying plate indicates the positions in which the operation of ligating an artery in its continuity is most commonly performed. There is no point at which ligatures are applied which is not represented, with the exception of a few which are very rarely chosen. With the explanations which follow, this diagram will serve as a guide to the practitioner who may be called upon in an emergency to perform an operation in regard to which he has grown somewhat rusty from want of practice.

1. Ligation of the Temporal Artery.—The temporal artery is the continuation of the external carotid from the level of the lobe of the ear to the level of the eyebrow, where it divides into its terminal branches. The point at which it is ligated is a little in front of the anterior curve of the concha. A vertical incision, about an inch in length, is made in front of the concha. The artery is easy to find, as it is very superficial at this part of its course. Its accompanying vein lies to the outer side, and the temporal branch of the auriculo-temporal nerve lies in front of it.

2. Ligation of the Occipital Artery. The occipital artery comes off from the posterior part of the external carotid at about the level of the angle of the lower jaw. It curves backward under and to the inner side of the mastoid process, at the back of which it comes out through the attachment of the trapezius muscle, and winds in a tortuous course upward over the occipital bone. The point at which it is ligated is about an inch behind the ear. An incision an inch and a half long, parallel to the posterior edge of the mastoid process, will expose it. Its accompanying vein lies to the outer side of it. There are but few occasions when it is necessary to ligate this artery, as its position makes it very easy to stop the flow of blood through it by pressure with a suitable compress and bandage.

3. Ligation of the Facial Artery. The facial artery comes off from the external carotid just below the level of the angle of the lower jaw, and passes forward below this bone to a point just in front of the insertion of the masseter muscle. Here it sends a branch forward, the submental, while the main trunk curves up over the lower jaw and passes in a


tortuous course toward the angle of the mouth. The point at which it is ligated is that at which it ascends in front of the masseter muscle. An incision an inch in length will expose it. Its accompanying vein lies to the outer side.

4. Ligation of the Lingual Artery. The lingual artery comes off from the external carotid at the level of the great corn of the hyoid bone; it passes forward under the posterior belly of the digastric muscle, and the stylo-glossus and hyo-glossus muscles, to the root of the tongue and thence to its further distribution. It is in some respects the most difficult artery to ligate in the whole body. The point at which it is ligated is the second part of its course, under the hyo-glossus muscle. An incision is made about half an inch above, and parallel to, the great horn of the hyoid bone, and extending from just in front of the anterior edge of the sterno-cleido-mastoid muscle to near the middle line of the neck, half an inch below the base of the jaw. The skin and subcutaneous fat having been divided, the capsule of the submaxillary gland will be met. This capsule must be divided, and the gland carefully drawn up over the jaw. In doing this great care must be taken not to injure the facial artery and vein, which pass through the gland. The posterior part of the capsule of the gland must next be divided. This will expose the shining aponeurosis which constitutes the pulley of the digastric muscle, and attaches it to the great horn of the hyoid bone, near the insertion of the stylo-hyoid muscle. Behind these passes the hypoglossal nerve. This must be delicately detached from the sheath of the hyo-glossus muscle, which lies below it, and pushed up out of the way. As the next step, fix the pulley of the digastric with a tenaculum and gently insinuate a director behind the posterior edge of the hyo-glossus muscle, and push it forward between this and the middle constrictor of the pharynx, which lies underneath it, in a line parallel to, and just above, the greater horn of the hyoid bone. The hyo-glossus muscle is then to be divided on the director, and the lingual artery is exposed. In placing the ligature the needle should be passed from above downward, so as not to injure the hypoglossal nerve. The important guides in this ligation are the hyoid bone, the pulley of the digastric muscle, and the hypoglossal nerve.

5 and 6. Ligation of the Common Carotid Artery.—For ligation, the common carotid artery is divided into two portions by the narrow omo-hyoid muscle, which crosses it near its middle. (Shown in the cut.) The upper portion is comparatively superficial, and occupies what on this account is called the triangle of election. The lower portion lies deeper, and occupies what for this reason is called the triangle of necessity. (The cut represents the incisions for both these ligations united.)

5. Ligation of the Common Carotid Artery in the Triangle of Election.—The patient should be recumbent, with the head extended and turned toward the opposite side. An incision should be made along the anterior edge of the sterno-cleido-mastoid muscle, from about an inch below the angle of the lower jaw to a little below the cricoid cartilage. The superficial fascia, the platysma-myoides muscle, and the deep fascia are then to be divided on a director. The sterno-mastoid muscle must then be held back, and the deep layer of the deep fascia picked up and carefully divided on a director. This will expose the sheath of the blood-vessels and perhaps the *descendens noni* nerve. Avoiding the latter, the sheath of the vessels is to be picked up with forceps and nicked with a scalpel. A director is then to be introduced, and the sheath slit up upon it. The artery is now to be separated from the internal jugular vein, which lies to the outer side, and from the pneumogastric nerve, which lies back of and between the artery and vein. The ligature should be passed from without inward.

6. Ligation of the Common Carotid Artery in the Triangle of Necessity.—The position of the patient should be the same as for the operation just described. The incision should be made along the inner edge of the sterno-mastoid muscle, and from the middle of the neck to the top of the sternum. The skin, superficial fascia, and platysma-myoides muscle are to be divided. Then the deep fascia is to be divided on a director, avoiding injury to the anterior jugular vein. When the sides of the wound are



drawn asunder, another layer of the deep fascia will be exposed, which must be lifted with forceps, nicked, and then carefully divided on a director. Under this lies the sheath of the blood-vessels, and upon it the *descendens novi* nerve. The subsequent steps of the operation are the same as those described for ligation in the triangle of election.

7. Ligation of the Subclavian Artery.—The point of election for ligation of this artery is the outer side of the *scalenus anticus* muscle. An incision is made parallel to and just above the clavicle, from the outer half of the clavicular insertion of the sterno-cleido-mastoid muscle to the anterior edge of the trapezius muscle, taking care not to go deep enough to cut the external jugular vein. The skin, superficial fascia, and the *platysma-myoides* muscle are to be divided. Then the external jugular vein is to be drawn aside. If this be impossible, it must be divided between two ligatures. The loose connective tissue must now be carefully broken up with a director and a search made for the artery. It must be remembered that it lies to the outer side of the *scalenus anticus* muscle and to the inner side of the cords of the brachial plexus. When found, it is to be very cautiously separated from its attachments to the pleura, and the ligature passed from below upward. The greatest care is necessary to avoid injuring the vein or the pleura, or including a nerve in the ligature.

8 and 9. Ligation of the Axillary Artery.—This artery is ligated in its first portion just after passing over the first rib, and in its third portion in the axilla.

8. Ligation of the First Portion of the Axillary Artery.—The arm is drawn away from the side, and an incision about three inches in length is made in the line which marks the division between the upper and lower parts of the *pectoralis major* muscle. The superficial and deep fascia must be divided on a director. The muscular interspace is then to be sought for, and the two parts of the muscle separated, working up toward the clavicle, and not directly backward. As this is done, the arm should be brought back to the side, so as to relax the part. The *costo-coracoid* membrane is next to be found and opened on a director. The artery lies beneath it. It is to be carefully isolated from the axillary vein, which lies on its inner, or lower, side, and the axillary nerves, which lie to its outer side. The ligature should be passed from within outward, or from below upward.

9. Ligation of the Axillary Artery in the Axilla.—This is the most accessible part of the artery. It is reached by drawing the arm away from the body, and making an incision about three inches long over the course of the vessel, at the junction of the anterior and middle thirds of the axilla. The deep fascia must be divided on a director, and the artery will be found beneath it, with the axillary vein on the ulnar side (toward the back of the arm), and the median nerve on the radial side (toward the front of the arm). The ligature should be passed from the ulnar toward the radial side, so as not to injure the vein.

10. Ligation of the Brachial Artery. in the middle of the arm.—An incision about two inches in length is to be made along the inner edge of the swell of the biceps muscle, at the middle of the arm. After dividing the skin and superficial fascia, and any fat that may be met with, the fibrous membrane which encloses the biceps muscle will be encountered. This is to be divided on a director, and the loose connective tissue along the edge of the muscle broken up, when the artery will be exposed. Care must be taken not to divide the median nerve, which crosses the brachial artery near its middle, passing from the radial toward the ulnar side, and not to include it in the ligature. The *venae comites* lie one on each side of the artery.

11. Ligation of the Radial Artery. in the upper third.—An incision is to be made in a line drawn from the middle of the bend of the elbow to the point at which the pulse can be felt at the wrist. The median vein is to be avoided in incising the superficial fascia. When the *lat. fasc.* has been exposed, a yellow line, marking the division between the *supinator longus*, on the radial side, and the *pronator radii teres*, on the ulnar side, is to be looked for. The fascia is to be opened along this line.

and the artery will be found below it. The radial nerve lies to the radial side of the artery, but so far away as to be in little danger of being injured.

12. Ligation of the Ulnar Artery.—In the lower third. An incision is to be made in the depression between the flexor sublimis digitorum and the flexor carpi ulnaris, just above the carpus. The skin and superficial fascia are to be divided, and then the deep fascia is to be picked up with the forceps and divided on a director. The artery will be found lying on the pronator quadratus, alongside the tendon of the flexor carpi ulnaris, with the ulnar nerve on its ulnar side. The ligature should be passed from the ulnar toward the radial side, avoiding the nerve.

13. Ligation of the External Iliac Artery.—An incision is to be made nearly parallel to Poupart's ligament, and about an inch above it, extending from about an inch above the anterior superior spinous process of the ilium to a point near the external abdominal ring. This incision should curve slightly upward at both ends. The skin and superficial fascia are to be divided, and ligatures to be applied to any vessels which bleed. The tendon of the external oblique muscle is to be divided, and then the internal oblique and the transversalis muscles. The separation of the transversalis fascia is a work of the utmost delicacy, to be effected with the forceps and the finger, the latter restraining and protecting the peritoneum. A finger passed down into the pelvis through the opening will discover the artery. The accompanying iliac vein lies to the inner side of the artery, and the anterior crural nerve at some distance to its outer side. The ligature should be passed from within outward.

14. Ligation of the Femoral Artery in Scarpa's Triangle.

—The point of election for ligation of the femoral artery is at the apex of Scarpa's triangle. To accomplish it, the leg should be flexed at the knee, and the thigh slightly flexed upon the abdomen, abducted and rotated outward, so as to lie on its outer side. An incision should then be made, over the course of the artery, in a line bisecting Scarpa's triangle from the middle of its base to its apex. This incision should be about three inches long, and its middle should be over the apex of the triangle. The skin, superficial fascia, and fat having been divided, the inner edge of the sartorius muscle must be found and drawn outward. The artery will now be found, with the vein to the inner side and a little behind it, and several nerves on the outer side. The artery should be carefully isolated after opening its sheath, and the ligature passed from within outward.

15. Ligation of the Femoral Artery in Hunter's Canal.—

This rare ligation is performed by placing the limb in the position for the former operation, and making an incision from the apex of Scarpa's triangle toward the back part of the internal condyle of the femur, avoiding the saphena vein. After passing through the skin, superficial fascia, and fat the deep fascia will be seen, binding down the sartorius muscle. This membrane is to be opened on a director and the muscle drawn backward. This will disclose the aponeurotic sheath which covers the femoral artery at this point. This sheath must be opened, and the artery will be exposed. In ligating it, care must be taken not to injure the long saphenous nerve, which accompanies it, or the femoral vein, which here lies behind it and toward its outer side. The ligature should be passed very carefully from without inward.

16. Ligation of the Anterior Tibial Artery, in the middle

third.—An incision is to be made, about three inches long, in a line passing from a point midway between the tubercle of the tibia and the head of the fibula, to a point midway between the two malleoli. When the deep fascia has been reached, a yellow line may be found, indicating the septum between the tibialis anticus muscle and the extensor longus digitorum. Along this line the aponeurosis is to be divided on a director, and the two muscles carefully separated until the extensor proprius pollicis is met with, when this is to be separated from the tibialis anticus until the artery is reached. It is accompanied by two vene comites, which it is not important to regard if they are difficult to separate from the artery. The anterior tibial nerve lies either in front or at the outer side of the artery. To avoid injuring it, it should be pressed to the outer side of the wound and the ligature should be passed from without inward.

17. Ligation of the Posterior Tibial Artery.—In the middle third.—An incision three inches long is to be made parallel to the inner border of the tibia and about half an inch back of it, and carried down to the superficial fascia. This is to be carefully opened, avoiding injury to the internal saphenous vein and nerve, which lie in the track of the wound. They must be found and drawn aside. Then the deep fascia is to be divided on a director, and the inner border of the soleus muscle is to be pressed away from the tibia, when the vessel will be exposed. It is accompanied by two venæ comites and the posterior tibial nerve, which lies to the fibular side of it. The veins and nerve being separated from the artery, the ligature should be passed from the fibular toward the tibial side.

18. Ligation of the Posterior Tibial Artery behind the Internal Malleolus.—A slightly curving incision an inch and a half long is to be made behind the internal malleolus, and about half-way between it and the tendo Achillis. The skin and superficial fascia having been divided, the deep fascia is to be lifted and divided on a director. The artery will then be exposed, accompanied by its venæ comites, and the posterior tibial nerve, which lies to the back of it and toward the fibular side. The ligature should be passed from behind forward.

19. Ligation of the Dorsalis Pedis Artery.—An incision an inch long is to be made on the back of the foot between the inner tendon of the extensor brevis digitorum and the outer side of the tendon of the extensor longus pollicis. (These tendons can be mapped out by manipulating the great toe.) After dividing the skin, the superficial fascia, and the deep fascia, the artery will be exposed. The continuation of the anterior tibial nerve lies to the fibular side of the artery. The ligature should be passed from the fibular toward the tibial side.

NOTE.—It is important in ligating arteries to make all the incisions carefully, to divide all deep structures on a director, to pass the ligature first between the artery and any important vein close to it, so as afterward to pass away from the dangerous neighborhood, to be sure not to tie a vein, or a nerve, or anything else for an artery. Opinions differ as to the best material for ligatures. Good carbolized catgut, if well applied, is the best material. But, as it is not always easy to get catgut which can be relied upon, nor can everybody apply it just as it should be applied, the material most generally applicable is unquestionably stout silk or linen thread. This should be well soaked in a solution of carbolic acid or of corrosive sublimate before being used. The knots should be made firm. In the case of catgut the ends should be cut off short and the whole wound closed, so as to secure immediate union. When silk or linen thread is used, it is usually best to cut one end off short and to leave the other hanging out of the wound.

After ligating a main artery, the parts below it should be kept absolutely quiet, and their warmth maintained by artificial means, so as to diminish the liability to gangrene.

THE
MEDICAL NEWS
VISITING LIST.
RECORDS OF PRACTICE.
1886.

NAME.

Su. M. Tu. W. Th. F. S.

Record.

January, 1886.

71

MEMORANDA

CHARGE

PREP
L.

NAME.

Su.	M.	Tu.	W.	Th.	F.	Sa.
3	1	5	6	7	8	

*William T. ...**W. T. ...**W. T. ...*

25

Record.

January, 1886.

15

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

Su.	M.	Tu.	W.	Th.	F.	Sa.
10	11	12	13	14	15	16

The ...
 ...
 ...
 ...
 ...

Record.

January, 1886.

၁၆

MEMORANDA

CHARGE.

Page in
Ledger.

NAME.

Su.	M.	Tu.	W.	Th.	F.	Sa.
17	18	19	20	21	22	23

1. ...
 2. ...
 3. ...
 4. ...
 5. ...
 6. ...
 7. ...
 8. ...
 9. ...
 10. ...
 11. ...
 12. ...
 13. ...
 14. ...
 15. ...
 16. ...
 17. ...
 18. ...
 19. ...
 20. ...
 21. ...
 22. ...
 23. ...
 24. ...
 25. ...
 26. ...
 27. ...
 28. ...
 29. ...
 30. ...
 31. ...

MEMORANDA.

CHARGE

Page in
Ledger

January, 1886.

Daily

January-February, 1886.

Daily

NAME.

Su M. Tu. W. Th F Sa

1 2 3 4 5 6

Jan 1st		X	X	X	X	X	
Jan 2nd			/				/
Jan 3rd							
Jan 4th							
Jan 5th							
Jan 6th							
Jan 7th							
Jan 8th							
Jan 9th							
Jan 10th							
Jan 11th							
Jan 12th							
Jan 13th							
Jan 14th							
Jan 15th							
Jan 16th							
Jan 17th							
Jan 18th							
Jan 19th							
Jan 20th							
Jan 21st							
Jan 22nd							
Jan 23rd							
Jan 24th							
Jan 25th							
Jan 26th							
Jan 27th							
Jan 28th							
Jan 29th							
Jan 30th							
Jan 31st							

MEMORANDA

CHARGE.

Page in
Ledger*4. Dec. 1886*

MEMORANDA.

CHARGE

Page in
Ledger

Patient of Dr. O'Hara. In catarrh of
urine suspected

February, 1886.

Daily

[illegible]

Record.

February, 1886.

65

MEMORANDA.

CHARGE.

Page in
Ledger

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

Su.	M.	Tu.	W.	Th.	F.	Sa.
28	1	2	3	4	5	6

The following is a list of
 the names of the
 persons who have been
 admitted to the
 membership of the
 Association since the
 last meeting.

The following is a list of
 the names of the persons who
 have been admitted to the
 membership of the Association
 since the last meeting.

Record.

February - March, 1886.

49

MEMORANDA.

CHARGE.

Page in
Ledger.

March, 1886.

Daily

NAME

Su. M. Tu. W. Th. F. S.
7 8 9 10 11 12 1

17 18

1

Chertington

10

17 18 19 20 21 22 23 24 25 26 27 28 29 30 31

1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31.

Record.

March, 1886.

71

MEMORANDA

CHARGE

Page 10

L. L.

NAME

| Su | M. | Tu. | W | Th | F | S |
|----|----|-----|----|----|----|----|
| 14 | 15 | 16 | 17 | 18 | 19 | 20 |

| | | | | | | |
|--|--|--|---|---|--|--|
| | | | X | X | | |
|--|--|--|---|---|--|--|

March 23rd. Sunday. Clear. Windy. No rain. With cold rain.

March 24th. Monday. Clear. Windy. No rain. With cold rain.

March 25th. Tuesday. Clear. Windy. No rain. With cold rain.

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 21 | 22 | 23 | 24 | 25 | 26 | 27 |

X

Y

12

Record.

March, 1886.

75

MEMORANDA

CHARGE

Page in
Ledger

100 2 00

5

NAME.

| Su | M | Tu | W | Th | F | Sa |
|----|----|----|----|----|---|----|
| 28 | 29 | 30 | 31 | 1 | 2 | 3 |

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 |
|---|---|---|---|---|---|---|---|---|----|

1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31.

1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31.

1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31.

1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31.

1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31.

MEMORANDA

CHARGE

Page in
Ledger.

April, 1886.

Daily

NAME

Su. M. Tu. W. Th. F. Sa.
1 2 3 4 5 6 7 8 9 10

[Faint handwritten notes, possibly names or dates, mostly illegible due to fading.]

[Faint handwritten notes in the right margin, possibly dates or initials.]

Record.

April, 1886.

79

MEMORANDA

CHARGE

Page in
Ledger

April, 1886.

Daily

NAME

| Su. | M. | Tu. | W | Th. | F | Sa |
|-----|----|-----|----|-----|----|----|
| 11 | 12 | 13 | 14 | 15 | 16 | 17 |

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

A. P. Smith

Record.

April, 1886.

81

MEMORANDA.

CHARGE.

Page in
Ledger.

April, 1887.

Daily

NAME.

| Su | M | Tu | W | Th | F | Sa |
|----|----|----|----|----|----|----|
| 18 | 19 | 20 | 21 | 22 | 23 | 24 |

R. P. Ketchum
 and Tracy
 went to
 Hunt

C C C C

Record.

April, 1886.

83

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 25 | 26 | 27 | 28 | 29 | 30 | |

1. *My first*
letter to you
was written
on May 1st
1886.

Record.

April—May, 1886.

85

MEMORANDA

CHARGE

Page in
Ledger

May, 1886.

Daily

| NAME | Su. | M | Tu. | W. | Th. | F. | Sa. |
|------|-----|---|-----|----|-----|----|-----|
| | 2 | 3 | 4 | 5 | 6 | 7 | 8 |

MEMORANDA.

CHARGE

Page in
Ledger

Sept. 21st died
P.M. 1886.

transf

NAME

| Su | M | Tu | W | Th | F | Sa |
|----|----|----|----|----|----|----|
| 9 | 10 | 11 | 12 | 13 | 14 | 15 |

[Faint handwritten notes, possibly bleed-through from the reverse side of the page. The text is largely illegible due to fading and the quality of the scan.]

MEMORANDA

CHARGE

Page in
Ledger.

May, 1856.

Daily

NAME

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 16 | 17 | 18 | 19 | 20 | 21 | 22 |

Handwritten notes in the left margin, including the word "C" and some illegible cursive text.

MEMORANDA

CHARGE

Page in
Ledger

NAME.

Su M Tu W Th F Sa
23 24 25 26 27 28 29

[Faint handwritten text, likely bleed-through from the reverse side of the page. The text is mostly illegible due to fading.]

Record.

May, 1886.

93

MEMORANDA.

CHARGE.

Page in
Ledger

NAME

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 30 | 31 | 1 | 2 | 3 | 4 | 5 |

MEMORANDA.

CHARGE.

Page in
Ledger

Record.

June, 1886.

97

MEMORANDA.

CHARGE

Page in
Ledger

June, 1886.

Daily

Record

NAME

| Su | M. | Tu. | W. | Th | F. | Sa |
|----|----|-----|----|----|----|----|
| 13 | 14 | 15 | 16 | 17 | 18 | 19 |

MEMORANDA.

CHARGE.

Page in
Ledger

NAME.

| Su. | M. | Tu. | W | Th | F. | Sa |
|-----|----|-----|----|----|----|----|
| 20 | 21 | 22 | 23 | 24 | 25 | 26 |

Handwritten notes:
June 20, 1888
June 21, 1888

MEMORANDA.

CHARGE

Page in
Ledger.

NAME

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 27 | 28 | 29 | 30 | 1 | 2 | 3 |

See ledger notebook

1 2

MEMORANDA.

CHARGE.

Page in
Ledger.

| NAME. | Su. | M. | Tu. | W. | Th. | F. | Sa. |
|----------------------------------|-----|----|-----|----|-----|----|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| Mr. W. H. Hamilton | | | | | | | Dr. |
| Lt. Col. Landon 1139 Girard St | | | | | | | Dr. |
| Mr. Devine | | | | | | | C 1 |
| Mr. Martin (Dr. with Dr. Morris) | | | | | | | C OH |
| Mr. J. J. J. | | | | | | | |

+ Lucie - at 11 Littlefinger life
 des. The 1st segment completely in head
 segment constantly very raised. Last segment
 in the neighborhood. Abundant
 ability. Susceptible of hysterical movement in the
 lower inf 65-66. T. 101°. Drag. moving

Record.

July, 1886.

105

MEMORANDA.

CHARGE


Page in
Ledger.

NAME

| Su. | M | Tu. | W. | Th | F. | Sa |
|-----|----|-----|----|----|----|----|
| 11 | 12 | 13 | 14 | 15 | 16 | 17 |

Mrs. Allen no actual work:
 Pittsburgh, returned Cap
 no work 8

Last day. S. side of river.
 shells & bones & wood only. S. side of river.
 no. 1. later on left side with sand. 3rd
 shell below T. 101. P. 120. P. 120. P. 120.
 no remains, no tubercles. Tub. 120. P. 120.



MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Su. | M. | Tu. | W. | Th | F. | Sa |
|-----|----|-----|----|----|----|----|
| 18 | 19 | 20 | 21 | 22 | 23 | 24 |

3

73

MEMORANDA.

CHARGE.

Page in
Ledger

NAME

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 25 | 26 | 27 | 28 | 29 | 30 | 31 |

Record.

July, 1886.

111

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 |

Record.

August, 1886.

113

MEMORANDA.

CHARGE.

Page in
Ledger

NAME

| Sp. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 8 | 9 | 10 | 11 | 12 | 13 | 14 |

Record.

August, 1886.

115

MEMORANDA.

CHARGE.

Page in
Ledger

NAME

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 15 | 16 | 17 | 18 | 19 | 20 | 21 |

Record.

August, 1886.

117

MEMORANDA.

CHARGE

Page in
Ledger

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 22 | 23 | 24 | 25 | 26 | 27 | 28 |

Record.

August, 1886.

119

MEMORANDA.

CHARGE.

Page in
Ledger

NAME.

| Sa. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 29 | 30 | 31 | 1 | 2 | 3 | 4 |

W. C. 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600, 610, 620, 630, 640, 650, 660, 670, 680, 690, 700, 710, 720, 730, 740, 750, 760, 770, 780, 790, 800, 810, 820, 830, 840, 850, 860, 870, 880, 890, 900, 910, 920, 930, 940, 950, 960, 970, 980, 990, 1000.

a.

c.

W. C. 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, 300, 310, 320, 330, 340, 350, 360, 370, 380, 390, 400, 410, 420, 430, 440, 450, 460, 470, 480, 490, 500, 510, 520, 530, 540, 550, 560, 570, 580, 590, 600, 610, 620, 630, 640, 650, 660, 670, 680, 690, 700, 710, 720, 730, 740, 750, 760, 770, 780, 790, 800, 810, 820, 830, 840, 850, 860, 870, 880, 890, 900, 910, 920, 930, 940, 950, 960, 970, 980, 990, 1000.

Record.

August—September, 1886.

121

MEMORANDA.

CHARGE.

Page in
Ledger.

| NAME | Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----------------------|-----|----|-----|----|-----|----|-----|
| | 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| Her. 4/12 | | | | | Q | | 1/1 |
| Her. 4/12 | | | | | Q | | 1/1 |
| Her. 4/12 | | | | | Q | | 1/1 |
| C. Smith. 6/12 | | | | | | | 1/1 |
| * P. 4/12 009 P. 4/12 | | | | | | | 1/1 |
| 1/12 4/12 | | | | | | | 1/1 |

Chronic 4/12 1/1

Record.

September, 1886.

123

MEMORANDA

CHARGE.

Page in
Ledger.

MEMORANDA.

CHARGE

Page in
Ledger.

NAME

| Su. | M. | Tu. | W | Th. | F | Sa |
|-----|----|-----|----|-----|----|----|
| 19 | 20 | 21 | 22 | 23 | 24 | 25 |

Mr Phelps

c

Mr W. L. G. with Mr. Allen

c

Record.

September, 1886.

127

MEMORANDA.

| CHARGE

Page in
Ledger.

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 26 | 27 | 28 | 29 | 30 | 1 | 2 |

1st of Sept.
 1886. 10th Sept.

C

P

, 0

Record.

September-October, 1886.

129

MEMORANDA.

1 CHARGE.

Page in
Ledger.

NAME.

| Su | M | Tu | W | Th | F | Sa |
|----|---|----|---|----|---|----|
| 3 | 4 | 5 | 6 | 7 | 8 | 9 |

W. N. S. S. S. S.

C

Record.

October, 1886.

131

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Su | M. | Tu. | W. | Th | F. | Sa |
|----|----|-----|----|----|----|----|
| 10 | 11 | 12 | 13 | 14 | 15 | 16 |

L. Brown 1439 4 76
for 10

10

10

Record.

October, 1886.

133

MEMORANDA.

CHARGE

Page in
Ledger

MEMORANDA

CHARGE

Page in
Ledger

from the edge not distinct
inter 20, then the soft bottom
then skin.

3.
... ..
the a

... .. by

Others should be considered.

1. Lympho sarcoma retroperitoneal
... ..

2. Pancreatic

3.
... ..
... ..

... .. growth - eccentric
... .. points
... ..
... ..

W.O. 18/10/86

... ..
... ..

NAME

| Su. | M | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 21 | 25 | 26 | 27 | 28 | 29 | 31 |

New Hampshire 943 N 711
 St. Kitts
 St. Kitts

c c c

at 32. 1305 with 1000 yds. from.

It was 2 yrs. 2 yrs. ago with a mad dog. case
 reported by Dr. Hamilton in 1885 for St. Kitts.
 Bladder with 2 yrs. 2 yrs. ago. called
 himself. 4 y. ago. had a dec. attack of St. Kitts
 in 1885 at base. all 10 weeks. in Cal. 20 months
 24 y. very little cough. From 1885 to 1886
 2 yrs. for 3 mos. He had several attacks
 twice. had. for 1885 to 1886.

He seemed to be 2 yrs. 2 yrs. ago. present
 Maltese. weather.

dump. It seems pretty clear. a few notes. at
 mar. of sternum. left. 5 notes whiffing with
 in 1885. in 1885. 23 of 23 notes. 23 notes
 in 2nd. very abundant. interruption in 1885
 marked. 5-6 notes. 1885. 1885.

MEMORANDA

CHARGE.

Page in
Ledger.

at river a well a bare relative flatness,
deeper part breaching a greater
color in deep appearance.

none.

2. thin

7. none part 3. large

| NAME | Su. | M. | Tu. | W. | Th. | F. | Sa. |
|------|-----|----|-----|----|-----|----|-----|
| | 31 | 1 | 2 | 3 | 4 | 5 | 6 |

Dr Boardman

c c

News Newspaper

x

Her Morris Mr Hayes 40¢ each

c.

Dr Her Blumens. Ch. 1 8 16 25

c

Bl. color 38 R p c. m. 3 100000

was at 12th. 15. color R. about 3 0000 3 weeks ago

Record.

October—November, 1886.

149

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME

Su. M. Tu. W. Th. F. Sa.
7 8 9 10 11 12 13

Mr John M. Cord 2004 Spruce

C

Record.

November, 1886.

141

MEMORANDA

CHARGE.

Page in
Ledger

| NAME. | Su. | M. | Tu. | W. | Th. | F. | Sa. |
|----------------------------------|-----|----|-----|----|-----|----|-----|
| | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
| C H Muller Chesh. Hall | e | | | | | | |
| c or ghy Swedenboro. n g | | | E | | | | J |
| Mr Mellan 237 N 16 | | | C | | | | |
| W - n clord 2004 Spruce | | | | | | | Cor |
| Mr H. and George 1730 Sp Garden | | | | | | C | |
| Leut. H. Mearns 1730 Sp Garden | | | | | | C | |
| Stirling Costigan 3115 Sp Garden | | | | | | C | |
| Dr. Mearns: gill. | | | | | | | I |

Record.

November, 1886.

143

MEMORANDA.

CHARGE.

Page in
Ledger

| NAME. | Su. | M. | Tu. | W. | Th. | F. | Sa. |
|---|-----|----|-----|----|-----|----|-----|
| | 21 | 22 | 23 | 24 | 25 | 26 | 27 |
| <i>Mon. 9.</i>
Mr. Wm Syckelmore 1425 ^{ans} | | | | | | | C |
| Capt J W Stratford | | | | | | | C |
| Mr. W. L. Lepper 967 Franklin | | | | | | | C |
| Mr. B. L. Lepper 1910 Green | | | | | | | C |

Record.

November, 1886.

145

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Sa. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 28 | 29 | 30 | 1 | 2 | 3 | 4 |

Mrs

142 5 Mont. ad C

Mrs M. W. de la Roche 967 Franklin C.

Mrs M. C.

C

Mrs M. C.

C C

Record.

November—December, 1886.

147

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME

| Su. | M | Tu. | W. | Th. | F | Sa. |
|-----|---|-----|----|-----|----|-----|
| 5 | 6 | 7 | 8 | 9 | 10 | 11 |

Mr Sime 2210 St James Place C.

Mr McCord C

Record.

December, 1886.

149

MEMORANDA.

CHARGE.

Page in
Ledger

| NAME | Su | M. | Th. | W. | Th. | F. | S. |
|--------------------------------|----|----|-----|----|-----|---------|----|
| | 12 | 13 | 14 | 15 | 16 | 17 | 18 |
| C. S. B. Bentley 20202 40 | | | | C | C | 10 p.d. | |
| P. Lang. name | | | | | C | 10 p.d. | |
| Mrs. Leary 138 2 10 | | | | C | | 200 | |
| P. F. name | | | | | C | | |
| Dr. Kelly 3021 Transit. and 25 | | | | | C | | |

after 2' and end upple. map with very possible
 no more after. Rungy and at once.
 Passes good ground of some. No palpitation. No
 extension. going up stream. A man is down
 the 20 by T. name.
 of T.

. the water was
 health. All a year
 Sat. and my stomach
 Put 40 C.

Record.

December, 1886.

151

MEMORANDA.

CHARGE.

Page 11
1. 1886

Record.

December, 1886.

153

MEMORANDA.

CHARGE.

Page in
Ledger

NAME

| Su | M | Tu | W | Th | F | Sa |
|----|----|----|----|----|----|----|
| 26 | 27 | 28 | 29 | 30 | 31 | |

Record.

December, 1886.

155

MEMORANDA.

CHARGE.

Page in
1542-1

100 PATIENT'S NAME.

Clinical

DATE

PULSE

TEMP.

RESP.

URINE.

STOOLS

TREATMENT CONDITION REMARKS

Chin

100

110

120

130

140

150

160

170

180

DATE

PULSE

TEMP

RESP

URINE

STOOLS

Record.

100

TREATMENT CONDITION REMARKS

130

PATIENT'S NAME.

Clinical

DATE

PULSE.

TEMP.

RESP.

URINE.

STOOLS

TREATMENT. CONDITION. REMARKS.

DATE

PULSE

TEMP

RESP

URINE

SIGNS

Record.

No.

TREATMENT.—CONDITION.—REMARKS.

| DATE. | PULSE. | TEMP. | RESP. | URINE. | STOOLS. |
|-------|--------|-------|-------|--------|---------|
|-------|--------|-------|-------|--------|---------|

TREATMENT — CONDITION. — REMARKS

| DATE. | PULSE | TEMP | RESP. | URINE. | STOOLS |
|-------|-------|------|-------|--------|--------|
|-------|-------|------|-------|--------|--------|

TREATMENT — CONDITION. — REMARKS

| DATE. | PULSE. | TEMP. | RESP. | URINE. | STOOLS |
|-------|--------|-------|-------|--------|--------|
|-------|--------|-------|-------|--------|--------|

TREATMENT.—CONDITION.—REMARKS.

DATE

PULSE

TEMP.

RESP.

URINE

STOOLS

| TREATMENT | LOCATION | REMARKS |
|-----------|----------|---------|
|-----------|----------|---------|

DATE.

PULSE.

TEMP.

RESP.

URINE.

STOOLS

| TREATMENT. | CONDITION. | REMARKS. |
|------------|------------|----------|
| 1 | 2 | 3 |
| 4 | 5 | 6 |
| 7 | 8 | 9 |
| 10 | 11 | 12 |
| 13 | 14 | 15 |
| 16 | 17 | 18 |
| 19 | 20 | 21 |
| 22 | 23 | 24 |
| 25 | 26 | 27 |
| 28 | 29 | 30 |
| 31 | 32 | 33 |
| 34 | 35 | 36 |
| 37 | 38 | 39 |
| 40 | 41 | 42 |
| 43 | 44 | 45 |
| 46 | 47 | 48 |
| 49 | 50 | 51 |
| 52 | 53 | 54 |
| 55 | 56 | 57 |
| 58 | 59 | 60 |
| 61 | 62 | 63 |
| 64 | 65 | 66 |
| 67 | 68 | 69 |
| 70 | 71 | 72 |
| 73 | 74 | 75 |
| 76 | 77 | 78 |
| 79 | 80 | 81 |
| 82 | 83 | 84 |
| 85 | 86 | 87 |
| 88 | 89 | 90 |
| 91 | 92 | 93 |
| 94 | 95 | 96 |
| 97 | 98 | 99 |
| 100 | 101 | 102 |
| 103 | 104 | 105 |
| 106 | 107 | 108 |
| 109 | 110 | 111 |
| 112 | 113 | 114 |
| 115 | 116 | 117 |
| 118 | 119 | 120 |
| 121 | 122 | 123 |
| 124 | 125 | 126 |
| 127 | 128 | 129 |
| 130 | 131 | 132 |
| 133 | 134 | 135 |
| 136 | 137 | 138 |
| 139 | 140 | 141 |
| 142 | 143 | 144 |
| 145 | 146 | 147 |
| 148 | 149 | 150 |
| 151 | 152 | 153 |
| 154 | 155 | 156 |
| 157 | 158 | 159 |
| 160 | 161 | 162 |
| 163 | 164 | 165 |
| 166 | 167 | 168 |
| 169 | 170 | 171 |
| 172 | 173 | 174 |
| 175 | 176 | 177 |
| 178 | 179 | 180 |
| 181 | 182 | 183 |
| 184 | 185 | 186 |
| 187 | 188 | 189 |
| 190 | 191 | 192 |
| 193 | 194 | 195 |
| 196 | 197 | 198 |
| 199 | 200 | 201 |
| 202 | 203 | 204 |
| 205 | 206 | 207 |
| 208 | 209 | 210 |
| 211 | 212 | 213 |
| 214 | 215 | 216 |
| 217 | 218 | 219 |
| 220 | 221 | 222 |
| 223 | 224 | 225 |
| 226 | 227 | 228 |
| 229 | 230 | 231 |
| 232 | 233 | 234 |
| 235 | 236 | 237 |
| 238 | 239 | 240 |
| 241 | 242 | 243 |
| 244 | 245 | 246 |
| 247 | 248 | 249 |
| 250 | 251 | 252 |
| 253 | 254 | 255 |
| 256 | 257 | 258 |
| 259 | 260 | 261 |
| 262 | 263 | 264 |
| 265 | 266 | 267 |
| 268 | 269 | 270 |
| 271 | 272 | 273 |
| 274 | 275 | 276 |
| 277 | 278 | 279 |
| 280 | 281 | 282 |
| 283 | 284 | 285 |
| 286 | 287 | 288 |
| 289 | 290 | 291 |
| 292 | 293 | 294 |
| 295 | 296 | 297 |
| 298 | 299 | 300 |
| 301 | 302 | 303 |
| 304 | 305 | 306 |
| 307 | 308 | 309 |
| 310 | 311 | 312 |
| 313 | 314 | 315 |
| 316 | 317 | 318 |
| 319 | 320 | 321 |
| 322 | 323 | 324 |
| 325 | 326 | 327 |
| 328 | 329 | 330 |
| 331 | 332 | 333 |
| 334 | 335 | 336 |
| 337 | 338 | 339 |
| 340 | 341 | 342 |
| 343 | 344 | 345 |
| 346 | 347 | 348 |
| 349 | 350 | 351 |
| 352 | 353 | 354 |
| 355 | 356 | 357 |
| 358 | 359 | 360 |
| 361 | 362 | 363 |
| 364 | 365 | 366 |
| 367</ | | |

| DATE. | PULSE. | TEMP. | RESP. | URINE. | STOOLS |
|-------|--------|-------|-------|--------|--------|
|-------|--------|-------|-------|--------|--------|

TREATMENT.—CONDITION.—REMARKS.

DATE.

PULSE.

TEMP.

RESP.

URINE.

STOOLS

TREATMENT.—CONDITION.—REMARKS.

| DATE. | PULSE | TEMP. | RESP. | URINE. | STOOLS. |
|-------|-------|-------|-------|--------|---------|
|-------|-------|-------|-------|--------|---------|

TREATMENT.—CONDITION.—REMARKS.

180

●

DATE.

PULSE.

TEMP.

RESP.

URINE.

STOOLS

TREATMENT. — CONDITION. — REMARKS

| DATE. | PULSE | TEMP. | RESP. | URINE. | STOOLS. |
|-------|-------|-------|-------|--------|---------|
|-------|-------|-------|-------|--------|---------|

TREATMENT - CONDITION - REMARKS

DATE.

TIME.

PATIENT'S NAME AND ADDRESS.

ATTENDING PHYSICIAN

REMARKS.

186

Consultation

DATE

TIME

PATIENT'S NAME AND ADDRESS

ATTENDING PHYSICIAN

REMARKS.

ISS

Consultation

DATE

TIME

PATIENT'S NAME AND ADDRESS

ATTENDING PHYSICIAN.

REMARKS.

PATIENT'S NAME

End of Last ' Date Exp'd ' Date of
Menstruat'n, Confinement Delivery

PATIENT'S NAME

| End of Last
Menstruat'n, Confinement | Date Exp'd /
Confinement | Date of
Delivery |
|---|-----------------------------|---------------------|
|---|-----------------------------|---------------------|

| PATIENT'S NAME. | End of Last Menstruat'n. | Date Exp'td Confinement | Date of Delivery |
|-----------------|--------------------------|-------------------------|------------------|
|-----------------|--------------------------|-------------------------|------------------|

PATIENT'S NAME.

End of Last Date Exp'd
Menstruat'n, ConfinementDate of
Delivery.

| PATIENT'S NAME. | End of Last Date Exp'd
'Menstruat'n. Confinement | Date of
Delivery. |
|-----------------|---|----------------------|
|-----------------|---|----------------------|

Obstetric Engagements.

195

PATIENT'S NAME

End of Last | Date Exp'd
Menstruat'n. Confinement

Date of
Delivery

| DATE. | NAME OF CHILD | SEX. | NO. OF
CHILD | COLOR. |
|-------|---------------|------|-----------------|--------|
|-------|---------------|------|-----------------|--------|

PARENTS' NAMES, OCCUPATION AND ADDRESS. REMARKS.

Date _____

NAME OF CHILD.

SEX.

No. of

('HILD.

COLOR.

Practice.

134

PARENTS' NAMES, OCCUPATION AND ADDRESS. REMARKS.

| DATE | NAME OF CHILD. | SEX. | No. of
CHILD. | COLOR |
|------|----------------|------|------------------|-------|
|------|----------------|------|------------------|-------|

PARENTS' NAMES, OCCUPATION AND ADDRESS.—REMARKS

| DATE. | NAME OF CHILD | SEX. | No. of
CHILD | COLOR |
|-------|---------------|------|-----------------|-------|
|-------|---------------|------|-----------------|-------|

PARENTS' NAMES. OCCUPATION AND ADDRESS.—REMARKS

DATE.

NAME OF CHILD

SEX.

No. of
CHILD

COLOR

[illegible]

DATE

NAME OF CHILD.

SEX.

NO. OF
CHILD.

COLOR.

| PARENTS' NAMES, OCCUPATION AND ADDRESS | REMARKS |
|--|---------|
|--|---------|

Vaccinations.

209

NAME.

DATE.

RESULT.


Vaccinations.

211

NAME.

DATE

RESULT



NAME AND ADDRESS OF DECEASED

AGE

Y

M.

D.

COLOR

Mr'd, Single,
or Widowed.

CAUSE OF DEATH

DATE

NAME AND ADDRESS OF DECEASED

AGE

Y.

M.

D.

COLOR

Mr. Buckland 1910 St.

Mr'd, Single,
or Widowed.

CAUSE OF DEATH

DATE

ADDRESSES OF PATIENTS

Mrs Kamenitz

800 Cornelian st

ADDRESSES OF PATIENTS.

1. *[faint handwriting]*

2. *[faint handwriting]*

3. *[faint handwriting]*

4. *[faint handwriting]*

5. *[faint handwriting]*

6. *[faint handwriting]*

7. *[faint handwriting]*

8. *[faint handwriting]*

9. *[faint handwriting]*

10. *[faint handwriting]*

11. *[faint handwriting]*

12. *[faint handwriting]*

13. *[faint handwriting]*

14. *[faint handwriting]*

15. *[faint handwriting]*

16. *[faint handwriting]*

ADDRESSES OF PATIENTS

ADDRESSES OF PATIENTS.

NURSES' ADDRESSES

NURSES' ADDRESSES

Cash Account.

| BILL SENT | NAME. | AMOUNT CHARGED. | AMOUNT PAID. | WHEN PAID. |
|-----------|---------------|-----------------|--------------|------------|
| 1 | W. A. Luchner | 1.00 | | |
| 2 | W. A. Luchner | 1.00 | | |
| 3 | W. A. Luchner | 1.00 | | |
| 4 | W. A. Luchner | 1.00 | | |
| 5 | W. A. Luchner | 1.00 | | |
| 6 | W. A. Luchner | 1.00 | | |
| 7 | W. A. Luchner | 1.00 | | |
| 8 | W. A. Luchner | 1.00 | | |
| 9 | W. A. Luchner | 1.00 | | |
| 10 | W. A. Luchner | 1.00 | | |
| 11 | W. A. Luchner | 1.00 | | |
| 12 | W. A. Luchner | 1.00 | | |
| 13 | W. A. Luchner | 1.00 | | |
| 14 | W. A. Luchner | 1.00 | | |
| 15 | W. A. Luchner | 1.00 | | |
| 16 | W. A. Luchner | 1.00 | | |
| 17 | W. A. Luchner | 1.00 | | |
| 18 | W. A. Luchner | 1.00 | | |
| 19 | W. A. Luchner | 1.00 | | |
| 20 | W. A. Luchner | 1.00 | | |
| 21 | W. A. Luchner | 1.00 | | |
| 22 | W. A. Luchner | 1.00 | | |
| 23 | W. A. Luchner | 1.00 | | |
| 24 | W. A. Luchner | 1.00 | | |
| 25 | W. A. Luchner | 1.00 | | |
| 26 | W. A. Luchner | 1.00 | | |
| 27 | W. A. Luchner | 1.00 | | |
| 28 | W. A. Luchner | 1.00 | | |
| 29 | W. A. Luchner | 1.00 | | |
| 30 | W. A. Luchner | 1.00 | | |
| 31 | W. A. Luchner | 1.00 | | |
| 32 | W. A. Luchner | 1.00 | | |
| 33 | W. A. Luchner | 1.00 | | |
| 34 | W. A. Luchner | 1.00 | | |
| 35 | W. A. Luchner | 1.00 | | |
| 36 | W. A. Luchner | 1.00 | | |
| 37 | W. A. Luchner | 1.00 | | |
| 38 | W. A. Luchner | 1.00 | | |
| 39 | W. A. Luchner | 1.00 | | |
| 40 | W. A. Luchner | 1.00 | | |
| 41 | W. A. Luchner | 1.00 | | |
| 42 | W. A. Luchner | 1.00 | | |
| 43 | W. A. Luchner | 1.00 | | |
| 44 | W. A. Luchner | 1.00 | | |
| 45 | W. A. Luchner | 1.00 | | |
| 46 | W. A. Luchner | 1.00 | | |
| 47 | W. A. Luchner | 1.00 | | |
| 48 | W. A. Luchner | 1.00 | | |
| 49 | W. A. Luchner | 1.00 | | |
| 50 | W. A. Luchner | 1.00 | | |
| 51 | W. A. Luchner | 1.00 | | |
| 52 | W. A. Luchner | 1.00 | | |
| 53 | W. A. Luchner | 1.00 | | |
| 54 | W. A. Luchner | 1.00 | | |
| 55 | W. A. Luchner | 1.00 | | |
| 56 | W. A. Luchner | 1.00 | | |
| 57 | W. A. Luchner | 1.00 | | |
| 58 | W. A. Luchner | 1.00 | | |
| 59 | W. A. Luchner | 1.00 | | |
| 60 | W. A. Luchner | 1.00 | | |
| 61 | W. A. Luchner | 1.00 | | |
| 62 | W. A. Luchner | 1.00 | | |
| 63 | W. A. Luchner | 1.00 | | |
| 64 | W. A. Luchner | 1.00 | | |
| 65 | W. A. Luchner | 1.00 | | |
| 66 | W. A. Luchner | 1.00 | | |
| 67 | W. A. Luchner | 1.00 | | |
| 68 | W. A. Luchner | 1.00 | | |
| 69 | W. A. Luchner | 1.00 | | |
| 70 | W. A. Luchner | 1.00 | | |
| 71 | W. A. Luchner | 1.00 | | |
| 72 | W. A. Luchner | 1.00 | | |
| 73 | W. A. Luchner | 1.00 | | |
| 74 | W. A. Luchner | 1.00 | | |
| 75 | W. A. Luchner | 1.00 | | |
| 76 | W. A. Luchner | 1.00 | | |
| 77 | W. A. Luchner | 1.00 | | |
| 78 | W. A. Luchner | 1.00 | | |
| 79 | W. A. Luchner | 1.00 | | |
| 80 | W. A. Luchner | 1.00 | | |
| 81 | W. A. Luchner | 1.00 | | |
| 82 | W. A. Luchner | 1.00 | | |
| 83 | W. A. Luchner | 1.00 | | |
| 84 | W. A. Luchner | 1.00 | | |
| 85 | W. A. Luchner | 1.00 | | |
| 86 | W. A. Luchner | 1.00 | | |
| 87 | W. A. Luchner | 1.00 | | |
| 88 | W. A. Luchner | 1.00 | | |
| 89 | W. A. Luchner | 1.00 | | |
| 90 | W. A. Luchner | 1.00 | | |
| 91 | W. A. Luchner | 1.00 | | |
| 92 | W. A. Luchner | 1.00 | | |
| 93 | W. A. Luchner | 1.00 | | |
| 94 | W. A. Luchner | 1.00 | | |
| 95 | W. A. Luchner | 1.00 | | |
| 96 | W. A. Luchner | 1.00 | | |
| 97 | W. A. Luchner | 1.00 | | |
| 98 | W. A. Luchner | 1.00 | | |
| 99 | W. A. Luchner | 1.00 | | |
| 100 | W. A. Luchner | 1.00 | | |

• 3-4 •

[illegible]

Cash Account.

| BILL
SENT | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|--------------|------------------------|--------------------|-----------------|---------------|
| Upaid. | | 2 | | |
| | | <u>300</u> | | Pd July 15 46 |
| | | | | |
| | | | | |
| | | | | |
| | | 75 00 | | Paid July 84 |
| | | <u>15</u> | | Pd July 11 46 |
| | | | | |
| | | | | |
| | | | | Pd June |
| | | | | Pd June |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | 20 00 | | Pd July |
| | | | | |
| | | <u>5</u> | | |
| | T. H. Handlin | <u>50 00</u> | | Pd July |
| | | 30 00 | | |
| | S. L. Leasing - 9 Bond | <u>10 00</u> | | Pd July |
| | Leasing 1000 Bond | 10 00 | | |
| | T. H. Leasing - 1000 | 100 00 | | |
| | Chas. A. Leasing | 5 00 | | |
| | W. Leasing 1000 Bond | <u>15 00</u> | | Pd July |
| | Handlin | 10 00 | | |
| | W. Leasing 1000 Bond | 10 00 | | |

Cash Account.

| BILL
SENT. | NAME | AMOUNT
CHARGED | AMOUNT
PAID | WHEN
PAID |
|---------------|------|-------------------|----------------|--------------|
|---------------|------|-------------------|----------------|--------------|

Handwritten entry: [Faint text]

Handwritten entry: [Faint text]

Handwritten entry: [Faint text]

Handwritten entry: [Faint text]

Handwritten entry: [Faint text]

Handwritten entry: [Faint text]

Handwritten entry: [Faint text]

38

2

Handwritten entry: [Faint text]

Handwritten entry: [Faint text]

Handwritten entry: [Faint text]

Handwritten entry: [Faint text]

Cash Account.

| BILL
SENT | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID |
|--------------|-------|--------------------|-----------------|--------------|
|--------------|-------|--------------------|-----------------|--------------|

| | | | | |
|------|-------------|--------|--|--|
| Self | Smith, 7000 | 140 00 | | |
| 10 | | | | |
| 20 | White | 10 00 | | |

Cash Account.

207

| DATE | NAME. | AMOUNT | AMOUNT | WHEN |
|-------|--------------------------------|---------|-------------------|------|
| SENT. | | CHARGED | PAID. | PAID |
| | From Oct. 1st 1886 | 78 86 | | |
| 12 00 | News Magazine. 287. | 25 00 | | |
| 10 00 | Wm. Stricklin | 20 00 | 10/5 Pd Dec 9 | |
| 10/50 | Mitchison Sale | 30 00 | 10/5 Pd 23/11/86 | |
| 1/86 | John Brown | 10 00 | | |
| 1/80 | F. T. R. Robt. | 2 00 00 | | |
| 1/80 | T. H. Hancock | 10 00 | Jan 5th | |
| 1/86 | Philip 2009 Park Ave. | 60 00 | Jan 6th | |
| 1/86 | John (at Dr. Keen's) | 15 00 | Jan 8th | |
| 1/86 | Titus Case J. C. Smith | 150 00 | Pd 11/11/86 | |
| 1/86 | Mrs. Kempfen | 10 00 | Pd | |
| 1/86 | Mrs. Kempfen | 18 00 | Jan 6 | |
| 1/86 | Mrs. M. Keyser 408 Arch | 20 00 | Pd. | |
| 1/86 | Wm. Blumhardt 816 N 5th | 10 00 | Jan 12/13 Pd. 176 | |
| 1/86 | J. M. McLeod 2004 Spruce | 40 00 | Jan 6 | |
| 1/86 | J. H. Miller Chestnut Hill | 25 00 | Jan 6 | |
| 1/86 | Mellon 237 N 16th St | 10 00 | Jan 6 | |
| 1/86 | Mrs. Bullen Swedenborg | 30 00 | Pd | |
| 1/86 | Harold Cramp 1706 Sp. Gard. | 20 00 | Cent. 12/15 | |
| 1/86 | Lieut. H. S. S. S. S. S. | 10 00 | Pd | |
| 1/86 | Storby Carlson 3115 Sp. Garden | 10 00 | | |
| 1/86 | Dr. S. K. K. K. K. | 15 00 | | |
| 1/86 | Capt. J. W. Shanker | 15 00 | Pd 24th | |
| 1/86 | Dr. M. W. S. S. S. | 10 00 | | |
| 1/86 | Dr. M. W. S. S. S. | 10 00 | Jan 14th Pd | |
| 1/86 | Dr. M. W. S. S. S. | 20 00 | Jan 15th Pd | |
| 1/86 | Dr. M. W. S. S. S. | 10 00 | Pd. | |
| 1/86 | Dr. M. W. S. S. S. | 3 00 | Pd | |

Cash Account.

| BILL
SENT. | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID |
|---------------|--------------------------|--------------------|-----------------|--------------|
| Dec | | | | |
| 15 | Mrs. Bley 3021 Grand Ave | | 10 00 | pd |
| 1 | Mr. Shipley 2 d. H det | | 10 00 | Jan 20 |
| 22 | Mr. Hahn 1806 Grand Ave | | 10 00 | Jan 31 |

| BILL
SENT. | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|---------------|-------|--------------------|-----------------|---------------|
|---------------|-------|--------------------|-----------------|---------------|

Cash Account.

231

| BILL
SENT. | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|---------------|-------|--------------------|-----------------|---------------|
|---------------|-------|--------------------|-----------------|---------------|

Jan. 10. 1886.

60

100

0

Nov. 10. May - for 12 mos

70 00

1886-87.

Nov 9. Bill. on letter.

50 00

" 12 Reg letter. 2. 50th

100 00

" 14 May

10 00

| BILL
SENT | NAME | AMOUNT
CHARGED. | AMOUNT
PAID | WHEN
PAID. |
|--------------|------|--------------------|----------------|---------------|
|--------------|------|--------------------|----------------|---------------|

| BILL
SENT. | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID |
|---------------|-----------------------------|--------------------|-----------------|--------------|
| 10/1 | J. B. ... | | 10 | |
| " 12 | Miss Humphreys | | 10 | 00 |
| | Med News 2nd, 94, 114 20.29 | | 26 | |
| 10/2 | College Cheque | | 333 | 84 |
| " 9 | Lee Brothers & Co | | 21 | 90 |
| " | Wm W. Kasper | | 20 | 00 |
| " 11 | Titus. Export trial | | 150 | 00 |
| " | Med News 132, 95, 264 | | 10 | 00 |
| " 16 | Miss Betty Swedesboro | | 30 | 00 |
| " 19 | Levent. Hetherington used | | 16 | |
| " 23 | Mitchell Estate | | 30 | 00 |
| " 24 | Capt. Shackford | | 15 | 00 |
| " | J. B. ... | | 5 | 00 |
| " | ... | | 10 | 00 |
| " | ... | | 15 | 00 |
| " | ... | | 20 | 00 |
| " 11 | ... | | 10 | 00 |
| " | 2026 & 4. ... | | 10 | 00 |
| " | ... | | 10 | 00 |
| " | J. H. ... | | 3 | 00 |
| " | ... | | 10 | 00 |
| " | H. B. Blumenthal | | 10 | 00 |
| | | | 708 | 20 |

235

| BILL | PAYMENT. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|---------|---|--------------------|-----------------|---------------|
| Oct. 11 | Subscription to Charleston Med Soc | | 20.00 | |
| 14 | do do subscription | | 5.00 | |
| | Said art. September | | 57.75 | Nov 1 |
| | Social art. semi-annual exp. | | 30.00 | |
| Nov. 11 | Three Club paper & art- | | 19.25 | Nov 2 |
| " | Haweco master endow art. sub. | | 28.00 | Nov 2 |
| | Chattie. apples | 11/16/ | 25.00 | |

Cash Account.

| BILL
SENT. | NAME | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID |
|---------------|------|--------------------|-----------------|--------------|
|---------------|------|--------------------|-----------------|--------------|

GENERAL MEMORANDA

| | | |
|-----|----------------------|--------|
| 1 | Jan 1 to Jan 31 1898 | 100.00 |
| 2 | Feb 1 to Feb 28 1898 | 100.00 |
| 3 | Mar 1 to Mar 31 1898 | 100.00 |
| 4 | Apr 1 to Apr 30 1898 | 100.00 |
| 5 | May 1 to May 31 1898 | 100.00 |
| 6 | Jun 1 to Jun 30 1898 | 100.00 |
| 7 | Jul 1 to Jul 31 1898 | 100.00 |
| 8 | Aug 1 to Aug 31 1898 | 100.00 |
| 9 | Sep 1 to Sep 30 1898 | 100.00 |
| 10 | Oct 1 to Oct 31 1898 | 100.00 |
| 11 | Nov 1 to Nov 30 1898 | 100.00 |
| 12 | Dec 1 to Dec 31 1898 | 100.00 |
| 13 | Jan 1 to Jan 31 1899 | 100.00 |
| 14 | Feb 1 to Feb 28 1899 | 100.00 |
| 15 | Mar 1 to Mar 31 1899 | 100.00 |
| 16 | Apr 1 to Apr 30 1899 | 100.00 |
| 17 | May 1 to May 31 1899 | 100.00 |
| 18 | Jun 1 to Jun 30 1899 | 100.00 |
| 19 | Jul 1 to Jul 31 1899 | 100.00 |
| 20 | Aug 1 to Aug 31 1899 | 100.00 |
| 21 | Sep 1 to Sep 30 1899 | 100.00 |
| 22 | Oct 1 to Oct 31 1899 | 100.00 |
| 23 | Nov 1 to Nov 30 1899 | 100.00 |
| 24 | Dec 1 to Dec 31 1899 | 100.00 |
| 25 | Jan 1 to Jan 31 1900 | 100.00 |
| 26 | Feb 1 to Feb 28 1900 | 100.00 |
| 27 | Mar 1 to Mar 31 1900 | 100.00 |
| 28 | Apr 1 to Apr 30 1900 | 100.00 |
| 29 | May 1 to May 31 1900 | 100.00 |
| 30 | Jun 1 to Jun 30 1900 | 100.00 |
| 31 | Jul 1 to Jul 31 1900 | 100.00 |
| 32 | Aug 1 to Aug 31 1900 | 100.00 |
| 33 | Sep 1 to Sep 30 1900 | 100.00 |
| 34 | Oct 1 to Oct 31 1900 | 100.00 |
| 35 | Nov 1 to Nov 30 1900 | 100.00 |
| 36 | Dec 1 to Dec 31 1900 | 100.00 |
| 37 | Jan 1 to Jan 31 1901 | 100.00 |
| 38 | Feb 1 to Feb 28 1901 | 100.00 |
| 39 | Mar 1 to Mar 31 1901 | 100.00 |
| 40 | Apr 1 to Apr 30 1901 | 100.00 |
| 41 | May 1 to May 31 1901 | 100.00 |
| 42 | Jun 1 to Jun 30 1901 | 100.00 |
| 43 | Jul 1 to Jul 31 1901 | 100.00 |
| 44 | Aug 1 to Aug 31 1901 | 100.00 |
| 45 | Sep 1 to Sep 30 1901 | 100.00 |
| 46 | Oct 1 to Oct 31 1901 | 100.00 |
| 47 | Nov 1 to Nov 30 1901 | 100.00 |
| 48 | Dec 1 to Dec 31 1901 | 100.00 |
| 49 | Jan 1 to Jan 31 1902 | 100.00 |
| 50 | Feb 1 to Feb 28 1902 | 100.00 |
| 51 | Mar 1 to Mar 31 1902 | 100.00 |
| 52 | Apr 1 to Apr 30 1902 | 100.00 |
| 53 | May 1 to May 31 1902 | 100.00 |
| 54 | Jun 1 to Jun 30 1902 | 100.00 |
| 55 | Jul 1 to Jul 31 1902 | 100.00 |
| 56 | Aug 1 to Aug 31 1902 | 100.00 |
| 57 | Sep 1 to Sep 30 1902 | 100.00 |
| 58 | Oct 1 to Oct 31 1902 | 100.00 |
| 59 | Nov 1 to Nov 30 1902 | 100.00 |
| 60 | Dec 1 to Dec 31 1902 | 100.00 |
| 61 | Jan 1 to Jan 31 1903 | 100.00 |
| 62 | Feb 1 to Feb 28 1903 | 100.00 |
| 63 | Mar 1 to Mar 31 1903 | 100.00 |
| 64 | Apr 1 to Apr 30 1903 | 100.00 |
| 65 | May 1 to May 31 1903 | 100.00 |
| 66 | Jun 1 to Jun 30 1903 | 100.00 |
| 67 | Jul 1 to Jul 31 1903 | 100.00 |
| 68 | Aug 1 to Aug 31 1903 | 100.00 |
| 69 | Sep 1 to Sep 30 1903 | 100.00 |
| 70 | Oct 1 to Oct 31 1903 | 100.00 |
| 71 | Nov 1 to Nov 30 1903 | 100.00 |
| 72 | Dec 1 to Dec 31 1903 | 100.00 |
| 73 | Jan 1 to Jan 31 1904 | 100.00 |
| 74 | Feb 1 to Feb 28 1904 | 100.00 |
| 75 | Mar 1 to Mar 31 1904 | 100.00 |
| 76 | Apr 1 to Apr 30 1904 | 100.00 |
| 77 | May 1 to May 31 1904 | 100.00 |
| 78 | Jun 1 to Jun 30 1904 | 100.00 |
| 79 | Jul 1 to Jul 31 1904 | 100.00 |
| 80 | Aug 1 to Aug 31 1904 | 100.00 |
| 81 | Sep 1 to Sep 30 1904 | 100.00 |
| 82 | Oct 1 to Oct 31 1904 | 100.00 |
| 83 | Nov 1 to Nov 30 1904 | 100.00 |
| 84 | Dec 1 to Dec 31 1904 | 100.00 |
| 85 | Jan 1 to Jan 31 1905 | 100.00 |
| 86 | Feb 1 to Feb 28 1905 | 100.00 |
| 87 | Mar 1 to Mar 31 1905 | 100.00 |
| 88 | Apr 1 to Apr 30 1905 | 100.00 |
| 89 | May 1 to May 31 1905 | 100.00 |
| 90 | Jun 1 to Jun 30 1905 | 100.00 |
| 91 | Jul 1 to Jul 31 1905 | 100.00 |
| 92 | Aug 1 to Aug 31 1905 | 100.00 |
| 93 | Sep 1 to Sep 30 1905 | 100.00 |
| 94 | Oct 1 to Oct 31 1905 | 100.00 |
| 95 | Nov 1 to Nov 30 1905 | 100.00 |
| 96 | Dec 1 to Dec 31 1905 | 100.00 |
| 97 | Jan 1 to Jan 31 1906 | 100.00 |
| 98 | Feb 1 to Feb 28 1906 | 100.00 |
| 99 | Mar 1 to Mar 31 1906 | 100.00 |
| 100 | Apr 1 to Apr 30 1906 | 100.00 |

GENERAL MEMORANDA

SDA
forward

3208

京

1 1/2

44

1891. 10. 22

$\frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$

165

0 2 0

11. 11. 11.

31 Invention. 7:46

10

... 1st. 1st 2nd 3rd 4th 5th 6th 7th 8th 9th 10th 11th 12th 13th 14th 15th 16th 17th 18th 19th 20th 21st 22nd 23rd 24th 25th 26th 27th 28th 29th 30th 31st 32nd 33rd 34th 35th 36th 37th 38th 39th 40th 41st 42nd 43rd 44th 45th 46th 47th 48th 49th 50th 51st 52nd 53rd 54th 55th 56th 57th 58th 59th 60th 61st 62nd 63rd 64th 65th 66th 67th 68th 69th 70th 71st 72nd 73rd 74th 75th 76th 77th 78th 79th 80th 81st 82nd 83rd 84th 85th 86th 87th 88th 89th 90th 91st 92nd 93rd 94th 95th 96th 97th 98th 99th 100th 101st 102nd 103rd 104th 105th 106th 107th 108th 109th 110th 111th 112th 113th 114th 115th 116th 117th 118th 119th 120th 121st 122nd 123rd 124th 125th 126th 127th 128th 129th 130th 131st 132nd 133rd 134th 135th 136th 137th 138th 139th 140th 141st 142nd 143rd 144th 145th 146th 147th 148th 149th 150th 151st 152nd 153rd 154th 155th 156th 157th 158th 159th 160th 161st 162nd 163rd 164th 165th 166th 167th 168th 169th 170th 171st 172nd 173rd 174th 175th 176th 177th 178th 179th 180th 181st 182nd 183rd 184th 185th 186th 187th 188th 189th 190th 191st 192nd 193rd 194th 195th 196th 197th 198th 199th 200th 201st 202nd 203rd 204th 205th 206th 207th 208th 209th 210th 211th 212th 213th 214th 215th 216th 217th 218th 219th 220th 221st 222nd 223rd 224th 225th 226th 227th 228th 229th 230th 231st 232nd 233rd 234th 235th 236th 237th 238th 239th 240th 241st 242nd 243rd 244th 245th 246th 247th 248th 249th 250th 251st 252nd 253rd 254th 255th 256th 257th 258th 259th 260th 261st 262nd 263rd 264th 265th 266th 267th 268th 269th 270th 271st 272nd 273rd 274th 275th 276th 277th 278th 279th 280th 281st 282nd 283rd 284th 285th 286th 287th 288th 289th 290th 291st 292nd 293rd 294th 295th 296th 297th 298th 299th 300th 301st 302nd 303rd 304th 305th 306th 307th 308th 309th 310th 311th 312th 313th 314th 315th 316th 317th 318th 319th 320th 321st 322nd 323rd 324th 325th 326th 327th 328th 329th 330th 331st 332nd 333rd 334th 335th 336th 337th 338th 339th 340th 341st 342nd 343rd 344th 345th 346th 347th 348th 349th 350th 351st 352nd 353rd 354th 355th 356th 357th 358th 359th 360th 361st 362nd 363rd 364th 365th 366th 367th 368th 369th 370th 371st 372nd 373rd 374th 375th 376th 377th 378th 379th 380th 381st 382nd 383rd 384th 385th 386th 387th 388th 389th 390th 391st 392nd 393rd 394th 395th 396th 397th 398th 399th 400th 401st 402nd 403rd 404th 405th 406th 407th 408th 409th 410th 411th 412th 413th 414th 415th 416th 417th 418th 419th 420th 421st 422nd 423rd 424th 425th 426th 427th 428th 429th 430th 431st 432nd 433rd 434th 435th 436th 437th 438th 439th 440th 441st 442nd 443rd 444th 445th 446th 447th 448th 449th 450th 451st 452nd 453rd 454th 455th 456th 457th 458th 459th 460th 461st 462nd 463rd 464th 465th 466th 467th 468th 469th 470th 471st 472nd 473rd 474th 475th 476th 477th 478th 479th 480th 481st 482nd 483rd 484th 485th 486th 487th 488th 489th 490th 491st 492nd 493rd 494th 495th 496th 497th 498th 499th 500th 501st 502nd 503rd 504th 505th 506th 507th 508th 509th 510th 511th 512th 513th 514th 515th 516th 517th 518th 519th 520th 521st 522nd 523rd 524th 525th 526th 527th 528th 529th 530th 531st 532nd 533rd 534th 535th 536th 537th 538th 539th 540th 541st 542nd 543rd 544th 545th 546th 547th 548th 549th 550th 551st 552nd 553rd 554th 555th 556th 557th 558th 559th 560th 561st 562nd 563rd 564th 565th 566th 567th 568th 569th 570th 571st 572nd 573rd 574th 575th 576th 577th 578th 579th 580th 581st 582nd 583rd 584th 585th 586th 587th 588th 589th 590th 591st 592nd 593rd 594th 595th 596th 597th 598th 599th 600th 601st 602nd 603rd 604th 605th 606th 607th 608th 609th 610th 611th 612th 613th 614th 615th 616th 617th 618th 619th 620th 621st 622nd 623rd 624th 625th 626th 627th 628th 629th 630th 631st 632nd 633rd 634th 635th 636th 637th 638th 639th 640th 641st 642nd 643rd 644th 645th 646th 647th 648th 649th 650th 651st 652nd 653rd 654th 655th 656th 657th 658th 659th 660th 661st 662nd 663rd 664th 665th 666th 667th 668th 669th 670th 671st 672nd 673rd 674th 675th 676th 677th 678th 679th 680th 681st 682nd 683rd 684th 685th 686th 687th 688th 689th 690th 691st 692nd 693rd 694th 695th 696th 697th 698th 699th 700th 701st 702nd 703rd 704th 705th 706th 707th 708th 709th 710th 711th 712th 713th 714th 715th 716th 717th 718th 719th 720th 721st 722nd 723rd 724th 725th 726th 727th 728th 729th 730th 731st 732nd 733rd 734th 735th 736th 737th 738th 739th 740th 741st 742nd 743rd 744th 745th 746th 747th 748th 749th 750th 751st 752nd 753rd 754th 755th 756th 757th 758th 759th 760th 761st 762nd 763rd 764th 765th 766th 767th 768th 769th 770th 771st 772nd 773rd 774th 775th 776th 777th 778th 779th 780th 781st 782nd 783rd 784th 785th 786th 787th 788th 789th 790th 791st 792nd 793rd 794th 795th 796th 797th 798th 799th 800th 801st 802nd 803rd 804th 805th 806th 807th 808th 809th 810th 811th 812th 813th 814th 815th 816th 817th 818th 819th 820th 821st 822nd 823rd 824th 825th 826th 827th 828th 829th 830th 831st 832nd 833rd 834th 835th 836th 837th 838th 839th 8

1870

2

3

1

" " N. I. M. don

135

1 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 104

1

3.

1-19

" 18 New Holger

10

2. The 2 hours 30 min. Sept. 8. 1895.

20

July 1 and 2, 1891

15

" 6 } " has been over

5.

" 7 " then belly

1

H. W. Abbott

72

Mr Hawkins

15.

15

Wa Trau

300

11

W. L. Luchman

3

$\frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2}$

GENERAL MEMORANDA

July

Harvard \$ 50.50.00

23rd Chas. a. Lige. 3.00

Sept.

7. Mrs. Isollen 10.00

9. Med. Recs. 25.00. Rec. in 20.24.25 15.00

13. I. Hoffman 15.00

620.00

GENERAL MEMORANDA

| | |
|-----------------|-------|
| also Patent 812 | 5.00 |
| Johnson | 0.00 |
| 1878 2nd | 5.00 |
| 1878 3rd | 10.00 |
| 1878 4th | 10.00 |
| 1878 5th | 10.00 |
| 1878 6th | 10.00 |
| 1878 7th | 10.00 |
| 1878 8th | 10.00 |
| 1878 9th | 10.00 |
| 1878 10th | 10.00 |
| 1878 11th | 10.00 |
| 1878 12th | 10.00 |
| 1878 13th | 10.00 |
| 1878 14th | 10.00 |
| 1878 15th | 10.00 |
| 1878 16th | 10.00 |
| 1878 17th | 10.00 |
| 1878 18th | 10.00 |
| 1878 19th | 10.00 |
| 1878 20th | 10.00 |
| 1878 21st | 10.00 |
| 1878 22nd | 10.00 |
| 1878 23rd | 10.00 |
| 1878 24th | 10.00 |
| 1878 25th | 10.00 |
| 1878 26th | 10.00 |
| 1878 27th | 10.00 |
| 1878 28th | 10.00 |
| 1878 29th | 10.00 |
| 1878 30th | 10.00 |
| 1878 31st | 10.00 |
| 1878 32nd | 10.00 |
| 1878 33rd | 10.00 |
| 1878 34th | 10.00 |
| 1878 35th | 10.00 |
| 1878 36th | 10.00 |
| 1878 37th | 10.00 |
| 1878 38th | 10.00 |
| 1878 39th | 10.00 |
| 1878 40th | 10.00 |
| 1878 41st | 10.00 |
| 1878 42nd | 10.00 |
| 1878 43rd | 10.00 |
| 1878 44th | 10.00 |
| 1878 45th | 10.00 |
| 1878 46th | 10.00 |
| 1878 47th | 10.00 |
| 1878 48th | 10.00 |
| 1878 49th | 10.00 |
| 1878 50th | 10.00 |
| 1878 51st | 10.00 |
| 1878 52nd | 10.00 |
| 1878 53rd | 10.00 |
| 1878 54th | 10.00 |
| 1878 55th | 10.00 |
| 1878 56th | 10.00 |
| 1878 57th | 10.00 |
| 1878 58th | 10.00 |
| 1878 59th | 10.00 |
| 1878 60th | 10.00 |
| 1878 61st | 10.00 |
| 1878 62nd | 10.00 |
| 1878 63rd | 10.00 |
| 1878 64th | 10.00 |
| 1878 65th | 10.00 |
| 1878 66th | 10.00 |
| 1878 67th | 10.00 |
| 1878 68th | 10.00 |
| 1878 69th | 10.00 |
| 1878 70th | 10.00 |
| 1878 71st | 10.00 |
| 1878 72nd | 10.00 |
| 1878 73rd | 10.00 |
| 1878 74th | 10.00 |
| 1878 75th | 10.00 |
| 1878 76th | 10.00 |
| 1878 77th | 10.00 |
| 1878 78th | 10.00 |
| 1878 79th | 10.00 |
| 1878 80th | 10.00 |
| 1878 81st | 10.00 |
| 1878 82nd | 10.00 |
| 1878 83rd | 10.00 |
| 1878 84th | 10.00 |
| 1878 85th | 10.00 |
| 1878 86th | 10.00 |
| 1878 87th | 10.00 |
| 1878 88th | 10.00 |
| 1878 89th | 10.00 |
| 1878 90th | 10.00 |
| 1878 91st | 10.00 |
| 1878 92nd | 10.00 |
| 1878 93rd | 10.00 |
| 1878 94th | 10.00 |
| 1878 95th | 10.00 |
| 1878 96th | 10.00 |
| 1878 97th | 10.00 |
| 1878 98th | 10.00 |
| 1878 99th | 10.00 |
| 1878 100th | 10.00 |

495

The Medical News.

A NATIONAL WEEKLY PERIODICAL, CONTAINING 28 TO 32
QUARTO PAGES IN EACH ISSUE.

The continually increasing appreciation of THE MEDICAL NEWS by the profession throughout the country, encourages its managers to make, during the coming year, even more strenuous efforts to enhance its value to the practitioner.

Possessing a most efficient organization, THE NEWS unites the best features of the medical magazine and newspaper. Its large and able Editorial Staff discusses in each issue the important topics of the day in a thoughtful and scholarly manner, while its corps of qualified reporters and correspondents, covering every medical centre, insures that its readers shall be promptly and thoroughly posted upon all matters of interest in the world of medicine. On account of the position conceded to THE NEWS, it has become the medium chosen by the leading minds of the profession for the publication of their most important Original Contributions to medical science. The valuable instruction afforded in Clinical Lectures, and the rich experience gained in the leading Hospitals of the world are constantly laid before the readers of THE NEWS, and prompt and authentic reports of Society Proceedings are received from special reporters in various sections of the country by mail and telegraph. In the pages devoted to the Progress of Medical Science are found early notes of all important advances, gleaned from the principal journals of both hemispheres.

In short, every branch of medicine is adequately represented

LEA BROTHERS & CO., Publishers, Philadelphia.

MEDICAL PERIODICALS—1886.

in THE NEWS, and the details of plan and typography have been carefully studied in order to economize the time and secure the comfort of the reader in every possible way.

Price, FIVE DOLLARS, per Annum, In Advance.

* * * * *

COMMUTATION RATE.

To subscribers paying in advance for 1886: THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES (quarterly), and THE MEDICAL NEWS (weekly), to one address for \$9.00 per annum.

SPECIAL OFFERS.

THE MEDICAL NEWS VISITING LIST, arranged for 30 visits daily, may be obtained by any advance paying subscriber to either of the above periodicals, free by mail, by forwarding 75 cents (regular price \$1.00). THE MEDICAL NEWS VISITING LIST will be furnished with a Thumb-letter Index for quick reference, for 25 cents additional.

THE YEAR BOOK OF TREATMENT for 1885 (a complete abstract of the year's therapeutic advances, by a large number of eminent specialists), may be obtained by any advance paying subscriber to either of the above periodicals, free by mail, by forwarding 75 cents (regular price \$1.25).

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, THE MEDICAL NEWS, THE MEDICAL NEWS VISITING LIST, and THE YEAR BOOK OF TREATMENT for 1885, will be sent to one address, free by mail, for \$10.00, in advance.

Subscribers can obtain, at the close of each volume, cloth covers for THE JOURNAL (one annually), and for THE NEWS (one annually), free by mail, by remitting Ten Cents for THE JOURNAL cover, and Fifteen Cents for THE NEWS cover.

LEA BROTHERS & CO., Publishers, Philadelphia.

THE
MEDICAL NEWS
VISITING-LIST.

1887.

THIRTY PATIENTS PER WEEK.



PHILADELPHIA:
LEA BROTHERS & CO.
1886.

Entered according to Act of Congress, in the year 1886, by
LEA BROTHERS & CO.,
in the Office of the Librarian of Congress. All rights reserved.

WESTCOTT & THOMSON,
Stereotypers and Electrotypers, Philada.

WILLIAM J. DORNAN,
Printer, Philada

CONTENTS.

| | PAGE |
|--|---------|
| PREFACE | 5 |
| HOW TO KEEP ACCOUNTS | 7 |
| HOW TO KEEP THE VISITING-LIST | 7 |
| TABLE TO FIND DAY OF CONFINEMENT | 8 |
| SIGNS OF PREGNANCY | 8 |
| SIGNS OF DENTITION | 8 |
| WEIGHTS AND MEASURES (ORDINARY AND METRIC) | 9-10 |
| COMPARATIVE SCALES | 11 |
| EXAMINATION OF URINE | 12-13 |
| DISINFECTANTS | 13 |
| TABLE OF ERUPTIVE FEVERS | 14 |
| SOME REMEDIES NOT IN GENERAL USE | 15-18 |
| INCOMPATIBLES | 18 |
| ARTIFICIAL RESPIRATION | 19 |
| POISONS AND ANTIDOTES | 20-21 |
| TABLE OF DOSES | 22-26 |
| THERAPEUTIC TABLE | 27-43 |
| LIGATION OF ARTERIES | 43-48 |
| <hr/> | |
| DAILY RECORD OF PRACTICE | 50-155 |
| CLINICAL RECORD | 156-183 |
| CONSULTATION PRACTICE | 184-189 |
| OBSTETRIC ENGAGEMENTS | 190-195 |
| OBSTETRIC PRACTICE | 196-207 |
| VACCINATIONS | 208-211 |
| DEATHS | 212-215 |
| ADDRESSES OF PATIENTS | 216-219 |
| ADDRESSES OF NURSES | 220-221 |
| CASH ACCOUNT | 222-236 |
| GENERAL MEMORANDA | 237-240 |



PREFACE.

THIS Visiting List has been arranged so as to include a number of desirable features, some of which are now to be found scattered about in a variety of similar books, others of which are here presented for the first time. A visiting list should not be expected to contain a treatise on the practice of medicine. It should, however, furnish brief memoranda which may be useful in an emergency, or in the moments, which come to the best of men, when one cannot recall a fact with which he may be perfectly familiar. A visiting list should also be so arranged as to permit a physician to keep a clear record of his engagements and of his professional work, from which he may transfer to a ledger a permanent record which will have a legal status in case of a dispute. Both these objects have been aimed at in this Visiting List, the purchasers of which are strongly urged to complete their records by the use of a suitable ledger, since the imperfect accounts of visiting lists have been decided to have no legal value as evidence in the collection of debts.

The Visiting List for 1887 has been revised and improved. The list of remedies not in general use and the therapeutic table have been brought up to date by omitting some remedies which have not justified the expectation concerning them, and by inserting others which have been introduced during the last year.

In response to many requests, it is now issued in various sizes, to accommodate practices of thirty, sixty, and ninety patients per week, and another edition is published without dates (perpetual). Every requirement of the profession, it is believed, has now been met.

Introduced to the profession toward the close of 1885. The Medical News Visiting List was regarded by its projectors as an experiment, and the result was watched with considerable interest. The verdict on its popularity was not long in suspense, and the little venture is therefore established on a permanent basis.



HOW TO KEEP ACCOUNTS.

THOUGH most medical men have never had any business training, they will be able, by a careful observation of the two following suggestions, to keep accounts which will be accurate and valid in a court of law:

1. Keep your visiting list in such a way that you can fully explain it.
2. Transfer a record of your services at short intervals to a ledger, stating the items.

HOW TO KEEP THE VISITING LIST.

ENTER the names in the proper columns, and, in case of a new patient, the address (or husband's or parent's name and address) in the right-hand page under the head of "Memoranda." Under the proper date make a little dot • in lead pencil for a visit *to be paid*, and if at a special hour, write this down: 10 A. M. When a visit is paid make a diagonal stroke in ink, /. When several visits are paid a series of strokes, ///, may be made, or one with a little star */. and in the memorandum column the number of visits, "*4 visits," may be written in. A night-visit should be marked *N*. For an expected office case a small circle in lead pencil, ○, should be made, and when the patient has come an *O* may be written in ink. For a consultation write **C*, and in the "Memoranda" "Consultation with Dr. —" (whoever asked you in consultation). When you call a consultant mark the visit */. and in the "Memoranda" column write "*Dr. — in consultation." A confinement case may be marked *Cf.*, and anything special noted in column for memoranda. A vaccination should be marked *V*.

SIGNS.


| | |
|-----|--|
| • | Visit to be made. |
| / | Visit made. |
| // | Second Visit to be made same day. |
| // | Second Visit made. |
| N | Night Visit. |
| ○ | Office Visit expected. |
| O | Office Visit. |
| C | Consultation (when you are the consultant). |
| Cf. | Confinement Case. |
| V | Vaccination. |
| | Refers to a special note in column headed "Memoranda." |

TABLE TO FIND DAY OF CONFINEMENT.

AROUND the circle the months of the year are arranged in their order, with the number of days in each month. The number placed below each month gives the number of days that must be added to the nine preceding months in order to make up 280 days. Rule: Count backward three months from the last day of menstruation, and add to the date thus reached the number below that month.

Example: Last day of menstruation, Feb. 10th. Count backward three months to Nov. 10th and add 7 days = Nov. 17th.

It should never be forgotten that an exact reckoning of the date of confinement is impossible, errors of one or two weeks being sometimes made. For this reason it is not worth while to make any difference in the calculation for the extra day which falls in each leap year.



SIGNS OF PREGNANCY.

| TERM OF PREGNANCY. | Morning Sickness. | Suppression of the Menses. | Mammary Areola. | Enlargement of the Abdomen. | Fetal Movements. | Ballotement. | Uterine Souffle. | Fetal Heart Sounds. | Dusky hue of the Vagina. |
|----------------------|-------------------|----------------------------|-----------------|-----------------------------|------------------|--------------|------------------|---------------------|--------------------------|
| End of First Month.. | + | + | + | + | + | + | + | + | + |
| " Second " | + | + | + | + | + | + | + | + | + |
| " Third " | + | + | + | + | + | + | + | + | + |
| " Fourth " | + | + | + | + | + | + | + | + | + |
| " Fifth " | + | + | + | + | + | + | + | + | + |
| " Sixth " | + | + | + | + | + | + | + | + | + |
| " Seventh " | + | + | + | + | + | + | + | + | + |
| " Eighth " | + | + | + | + | + | + | + | + | + |
| " Ninth " | + | + | + | + | + | + | + | + | + |

DENTITION.

The first teeth usually appear at about the seventh month, but they may be cut earlier or later without any presumption of disease. The first or temporary set of teeth numbers twenty, and appears approximately as follows (the numbers indicate months):

| | | | | |
|---------|---------|-----------|---------|---------|
| Molars. | Canine. | Incisors. | Canine. | Molars. |
| 24-12 | 18 | 9-7-7-9 | 18 | 12-24 |

The permanent teeth appear about as follows (the numbers indicate years):

| | | | | | | |
|---------|-------------|---------|-----------|---------|-------------|---------|
| Molars. | Bicuspsids. | Canine. | Incisors. | Canine. | Bicuspsids. | Molars. |
| 25-13-6 | 10-9 | 11 | 8 7-7-8 | 11 | 9-10 | 6 13-25 |

THERMOMETRIC SCALES.

To convert Degrees of the Centigrade Scale into those of Fahrenheit: Multiply the number of the former by 9 and divide by 5; if the temperature was above the freezing-point (0°), add 32° ; if it was below the freezing-point, subtract the result from 32° .

To convert Degrees of the Fahrenheit Scale into those of Centigrade: If the temperature be above the freezing-point (32°), subtract 32; if it be below the freezing-point, subtract the number from 32 then multiply the remainder by 5 and divide by 9.

In taking temperatures, the thermometer should always be used in the same situation (axilla, mouth, etc.), and left in for the same length of time. It is important also, in keeping a record or making a report, to state the situation of the thermometer.

ORDINARY WEIGHTS AND MEASURES.

Troy, or Apothecaries', Weight.

| | | |
|-----------------|-----------------|--------------|
| 20 grains (gr.) | = 1 scruple (℥) | = 20 grains. |
| 3 scruples | = 1 drachm (ʒ) | = 60 " |
| 8 drachms | = 1 ounce (℥) | = 480 " |
| 12 ounces | = 1 pound (lb.) | = 5760 " |

Wine, or Apothecaries', Measure.

| | |
|------------------------|--|
| 60 minims (℥) | = 1 fluidrachm. |
| 8 fluidrachms (fldr.) | = 1 fluidounce. |
| 16 fluidounces (flox.) | = 1 pint. |
| 2 pints (O) | = 1 quart. |
| 4 quarts | = 1 gallon (Cong.) = 231 cubic inches. |

Linear Measure.

| | |
|-----------|-----------|
| 12 lines | = 1 inch. |
| 12 inches | = 1 foot. |

THE METRIC SYSTEM.

THIS system gets its name from the meter, which is the unit of linear measure. It is equal to 39.370432 inches, or a little over three and one-quarter feet.

The gram is the unit of weight, and is equal to 15.434 grains (Troy).

The liter is the unit of solid or fluid measure, and is equal to 1.0567 quarts.

The names of the higher weights and measures are formed by prefixing to the name of the unit the Greek terms deka, hecto, kilo, and myria. The names of the lower weights and measures are formed by prefixing to the name of the unit the Latin terms deci, centi, milli.

TABLE OF LINEAR MEASURES.

| | | |
|----------------------|-------------------------|-----------------------|
| 1 millimeter (m.m.) | = 1 : 1000 of a meter = | 0.03937 inch. |
| 1 centimeter (c.m.) | = 1 : 100 " | 0.3937 " |
| 1 decimeter (d.c.m.) | = 1 : 10 " | 3.937 " |
| 1 meter (m.) | | 39.37 " |
| 1 dekameter (d.k.m.) | = | = 32 feet 9.7 inches. |

METRIC WEIGHTS AND MEASURES.

Length :

| | | |
|------------------------|-------------------------|-----------------------------|
| 1 millimeter | | .0393704 inches. |
| 10 millimeters (m.m.) | = 1 centimeter | = .393704 " |
| 10 centimeters (c.m.) | = 1 decimeter | = 3.937043 " |
| 10 decimeters (d.c.m.) | = 1 Meter | = 39.370432 " |
| 10 meters (m.) | = 1 dekameter | = 32 feet 9.7 inches. |
| 10 dekameters (dkm.) | = 1 hektometer | = 328 " 1. " |
| 10 hektometers (h.m.) | = 1 kilometer | = 3280 " 10.4 " |
| 10 kilometers (k.m.) | = 1 myriameter (myr.m.) | = 6.2137 miles. |

Volume :

| | | |
|-----------------------|--|---|
| 1 milliliter (m.l.) | = 1 cubic centimeter | = 16.23 minims. |
| 10 milliliters | = 1 centiliter | = 2.71 fluidrachms. |
| 10 centiliters (c.l.) | = 1 deciliter | = 3.38 fluidounces. |
| 10 deciliters (dc.l.) | = 1 Liter (= 1 cubic decimeter) | = 33.81 fluidounces =
1.0567 quarts. |
| 10 liters (l.) | = 1 dekaliter | = 2.6417 gallons. |
| 10 dekaliters (dk.l.) | = 1 hektoliter | = 26.417 " |
| 10 hektoliters (h.l.) | = 1 kiloliter or stere | = 264.17 " |


Weight :

| | | |
|-----------------------|--|---------------------------------------|
| 1 milligram (m.gm.) | = .001 gram | = .015 grain Troy. |
| 10 " | = 1 centigram | = .154 " " |
| 10 centigrams (c.gm.) | = 1 decigram | = 1.543 " " |
| 10 decigrams (dc.gm.) | = 1 Gram | = 15.432 " " |
| 1 gram (gm.) | weight of 1 cubic centimeter of water at 104° F. | |
| 10 grams | = 1 dekagram | = 25 34.32 grains Troy. |
| 10 dekagrams (dk.gm.) | = 1 hektogram | = 35 103.235 " " |
| 10 hektograms (h.gm.) | = 1 kilogram (k.gm.) | = 2 lb. 83 72 " " |

A teaspoon holds about 5 cubic centimeters, and a tablespoon about 20 cubic centimeters. A Troy ounce equals about 30 grams, and an eight-ounce bottle holds about 250 cubic centimeters.

In prescribing, the French usually confine themselves to ordering grams and multiples of the gram for weights, and cubic centimeters and multiples of the cubic centimeter in measures.

COMPARATIVE SCALES, showing at a glance the exact equivalent of ordinary weights and measures in those of the Metric System, and vice versa.



The equivalents of fractions, whether large or small, may be found with great nicety by these scales. For instance, $\frac{1}{2}$ grain = $\frac{1}{4}$ of the metric equivalent of 7 grains, and 1-200 grain 1-4000 of the metric equivalent of 20 grains. This method is, of course, reversible.

EXAMINATION OF THE URINE.

THE most important matters to be investigated in an examination of the urine are—1, the amount passed in a given time; 2, the specific gravity; 3, the reaction; 4, the proportion of urates present; 5, the proportion of phosphates present; 6, the absence or presence of albumen, and the quantity if present; 7, the absence or presence of sugar, and the quantity if present; and, 8, the character of the microscopic sediment.

The examination may be conducted as follows:

1. The amount may be measured by collecting the whole amount passed in a given 24 hours.

2. The specific gravity may be ascertained with a "urinometer."

3. The reaction should be ascertained, immediately or soon after the passage of the sample, by means of slips of blue litmus-paper, which is turned red by acids, or by reddened litmus-paper, which is made blue by alkalies, or by turmeric-paper, which is made brown by alkalies.

4. The proportion of urates may be very well estimated by the amount of sediment and turbidity in an *acid* urine. A high specific gravity in an acid urine usually indicates the presence of a proportionate excess of urates. A turbidity which is cleared up by warming the urine also indicates an excess of urates.

5. The proportion of phosphates may be ascertained by adding to some urine, in a test-tube, half its bulk of a solution of caustic potash (1:8), and gently warming the mixture. If the deposit which forms is deeper than a third of an inch in a test-tube three-quarters of an inch wide with two inches of the mixture in it, the phosphates are in excess. If much less, they are below normal. Turbidity of the urine, which is cleared up by the addition of a little acetic acid, is to be set down to an excess of phosphates—unless the urine is stale, when it may be due to alkaline decomposition.

6. The presence of albumen is detected by the appearance of a whitish cloudiness or a curdy deposit upon boiling a little clear urine, after slightly acidulating it with a *few drops* of acetic acid. Another good test is to pour a little urine into a test-tube, then to draw up some nitric acid into a long, pointed glass tube, close the upper end with a forefinger, thrust the point to the bottom of the urine, and then withdraw the finger gradually, so as to permit the acid to flow out slowly at the bottom, under the urine, which will be raised up as the acid flows out. If there be albumen present, it will be disclosed by the formation of a layer of coagulation between the acid and the urine. A light, cloudy appearance a little higher than the line where the acid and the urine are in contact is to be attributed to a precipitation of urates by the acid, and is not a sign of albumen, which coagulates just at the line of contact of acid and urine.

7. The presence of sugar, in quantity sufficient to have any pathological significance, is easily ascertained by adding to some urine, in a test-tube, half its bulk of solution of caustic potash 1:8, and boiling the mixture thoroughly. Any noteworthy amount of sugar will cause a distinct deepening of the color, even to making it dark brown or black if the proportion of sugar be very large. A slight change (to a rum color) does not

indicate the presence of sugar, but occurs when there is none or only a physiological quantity present. After the boiling, some muriatic acid may be added, when, if the amount of sugar be large, the odor of burnt sugar may be perceived. A high specific gravity in a clear, syrupy-looking urine is apt to go with the presence of sugar in it.

8. The character of the sediment as revealed by the microscope is of more significance than any other recognizable characteristic of the urine in disease. To conduct this part of the examination requires a good microscope and the ability to recognize the various sedimentary bodies to be found in health or disease.

DISINFECTANTS.

Clothing, Bedding, etc.—If these are not of too much value, the best and simplest method of disinfection is to burn them. If they are worth saving, they may be subjected to prolonged baking or boiling—that is to say to dry or moist heat—which will effectually destroy any germs which they may hold. Or they may be exposed to the fumes of sulphur, or soaked in a solution of corrosive sublimate and permanganate of potassium, of each 1 ounce in 8 gallons of water.

Bed-pans, Chamber-pots, Water-closets, etc. These should be washed with boiling-hot water and rinsed with a solution of corrosive sublimate and permanganate of potassium, of each 2 drachms in a gallon of water. A quantity of this solution should always be left in each vessel. According to the most recent investigations, this is the most efficient disinfectant for fecal discharges which we have.

Rooms.—Thorough cleanliness, plenty of fresh air, and sunlight are the most important purifiers of inhabited rooms. To these may sometimes be added the exposure of chloride of lime in saucers. When a room can be given over to thorough disinfection, this may be done by removing all curtains and carpets (which must be disinfected in the manner described for clothing), washing all the solid parts with hot water and soft soap, drying thoroughly, then shutting all doors and windows and filling the room with the fumes of burning sulphur. Fumigation with sulphur is effected by placing tubs of water in the room and standing in the midst of the water a brick, and upon this an earthenware saucer (like that for a flower-pot) in which some sulphur has been placed. From 1 to 3 pounds of sulphur should be allowed for each 1000 cubic feet of air-space. The room should be kept closed for 24 hours, and then thoroughly aired.

Water-closets, Drains, and Sinks.—To disinfect these it is important to cleanse them as well as possible, and the best way to do this is to use a strong solution of lye. There are a number of preparations which may be used, one of which is well known as "Banner Lye." A strong solution of this may be poured into the water-closet pan and allowed to remain there for some time, or into the drain-pipes and sinks, and managed so as to remain in contact with the polluted parts as long as possible. After this the solution of corrosive sublimate and permanganate of potassium recommended for bed-pans may be used.

Privies.—The contents of privies should be frequently and well strewn with chloride of lime.

Carbolic acid, which is often used for disinfection, is a very weak and unreliable agent, except when used in strong concentration. As sometimes used, by being exposed in saucers in sick rooms, it has no value whatsoever.

TABLE OF ERUPTIVE FEVERS.

| Disease. | Manner of Invasion. | Characteristics of Eruption. | Course of Disease. |
|------------------------------|---|--|---|
| Scarlatina
Scarlet fever. | Usually sudden; chilliness; high fever; rapid pulse; flushed face; convulsions or coma; sore throat; "strawberry tongue." | Appears very early, often on second day. First on neck and chest. May become universal. May be only about joints and very trifling. Bright scarlet; punctate. Followed by desquamation; skin burning hot; sometimes itches. | Progresses steadily and rapidly. Temperature rises to 104° 106°. Defervescence gradual. Coma a dangerous sign. Albumen in urine common, and often troubles patient five or six days. |
| Rotheln
(German measles). | Usually <i>unaccompanied</i> . Slight fever or none; mild catarrhal symptoms; vomiting; slight sore throat; enlargement of cervical lymphatic glands almost always present. | Appears early. Often first symptom noted, first on face and neck, then on trunk. Rosy papules, scattered or clustered. Fades in one part and appears in another. Lasts about three days. | Usually mild. Disappears gradually. |
| Varicella
(Chicken-pox). | Usually <i>unaccompanied</i> . Slight fever or none; slight malaise or none. | Appears early. Usually on chest and shoulders, then on body and limbs and face. Distinct, clear, shot-sized vesicles; sometimes with distinct areolae and even umbilicated; appear in successive crops, usually few in number. | Vesicles disappear by drying up. May form pits and leave pits. |
| Varicella
(Smallpox). | Usually sudden; chill; high fever; rapid pulse; violent pain in head and back; debility; nut cons membranes often involved. | Appears on third or fourth day: first on face and spreads over whole body and limbs in a day or two. Fine shot-like papules; then vesicles; then umbilicated papules about eighth day developing adherent crusts; well marked areolae. Intense itching. Peculiar odor. | Symptoms sometimes moderate when eruption first appears, and grow worse as pustules mature. Often severe constitutional disturbance, great restlessness and desire to scratch; hemorrhage, defervescence gradual and slow. May be interrupted or exuberant. Crusts fall off, and often leave deep pits (pockmarks). Usually mild and regular. Temperature may be high from complications. Lasts about a week. Recovery rapid. |
| Morbilli
(Measles). | Generally sudden; chilliness or even convulsions; moderate fever; lachrymation; catarrh of respiratory passages; bronchitis. | Appears about fourth day; first on face, and spreads to body and limbs. May be limited; usually almost universal. Consists of rosy papules, arranged in parts of circles (crescentic); appears sometimes in successive crops. | |

SOME REMEDIES NOT IN GENERAL USE.

Adonidine.—The active principle of the *Adonis vernalis*, given in doses of $\frac{1}{8}$ – $\frac{1}{4}$ grain F. d. in pill, is a useful substitute for digitalis, and free from its bad effects.

Apiol has been found useful in the treatment of amenorrhœa: given in capsules, in doses of 3 grains F. d.

Antipyrine.—An artificial hydrocarbon, soluble in three parts of cold and one-half part of warm water. Useful as an antipyretic, especially in typhoid fever and rheumatism, although it does not modify the course of the disease. Does not produce deafness or dizziness. Best excipient is syrup of tolu. Dose, usually 20–30 grains at first, followed by 10-grain doses hourly until temperature has fallen decidedly.

Calx sulphurata.—Generally known as sulphide of calcium. Useful for boils and for acne. To be given in doses of $\frac{1}{8}$ grain, with an equal quantity of sugar of milk, every hour. Doses of $\frac{1}{4}$ grain have been given.

Cascara sagrada.—A laxative much praised in the treatment of chronic constipation. May be prescribed in solid or fluid extract. Dose of the former, 1–2 grains: of the latter, 10–60 minims.

Chrysarobin (Chrysophanic acid) is used in the treatment of skin diseases, especially stubborn cases of psoriasis and tinea capitis. It is used in ointment (strength 10 per cent.) and in solution in chloroform (7 grains to the fluidounce), to be well rubbed in. It makes a very indelible stain if it comes into contact with clothing.

Cocaine.—The alkaloid of the *Erythroxylon coca*. A valuable local anæsthetic, producing contraction of small bloodvessels, and so having hæmostatic properties. In the eye it causes dilatation of the pupil, lachrymation, and slight paralysis of accommodation. It is a good application to burns and for the pain and heat of rhus-poisoning. It has been used for coryza, hay fever, tonsillitis, ear-ache, piles, and irritation of the bladder and urethra. It may be applied to epithelioma or lupus before using arsenic or other caustic. For sea sickness it has been used successfully in doses of $\frac{1}{4}$ – $\frac{1}{2}$ grain in water: and also in teaspoonful doses of a solution of 2½ grains in 5 fluidounces of water, given every 2–3 hours. It may be applied to the skin or mucous membrane, or injected hypodermically, before minor operations.

Duboisia, Duboisina. The sulphate or hydrochlorate is used to dilate the pupil and to paralyze accommodation. Its effects are more prompt and less lasting than those of atropia. Used in the strength of 2–4 grains to the fluidounce of water. It often produces disagreeable, though transient, cerebral disturbance.

Eucalyptus, Oil of. is a valuable stimulating expectorant in doses of from 2–5 drops. It has also been used as an antiseptic by external application and by inhalation. For internal administration it can be put in capsules. 10 minims every 4 hours have proved useful in typhoid fever.

Euonymin is the active principle of the *Euonymus atropurpureus*, and is a valuable cholagogue. The dose is 2 grains, to be followed by an aperient the next morning.

Grindelia robusta.—Used in the form of fluid extract, dose 15 minims to a fluidrachm, for the treatment of dyspnoea dependent on heart disease, as well as asthma, hay asthma, and the asthmatic attacks of bronchitis, whooping cough, and emphysema.

Hamamelis (Witch-Hazel).—Used in the form of the fluid extract, dose 1-2 fluidrachms, as a hæmostatic and to give tone to the muscular walls of the veins. Has been used in internal hemorrhages and to cure varicose veins and varicose ulcers of the leg.

Homatropine Hydrobromate.—A safe substitute for atropia to dilate the pupil (for which instillation of a few drops of a solution of 1-2 grains in a fluidounce of water is sufficient), and to paralyze the muscles of accommodation (for which a strength of 24 grains to the fluidounce of water is required).

Hydrogen. Peroxide of, is a transparent oily liquid, commonly used in watery solution of 10-15 per cent. It has a high oxidizing power, and when injected into septic cavities is decomposed, the extra atom of oxygen attacking the septic material and disinfecting it. It has also been recommended for the treatment of diphtheria.

Hyoscin Hydrobromate.—Useful in insomnia of melancholia, acute delirious mania, alcoholism, neurasthenia, etc., to calm excitement in mental disorders, and in chorea and epilepsy. A good formula is: Hyosceine hydrobrom. gr. j; aq. distil. f3ix; alcohol f3j. 10 minims = $\frac{1}{16}$ gr. Dose, 3-7½ minims = $\frac{1}{16}$ to $\frac{1}{8}$ grain.

Iridin.—An extract of the blue flag. A useful cholagogue, like euonymin. Dose, 4 grains.

Lactic Acid.—Useful as a caustic for lupus and epithelioma. The surrounding part being protected with wax plaster, cotton or lint dipped in a saturated solution of lactic acid may be applied to the morbid tissue. In 12 hours this will be soft and pulpy, when it should be dressed with cold water for 24 or 48 hours. This process may be repeated until a cure is effected.

Naphthalin has been used as an antiseptic, and has been highly recommended in doses of 2 grains 3-4 times a day to prevent decomposition of the urine.

Nitro-glycerine.—Useful in relieving the pain of neuralgia and angina pectoris, as well as headache, vertigo, and epilepsy. It reduces albumen in nephritis, but should not be used when there is hypertrophy of the heart. It has been used in hiccough. It is usually administered in a 1 per cent. solution (water or alcohol), the dose beginning with 1 drop, and being increased up to 10 or more drops thrice daily, as it can be tolerated. It may be administered in pill form, made up with cacao-butter, and it has also been made into granules or tablets.

Nitrite of Amyl has effects similar to those of nitro-glycerine, but less lasting. Administered by inhalation, a few drops being poured on a

thin cloth and held before the nose, and withdrawn when the face becomes distinctly flushed or relief has been afforded. It is believed to be of value in averting attacks of epilepsy and to relieve whooping cough and the syncope of chloroform-poisoning. Can be given internally. $\mathcal{R}$. Amyl nitrite, $\mathcal{M}$. XXXV; ethylic alcohol, f3v; glycerin, q.s. ad f3jss. Dose, f3j in warm water f3ij.

Oleates.—Under this name certain combinations of metals and alkalis with oleic acid have been urged upon the profession. Of these, few are of much value. The oleate of zinc, the oleate of copper, and the oleate of mercury are excellent preparations. There is no evidence that oleates are better than the old preparations with lard when absorption of the drug is desirable.

Paraldehyde.—Used as a hypnotic in conditions of high excitement, as mania, produces sleep without tendency to coma, it being possible to arouse the patient at once, while he falls asleep again when let alone. Given with syrup of orange or a simple elixir; so that a tablespoonful of the mixture shall contain from 25 minims to a fluidrachm of paraldehyde. It does not depress the heart.

Permanganate of Potassium has been found useful in the treatment of amenorrhœa. It is given in pill form, combined with some non-oxidizable vehicle, such as kaolin (ointment). The dose is from 1-3 grains thrice daily, to be given before the time for menstruating.

Physostigma (Calabar Bean).—Used in the form of a tincture (dose, 12 minims) and an extract (dose, from $\frac{1}{2}$ grain). Useful in tetanus, strychnia-poisoning, paralysis of the insane, paraplegia, and locomotor ataxia. Also used in constipation due to atony of the muscular coat of the bowels. Sulphate of eserine (the active principle) is used, in $\frac{1}{4}$ -1 per cent. watery solution, by instillation into the eye, for glaucoma, staphylo ma, and to neutralize the effect of mydriatics. It often causes sharp supraorbital and nasal pain for half an hour after the application.

Picrotoxin has been highly recommended, in doses of $\frac{1}{10}$ - $\frac{1}{20}$ grain, for the night-sweats of phthisis.

Pilocarpine, the active principle of jaborandi, used in the form of the hydrochlorate, in doses of from $\frac{1}{2}$ - $\frac{1}{4}$ of a grain, produces great activity of the salivary and sweat glands, as well as increased secretion from the mucous surfaces and of urine from the kidneys. Used chiefly in dropsy and uræmia, to supplement and to assist the kidneys. It should be given with care in fatty heart and where there is valvular disease. The fluid extract of jaborandi is sometimes used, but it is not so exact a remedy as pilocarpine, which can be very well administered hypodermically.

Resorcin is an artificial hydrocarbon, used as an antiseptic and antipyretic in diphtheria, typhoid fever, and the fever of phthisis, and as an application to eczema. The dose is from 5-30 grains. It may be given in syrup of orange, well diluted. It produces profuse perspiration and its antipyretic action does not last long.

Santonin, the active principle of santonica, used as an anthelmintic, in doses of from 2-10 grains, depending upon the age of the patient.

Terebene.—An oily liquid prepared from turpentine, given in doses of 5-6 drops on a lump of sugar, or in an emulsion, every 2-4 hours, has proved very useful in winter cough (bronchitis). It also checks acidity and flatulence in the stomach.

Urethan.—A useful hypnotic. Dose, 15-60 grains at bedtime. Has been used successfully in doses of 4 grains every 2 hours in tetanus. It has been found by experiment to be antagonistic to strychnia.

Viburnum prunifolium Black Haw. The fluid extract of which the dose is 30-60 minims, is said to be valuable in the prevention of threatened abortion, and in menorrhagia. It is, however, by no means a certain remedy.

IMPORTANT INCOMPATIBLES.

Acacia (gum), with alcohol, iron, lead-water, and mineral acids.

Acids (mineral), with alkalis and relatively weak salts of other acids, such as bromides, chlorides, and iodides.

Alkalies, with acids and with relatively weak salts.

Arsenic, with tannic acid, salts and oxide of iron, and lime and magnesia.

Bitter infusions and tinctures, with salts of iron and lead.

Bromides, with acids, acid salts, or alkalis.

Calomel, with alkalis, lime-water, salts of iron and lead, and iodide of potassium.

Camphor (spirits of), with water.

Carbonates, with acids and acid salts.

Chlorides, with silver salts, lead salts, and alkalis.

Chloroform (except in minute quantity), with water.

Corrosive sublimate, with alkalis, lime-water, salts of iron and lead, iodide of potassium, albumen, gelatine, and vegetable astringents. It may, however, be advantageously combined with the tincture of the chloride of iron and the liq. acidi arseniosi of the new Pharmacopœia, or, in spite of the chemical reaction, with iodide of potassium, as in the famous Gibert's syrup.)

Digitalis, with iron and preparations containing tannic acid.

Iron (salts), with anything containing tannic acid. Tincture of the chloride of iron, with alkalis, carbonates, mucilages, and preparations containing tannic acid.

Mucilages, with acids, iron salts, and alcohol.

Potassium (iodide of), with all strong acids and acid salts. (See Corrosive Sublimate.)

Spirits of nitrous ether, with sulphate of iron, tincture of guaiacum, and most carbonates.

Tinctures of gums or resins, with water.

Vegetable preparations containing tannic acid, with salts of iron and lead.

ARTIFICIAL RESPIRATION Sylvester's Method).

FREE the body from any clothing that binds the neck, chest, or waist; turn it over upon the face for a moment, thrusting a finger into the mouth and sweeping it round, to bring away anything that may have gotten in or accumulated there. Then lay the body flat on the back, with something a few inches high under the shoulders, so as to cause the neck to

be stretched out and the chin to be carried from the chest. Draw the tongue well forward out of the mouth and let it be held by an assistant. (If there be no one to do this, a pencil or small stick may be thrust across the mouth on top of the tongue and back of the last teeth, to keep the mouth open and the tongue out of the throat. Place yourself on your knees behind the head, seize both arms near the elbows and sweep them round horizontally, away from the body and over the head till they meet above it; give a good, strong pull, and keep it up for a




FIG. 1.

few seconds. After this return the arms to their former position alongside the chest, and make strong pressure against the lower ribs, so as to drive the air out of the chest and effect an act of expiration. This need occupy but a second of time.

This plan, regularly carried out, will make about 16 complete acts of respiration in a minute. It should be kept up for a long time, and not aban-



FIG. 2.

doned until the heart has ceased to beat. It should be remembered that cessation of the pulse at the wrists amounts to nothing as a sign of death; and life is present when only a most acute ear can detect the sound of the heart. In a moderately thin person deep pressure with the finger-ends just below the lower end of the breastbone may sometimes reveal pulsation in the aorta when it cannot be found anywhere else.

POISONS AND ANTIDOTES.

THE following table contains suggestions for the proper treatment of every form of poisoning which is likely to occur. Though brief, they are all that will be needed to prompt the physician's memory:

| POISON. | TREATMENT. |
|---|--|
| <i>Nature unknown</i> | { Provoke repeated vomiting;
Give bland liquids;
Stimulate, if necessary. |
| <i>Acids</i> —
Sulphuric, {
Nitric, {
Muriatic, {
Oxalic, { | { Give an alkali;
Provoke vomiting;
Give bland fluids;
Secure rest;
Stimulate, if necessary. |
| <i>Prussic Acid</i> | { Give dilute hartshorn; chlorine-water: so-
lution of a salt of iron;
Produce vomiting and stimulate. |
| <i>Carbolic Acid</i> | { Give Epsom salts, dilute sulphuric acid,
and oil;
Produce vomiting and stimulate. |
| <i>Alkalies</i> —
Hartshorn, {
Soda, {
Potash, {
Lye, { | { Give an acid or a fixed oil;
Provoke vomiting;
Give bland liquids;
Secure rest;
Stimulate, if necessary. |
| <i>Arsenic</i> —
Paris green, {
Scheele's green, {
Fowler's solution, { | { Provoke vomiting;
Give hydrated oxide of iron or dialyzed
iron and salt;
Give dose of castor oil;
Secure rest;
Stimulate, if necessary. |
| <i>Acetate of Lead</i> | { Give Epsom salts or dilute sulphuric acid;
Provoke vomiting;
Give bland liquids;
Give dose of castor oil. |
| <i>Mercury</i> , {
<i>Corrosive Sublimate</i> , {
<i>Antimony</i> , {
<i>Tartar Emetic</i> , { | { Provoke vomiting;
Give some infusion containing tannic acid;
Give raw eggs and milk;
Give dose of castor oil;
Stimulate, if necessary. |
| <i>Copper Salts</i> | { Give albumen;
Provoke vomiting;
Give bland fluids. |
| <i>Phosphorus</i> | { Provoke vomiting;
Give five-grain doses sulphate of copper;
Give dose of magnesia, but <i>no</i> oil. |
| <i>Nitrate of Silver</i> {
(lunar caustic), { | { Give strong salt and water; } repeat many
Provoke vomiting; } times. |
| <i>Iodine</i> | { Provoke vomiting;
Give starch and water;
Give bland fluids. |

| POISON. | TREATMENT. |
|--|--|
| <i>Opium</i> —
Morphine,
Laudanum,
Paregoric, etc.,
Chloral, | { Provoke vomiting, repeatedly;
Give atropia, and strong coffee or tea;
Keep up the breathing, preferably with
battery. |
| <i>Strychnine</i> | { Provoke vomiting, once or twice;
Give chloral and bromide of potash;
Secure absolute quiet. |
| <i>Acute</i> | { Provoke vomiting;
Stimulate well. |
| <i>Jamestown Weed</i> ,
<i>Hemlock</i> ,
<i>Nightshade</i> (belladonna),
<i>Toadstool</i> ,
<i>Tobacco, etc., etc.</i> , | { { Provoke vomiting and give a purge;
Stimulate well. |
| <i>Alcohol</i> | { Provoke vomiting;
Give hartshorn and water. |
| <i>Decayed Meat or Vegetables</i> | { Provoke vomiting;
Give a purgative;
Give powdered charcoal. |
| <i>Poisonous Gases</i> —
Carbonic acid or oxide,
Sulphuretted hydrogen, | { { Fresh air;
Artificial respiration;
Stimulation. |

To provoke vomiting, warm water may be used with or without ground mustard (a tablespoonful to a pint of water), or ipecac (a teaspoonful of the powder or a tablespoonful or so of the syrup), or a finger may be thrust down the throat. It is best to give large quantities (that is, a pint at a time) of warm water whenever vomiting is to be excited.

Bland liquids are milk, raw eggs, some sort of oil, gruel, etc.

Stimulants are tea, coffee, whiskey, wine, etc., or hartshorn and water. Of this a teaspoonful in a teacupful of water will be enough for a dose. In making tea or coffee one must not wait to do it as if for the table, but mix hot water and the leaves or grounds, squeeze them well, stir together, and give the whole—leaves, grounds, everything.

Alkaline antidotes which are most likely to be at hand are hartshorn and water a tablespoonful in two teacupfuls of water, soap and water, lime, whiting, soda, chalk, tooth-powder, plaster, magnesia, white-wash, and even wood-ashes.

Acid antidotes most commonly accessible are vinegar and lemon-juice.

In giving an antidote it is well to remember that it is not always necessary to wait for it to dissolve, but that it may be stirred up in any fluid at hand (except oil), and swallowed immediately.

TABLE OF DOSES OF REMEDIES MOST FREQUENTLY ADMINISTERED.

For hypodermic use the dose should be half that used by the mouth.

For use by rectum the dose should be twice that by the mouth.

Doses for Children.—Dr. Young's rule: Add 12 to the age, and divide by the age to give the denominator of a fraction; the numerator of which is 1. Example: For a child two years old, $\frac{12+2}{2} = \frac{1}{4}$ th; the dose should be $\frac{1}{4}$ th that for an adult. In giving powerful medicines and opium still smaller doses must be used for children.

| REMEDY. | DOSE. | REMEDY. | DOSE. |
|---------------------------------|---------------------------------------|--------------------------------|--------------------------------------|
| Abstract. belladonnæ..... | $\frac{1}{2}$ grain. | Ant. et pot. tartr; diaph..... | $\frac{1}{10}$ to $\frac{1}{2}$ gr. |
| cannab. ind..... | 1 grain. | emetic..... | 1 to 2 grs. |
| coni..... | 1 to 3 grains. | Antimonii oxysulphuret..... | $\frac{1}{2}$ to 2 grs. |
| gelsemii..... | 1 grain. | Antipyrin..... | $\frac{1}{5}$ to 30 grs. |
| hyoscyami..... | 2 grains. | Apomorph. hydrochlor..... | $\frac{1}{30}$ to $\frac{1}{2}$ gr |
| nuc. vom..... | 1 grain. | Aqua ammoniæ..... | 5 to 30 mins. |
| podophylli..... | 1 grain. | amygd. amar..... | 2 fl. drachms. |
| Acet. opii..... | 5 minims. | camphoræ..... | 1 to 4 fl. drs. |
| scillæ..... | 10 to 30 mins. | chlori..... | 1 to 4 fl. drs. |
| Acid. acet. dil..... | 60 minims. | creasoti..... | 1 to 4 fl. drs. |
| arsenici..... | $\frac{1}{2}$ to $\frac{1}{4}$ grain. | laurocerasi..... | 6 to 30 mins. |
| benzoic..... | 5 to 15 grs. | Argenti nitras..... | $\frac{1}{4}$ to $\frac{1}{2}$ gr. |
| boric..... | 5 to 10 grs. | Arsenii iodidum..... | $\frac{1}{12}$ to $\frac{1}{10}$ gr. |
| carbolic..... | 1 to 3 grs | Asafetida..... | 5 to 20 grs. |
| gallic..... | 3 to 15 grs. | Atropina..... | $\frac{1}{12}$ to $\frac{1}{6}$ gr. |
| hydrobrom. 34 grs..... | 10 to 15 grs. | Atropinæ sulph..... | $\frac{1}{12}$ to $\frac{1}{6}$ gr. |
| hydrobrom. dil..... | 15 to 60 mins. | Auri et sodii chlorid..... | $\frac{1}{32}$ to $\frac{1}{16}$ gr. |
| hydrochlor..... | 3 to 10 mins. | Bismuthi citras..... | 3 to 15 grs. |
| hydrochlor. dil..... | 10 to 30 mins. | et ammon. citr..... | 1 to 15 grs. |
| hydrocyan. dil..... | 2 to 6 mins. | subnitr..... | 3 to 15 grs. |
| lactic..... | 15 to 60 grs. | valer..... | 1 to 3 grs. |
| nitric..... | 3 to 10 mins. | Calcina..... | 1 to 5 grs. |
| nitr. dil..... | 10 to 30 mins. | Calcii hypophosphis..... | 3 to 15 grs. |
| nitro-hydrochlor..... | 3 to 10 mins. | iodidum..... | 1 to 3 grs. |
| nitro-hydrochlor. dil..... | 5 to 20 mins. | phosphas..... | 15 to 30 grs. |
| phosphoric (50 grs..... | 3 to 15 grs. | Calx sulphurata..... | $\frac{1}{2}$ to 1 gr. |
| phosphoric dil..... | 10 to 20 mins. | Cambogium..... | 1 to 4 grs. |
| salicylic..... | 5 to 15 grs. | Camphora..... | 3 to 10 grs. |
| sulphuric..... | 5 to 10 mins. | monobrom..... | 2 to 5 grs. |
| sulphuric. dil..... | 10 to 20 mins. | Cantharis..... | $\frac{1}{4}$ to 2 grs. |
| sulphuric. arom..... | 5 to 30 mins. | Capsicum..... | 1 to 3 grs. |
| sulphuros..... | 30 to 60 mins. | Castoreum..... | 6 to 15 grs. |
| tannic..... | 2 to 15 grs. | Catechu..... | 15 to 30 grs. |
| Aconitina (white crystals)..... | $\frac{1}{60}$ to $\frac{1}{30}$ gr. | Ceri nitras..... | 1 to 3 grs. |
| Aloe..... | 2 to 5 grs. | oxalas..... | 1 to 3 grs. |
| Aloinum..... | 1 to 3 grs. | Chinoidinum..... | 3 to 30 grs. |
| Alumen..... | 5 to 15 grs. | Chloral..... | 10 to 30 grs. |
| Ammonii benzoas..... | 10 to 20 grs. | Chloroformum..... | 1 to 5 mins |
| bronid..... | 5 to 30 grs. | Chrysarobinum..... | 3 to 15 grs. |
| carb..... | 3 to 10 grs. | Cinchona..... | 15 to 60 grs. |
| chlorid..... | 5 to 30 grs. | Cinchonidina..... | 1 to 30 grs. |
| iodid..... | 3 to 15 grs. | Cinchonina..... | 1 to 30 grs. |
| phosph..... | 5 to 20 grs. | Cinnamomum..... | 6 to 30 grs. |
| valer..... | 3 to 15 grs. | Cocaine hydrochlorate..... | 4 to 8 solut. dr. |
| Amyl nitris..... | 2 to 5 mins. | Codeina..... | $\frac{1}{4}$ to 2 grains. |

| REMEDY. | DOSE. |
|--------------------------------------|------------------------------------|
| Confectio sennæ..... | 1 to 2 grs. |
| Coniina and its salts..... | $\frac{1}{4}$ to $\frac{1}{2}$ gr. |
| Copaiba..... | 15 to 60 mins. |
| Cotula..... | 1 to 1 gr. |
| Creatum..... | 1 to 3 mins. |
| Cræta præpar..... | 15 to 75 grs. |
| Croton chloral..... | 1 to 10 grs. |
| Cubela..... | 15 to 60 grs. |
| Cupri acetat..... | $\frac{1}{2}$ to 6 grs. |
| sulphas..... | $\frac{1}{2}$ to 10 grs. |
| Cuprum ammon..... | $\frac{1}{2}$ to 1 gr. |
| Curare..... | $\frac{1}{2}$ to $\frac{1}{4}$ gr. |
| Decoct. aloes comp..... | 1 to 2 fl. ozs. |
| sarsap comp..... | 2 to 6 fl. ozs. |
| Digitalinum..... | $\frac{1}{2}$ to $\frac{1}{4}$ gr. |
| Digitalis..... | 1 to 3 grs. |
| Duboisina, and its salts..... | 15 to 30 gr. |
| Elaterinum; U.S.P., 1880..... | $\frac{1}{2}$ to $\frac{1}{4}$ gr. |
| Elaterium; U.S.P., 1870..... | $\frac{1}{2}$ to 4 gr. |
| Ergota..... | 15 to 60 grs. |
| Ergotinum..... | 2 to 8 grs. |
| Eserina and its salts..... | $\frac{1}{2}$ to $\frac{1}{4}$ gr. |
| Euonymin..... | 2 to 4 grs. |
| Extr. acori. rad.; U.S.P., 1880..... | $\frac{1}{2}$ to $\frac{1}{4}$ gr. |
| aconiti (rad.) fluid..... | 1 to 5 mins. |
| aloes aquos..... | 1 to 3 grs. |
| angusture fl..... | 15 to 45 mins. |
| angelicæ rad. fl..... | 30 to 60 mins. |
| anemidis..... | 2 to 10 grs. |
| anemidis fl..... | 30 to 60 mins. |
| apocyni cannab. fl..... | 8 to 30 mins. |
| arece fl..... | 15 to 75 mins. |
| arnicæ flor..... | 3 to 8 grs. |
| arnicæ fl..... | 5 to 15 mins. |
| arnicæ rad..... | 2 to 5 grs. |
| arnicæ rad. fl..... | 5 to 15 mins. |
| aromat. fl..... | 30 to 60 mins. |
| aspidospermæ fl..... | 15 to 45 mins. |
| aurantii cort. fl..... | $\frac{1}{2}$ to 2 fl. drs. |
| azedarach fl..... | 15 to 75 mins. |
| baptisæ fl..... | 7 to 30 mins. |
| bellad. fol. fl..... | 3 to 6 mins. |
| bellad. rad..... | $\frac{1}{2}$ to 1 gr. |
| bellad. rad. fl..... | 1 to 3 mins. |
| brayeræ fl..... | 2 to 4 fl. drs. |
| bryoniæ fl..... | 15 to 60 mins. |
| buchu fl..... | $\frac{1}{2}$ to 2 fl. drs. |
| calami fl..... | 15 to 60 mins. |
| calend..... | 15 to 60 mins. |
| calumbæ..... | 3 to 10 grs. |
| calumbæ fl..... | 15 to 60 mins. |
| canelle fl..... | 15 to 60 mins. |
| cannab. Amer. fl..... | 3 to 15 mins. |
| cannab. ind..... | $\frac{1}{2}$ to 1 gr. |
| cannab. ind. fl..... | 3 to 6 mins. |
| cantharidis fl..... | 1 to 3 mins. |
| capsici fl..... | 1 to 3 mins. |
| cardam. comp. fl..... | 15 to 45 mins. |
| cascarilla fl..... | $\frac{1}{2}$ to 2 fl. drs. |
| caulophylli fl..... | 15 to 30 mins. |
| chinaph. fl..... | $\frac{1}{2}$ to 1 fl. drs. |
| cimicifuge fl..... | 8 to 30 mins. |
| cinchona..... | 15 to 30 grs. |
| cinchona fl..... | 30 to 60 mins. |

| REMEDY. | DOSE. |
|------------------------------|---|
| Extr. cinchona arom. fl..... | 30 to 60 mins. |
| cinchonæ comp. fl..... | $\frac{1}{2}$ to $\frac{1}{2}$ fl. drs. |
| cocculi fl..... | 1 to 3 mins. |
| coleb. rad..... | $\frac{1}{2}$ to 1 fl. grs. |
| coleb. rad. fl..... | 3 to 15 mins. |
| coleb. sem. fl..... | $\frac{1}{2}$ to 10 mins. |
| colocynth..... | 1 to 5 grs. |
| colocynth. comp..... | 1 to 5 grs. |
| coni fol. fl..... | 3 to 15 mins. |
| convallariæ rad. fl..... | 15 to 30 mins. |
| corn. flor. fl..... | 30 to 60 mins. |
| coto fl..... | 1 to 15 mins. |
| cubebæ fl..... | 15 to 30 mins. |
| cypripedii fl..... | 15 to 60 mins. |
| damiana fl..... | $\frac{1}{2}$ to 2 fl. drs. |
| digitalis..... | $\frac{1}{2}$ to $\frac{1}{4}$ gr. |
| digitalis fl..... | 1 to 6 mins. |
| droseræ fl..... | 5 to 10 mins. |
| dulcamaræ..... | 5 to 15 grs. |
| dulcamaræ fl..... | 1 to 2 fl. drs. |
| eigete..... | 1 to 8 grs. |
| ergotæ fl..... | 15 to 60 mins. |
| erythroxyl fl..... | $\frac{1}{2}$ to 2 fl. drs. |
| eucalypti fl..... | 15 to 60 mins. |
| euonymi fl..... | 15 to 60 mins. |
| eupatorii fl..... | 30 to 60 mins. |
| euphorb. ipec. fl..... | 5 to 30 mins. |
| gallæ fl..... | $\frac{1}{2}$ to 2 fl. drs. |
| gelsem..... | 2 to 8 mins. |
| gelsem fl..... | 1 to 8 mins. |
| gent fl..... | 30 to 60 mins. |
| gent. comp. fl..... | 30 to 60 mins. |
| gerani fl..... | 15 to 30 mins. |
| gilleme fl..... | 15 to 30 mins. |
| gossypii fl..... | 15 to 45 mins. |
| granati rad. cort. fl..... | $\frac{1}{2}$ to 2 fl. drs. |
| grind. rob. fl..... | 30 to 60 mins. |
| grind. squarr. fl..... | 30 to 60 mins. |
| guaiaci ligni fl..... | 30 to 60 mins. |
| guaranæ fl..... | 15 to 30 mins. |
| hæmatoxyl..... | 8 to 30 grs. |
| hæmatoxyl fl..... | 30 to 60 mins. |
| hamamelidis fl..... | 60 to 90 mins. |
| helleb. nigris..... | $\frac{1}{2}$ to 3 grs. |
| helleb. nigris. fl..... | 5 to 15 mins. |
| heloniæ fl..... | 8 to 30 mins. |
| hepaticæ fl..... | 30 to 60 mins. |
| humuli..... | 3 to 15 grs. |
| humuli fl..... | 30 to 60 mins. |
| hydrangeæ fl..... | 30 to 60 mins. |
| hydrastis..... | 3 to 10 grs. |
| hydrastis fl..... | 8 to 30 mins. |
| hyoscyami fol. fl..... | 3 to 30 mins. |
| hyoscyami sem. fl..... | 2 to 8 mins. |
| ignatie..... | $\frac{1}{2}$ to $\frac{1}{2}$ grs. |
| ignatie fl..... | 1 to 6 mins. |
| ipæac. fl..... | 3 to 60 mins. |
| iris versicol..... | 3 to 6 grs. |
| irid. versicol. fl..... | 15 to 30 mins. |
| jalapæ; U.S.P., 1870..... | 5 to 10 grs. |
| japagæ fl..... | 15 to 60 mins. |
| juglandis..... | 15 to 30 grs. |
| juglandis fl..... | $\frac{1}{2}$ to 2 fl. drs. |
| junip. fl..... | 30 to 60 mins. |

| REMEDY. | DOSE. | REMEDY. | DOSE. |
|--|-----------------------------|------------------------------------|------------------------------------|
| Ext. kino fl. | 15 to 30 mins. | Extr. spigeliae et sennae fl. | $\frac{1}{2}$ to 2 fl. drs. |
| krameria 5 to 15 grs. | | stillingia fl. | $\frac{1}{2}$ to 2 fl. drs. |
| krameria fl. | 30 to 60 mins. | stillingia comp. fl. | $\frac{1}{2}$ to 2 fl. drs. |
| laetia 5 to 15 grs. | | stramonii fol. alc. | $\frac{1}{2}$ to $\frac{1}{2}$ gr. |
| laetia fl. | 15 to 60 mins. | stramonii sem. | $\frac{1}{2}$ to $\frac{1}{2}$ gr. |
| laetia fl. | 8 to 30 mins. | stramonii fl. | 1 to 6 mins. |
| leptandra 3 to 10 grs. | | sumbul fl. | 15 to 60 mins. |
| leptandra fl. | 30 to 60 mins. | taraxac. | 5 to 10 grs. |
| lobelia fl. | 5 to 30 c.c. | taraxac. fl. | $\frac{1}{2}$ to 2 fl. drs. |
| lobelia sem. fl. | 3 to 15 c.c. | trit. rep. fl. | 1 to 4 fl. drs. |
| lupulini fl. | 10 to 30 mins. | urticae rad. fl. | 5 to 15 mins. |
| lycopi fl. | 5 to 30 mins. | ustilag. maid. fl. | 15 to 60 mins. |
| multi. | 1 to 2 $\frac{1}{2}$ drs. | uva ursi fl. | 30 to 60 mins. |
| matico fl. | 30 to 60 mins. | valerian. | 5 to 15 grs. |
| mezerri $\frac{1}{2}$ to 1 gr. | | valer. fl. | 30 to 60 mins. |
| mezerri fl. | 3 to 10 mins. | veratr. vir. fl. | 2 to 8 mins. |
| nectandra fl. | 1 to 4 fl. drs. | viburni [prunifol.] fl. | 1 to 2 fl. drs. |
| nuc. vom. | $\frac{1}{2}$ to 1 gr. | xanthoxyli cort. fl. | 15 to 30 mins. |
| nuc. vom. fl. | 1 to 5 mins. | xanthoxyli fruct. fl. | 15 to 30 mins. |
| oppi. | 1 to 1 gr. | zingiberis fl. | 8 to 30 mins. |
| papaveris. | $\frac{1}{2}$ to 2 grs. | Fel bovis purif. | 3 to 6 grs. |
| papaveris fl. | 15 to 45 mins. | Ferri arsen. | $\frac{1}{2}$ to $\frac{1}{2}$ gr. |
| pareire fl. | 30 to 60 mins. | benzoas. | 1 to 5 grs. |
| petroselin. fl. | 1 to 2 fl. drs. | bronid. | 1 to 5 grs. |
| physostigma $\frac{1}{2}$ to $\frac{1}{2}$ gr. | | carb. sacch. | 4 to 15 grs. |
| physostigma fl. | 1 to 5 mins. | chlorid. | 1 to 5 grs. |
| phytolacca baccar. fl. | 5 to 30 mins. | citr. | 5 to 10 grs. |
| phytolacca rad. | 1 to 3 grs. | et ammon. citr. | 5 to 10 grs. |
| phytolacca rad. fl. | 5 to 30 mins. | et ammon. sulph. | 5 to 10 grs. |
| pilocarpi fl. | 15 to 60 mins. | et ammon. tartr. | 5 to 10 grs. |
| piper. nigr. fl. | 15 to 45 mins. | et cinchonid. citr. | 5 to 10 grs. |
| piscidia fl. | 15 to 60 mins. | et pot. tartr. | 5 to 30 grs. |
| podophylli $\frac{1}{2}$ to 1 gr. | | et quin. citr. | 5 to 10 grs. |
| podophylli fl. | 8 to 30 mins. | et strychn. citr. | 1 to 5 grs. |
| prun. virg. fl. | 30 to 60 mins. | hypophosphis. | 5 to 10 grs. |
| pulsatilla fl. | 2 to 10 mins. | iodidum. | 1 to 5 grs. |
| quassia. | 1 to 5 grs. | iodidum sacch. | 2 to 10 grs. |
| quassia fl. | 30 to 60 mins. | lactas. | 1 to 3 grs. |
| quercus fl. | 30 to 60 mins. | oxalas. | 1 to 3 grs. |
| rhei. | 5 to 15 grs. | oxid. hydrat. | $\frac{1}{2}$ to 2 ozs. |
| rhei fl. | 15 to 45 mins. | phosphas. | 1 to 5 grs. |
| rhois arom. fl. | 15 to 60 mins. | pyrophosphas. | 1 to 5 grs. |
| rhois glabr. cort. fl. | 30 to 60 mins. | subcarb. | 5 to 30 grs. |
| rhois glabr. fruct. fl. | 30 to 60 mins. | sulphas. | 1 to 3 grs. |
| rhois toxicod. fl. | 1 to 6 mins. | sulphas exsicc. | 1 to 3 grs. |
| ricini fol. fl. | $\frac{1}{2}$ to 2 fl. drs. | valer. | 1 to 3 grs. |
| rosa fl. | $\frac{1}{2}$ to 2 fl. drs. | Ferrum dialysatum. | 5 to 15 mins. |
| rubi fl. | 15 to 50 mins. | redactum. | 1 to 5 grs. |
| sabinae fl. | 5 to 15 mins. | Guarana 10 to 30 grs. | |
| salicis fl. | $\frac{1}{2}$ to 2 fl. drs. | Hydrarg. chlorid. corros. | $\frac{1}{2}$ to $\frac{1}{2}$ gr. |
| salvia fl. | $\frac{1}{2}$ to 2 fl. drs. | chlorid. mite. | $\frac{1}{2}$ to 10 grs. |
| sanguin. fl. | 5 to 15 mins. | cyanid. | $\frac{1}{2}$ to $\frac{1}{2}$ gr. |
| santali citr. fl. | 1 to 2 fl. drs. | iodid. flav. | $\frac{1}{2}$ to 1 gr. |
| santonica fl. | 15 to 60 mins. | iodid. rubr. | $\frac{1}{2}$ to $\frac{1}{2}$ gr. |
| sarsap. fl. | $\frac{1}{2}$ to 2 fl. drs. | iodid. vir. | $\frac{1}{2}$ to 1 gr. |
| sassafras fl. | 1 to 2 fl. drs. | oxid. flav. | $\frac{1}{2}$ to $\frac{1}{2}$ gr. |
| scilla fl. | 5 to 30 mins. | oxid. nigr. | $\frac{1}{2}$ to 1 gr. |
| scilla comp. fl. | 5 to 30 mins. | oxid. rubr. | $\frac{1}{2}$ to $\frac{1}{2}$ gr. |
| scoparii fl. | $\frac{1}{2}$ to 1 fl. dr. | subsulphas flav. | $\frac{1}{2}$ to 1 gr. |
| senega fl. | 8 to 15 mins. | c. creta. | 1 to 8 grs. |
| sennae fl. | 1 to 4 fl. drs. | Hyosciamina, and salts. | $\frac{1}{2}$ to $\frac{1}{2}$ gr. |
| serpent. fl. | 30 to 60 mins. | Infusum digitalis. | 2 to 4 fl. drs. |
| simarubae fl. | 15 to 30 mins. | senna comp. | 1 to 2 fl. ozs. |
| spigelia fl. | 15 to 60 mins. | Iodoformum. | 1 to 3 grs. |

| REMEDY. | DOSE. | REMEDY. | DOSE. |
|-----------------------------|-----------------|------------------------------|---------------|
| Iodum..... | 1 to 1 gr. | Pepsinum purum..... | 15 grains. |
| Ipecacuanha) expect..... | 1 to 1 gr. | saccharatum..... | 30 grains. |
|) emet..... | 15 to 30 grs. | Phosphorus..... | 1 to 1 gr. |
| Iridium..... | 2 to 1 grs. | Physostigmine salicylus..... | 1 to 1 gr. |
| Jalap..... | 15 to 30 grs. | sulphas..... | 1 to 1 gr. |
| Kanaba..... | 1 to 2 drs. | Picrotoxinum..... | 1 to 1 gr. |
| Kino..... | 8 to 30 grs. | Pilocarpina, and salts..... | 1 to 1 gr. |
| Lactucarium..... | 8 to 15 grs. | Pil. aloes..... | 1 to 3 pills. |
| Liq. ammon. acet..... | 2 to 1 fl. drs. | aloes et asafet..... | 2 to 5 pills. |
| acidi arseniosi..... | 2 to 7 mins. | aloes et ferri..... | 1 to 3 pills. |
| arsen. et hydr. iod..... | 2 to 7 mins. | aloes et mast..... | 1 to 3 pills. |
| ferri chloridi..... | 2 to 10 mins. | aloes et myrrhæ..... | 2 to 5 pills. |
| ferri dialys..... | 10 to 15 mins. | antim. comp..... | 1 to 3 pills. |
| ferri nitrat..... | 8 to 15 mins. | asa-fetida..... | 1 to 6 pills. |
| pepsini..... | 1 to 4 fl. drs. | cathart. comp..... | 1 to 4 pills. |
| potassæ..... | 5 to 30 mins. | ferri comp..... | 2 to 5 pills. |
| potassii arseni..... | 3 to 7 mins. | ferri iodidi..... | 1 to 4 pills. |
| potassii citrat..... | 2 to 1 fl. drs. | opii..... | 1 to 2 pills. |
| sodæ..... | 5 to 30 mins. | phosphori..... | 1 to 4 pills. |
| sodii arseniatis..... | 3 to 7 mins. | rhei..... | 2 to 5 pills. |
| Lithii benzoas..... | 2 to 5 grs. | rhei comp..... | 2 to 5 pills. |
| bromid..... | 1 to 3 grs. | Piperinum..... | 1 to 8 grs. |
| carb..... | 2 to 6 grs. | Plumbi acetæ..... | 1 to 3 grs. |
| citr..... | 2 to 5 grs. | noidum..... | 1 to 4 grs. |
| salicylus..... | 2 to 8 grs. | Potassii acetæ..... | 15 to 60 grs. |
| Lupulinum..... | 5 to 10 grs. | bicarb..... | 8 to 60 grs. |
| Magnesiæ..... | 15 to 60 grs. | bitart..... | 1 to 2 drs. |
| Magnesiæ carb..... | 1 to 60 grs. | bromid..... | 8 to 60 grs. |
| citr. gran..... | 2 to 1 drs. | carb..... | 8 to 30 grs. |
| sulphas..... | 2 to 8 drs. | chloras..... | 5 to 30 grs. |
| sulphis..... | 8 to 30 grs. | citræ..... | 15 to 60 grs. |
| Mangani sulphas..... | 2 to 10 grs. | cyanid..... | 1 to 1 gr. |
| Manna..... | 1 to 2 ozs. | et sodii tart..... | 1 to 1 oz. |
| Massa copaiab..... | 5 to 30 grs. | hypophosphis..... | 5 to 15 grs. |
| ferri carb..... | 5 to 1 grs. | iodid..... | 5 to 60 grs. |
| hydrarg..... | 1 to 10 grs. | nitras..... | 8 to 15 grs. |
| Mist. ammoniaci..... | 1 to 8 fl. drs. | sulphidum..... | 1 to 10 grs. |
| asa-fetida..... | 2 to 4 fl. drs. | sulphis..... | 1 to 30 grs. |
| chloroformi..... | 2 to 4 fl. drs. | tartas..... | 1 to 8 drs. |
| cretæ..... | 1 to 4 fl. drs. | Pulv. antimonialis..... | 3 to 10 grs. |
| ferri comp..... | 1 to 2 fl. ozs. | aromat..... | 5 to 30 grs. |
| ferri et amm. acet..... | 1 to 4 fl. drs. | cretæ comp..... | 8 to 30 grs. |
| glycyrrh. comp..... | 1 to 4 fl. drs. | glycyrrh. comp..... | 30 to 60 grs. |
| magnes. et asafet..... | 1 to 4 fl. drs. | ipecac. et opii..... | 5 to 15 grs. |
| potassii citr..... | 1 to 1 fl. oz. | jalapæ comp..... | 30 to 60 grs. |
| rhei et sodæ..... | 1 to 1 fl. oz. | morphinæ comp..... | 8 to 15 grs. |
| Morphina and its salts..... | 1 to 1 gr. | rhei comp..... | 30 to 60 grs. |
| Narcaina..... | 1 to 2 grs. | Quinidina (and salts)..... | 1 to 30 grs. |
| Nitro-glycerinum..... | 1 to 1 gr. | Quinina (and salts)..... | 1 to 30 grs. |
| Nux vomica..... | 1 to 5 grs. | Quinine arsenias..... | 1 to 1 gr. |
| Oleoresina capsici..... | 1 to 1 gr. | Resina copaiabæ..... | 2 to 10 grs. |
| cubebæ..... | 5 to 30 grs. | jalapæ..... | 2 to 5 grs. |
| lupulini..... | 2 to 5 grs. | podophylli..... | 1 to 1 gr. |
| piperis..... | 1 to 3 grs. | scammonii..... | 2 to 10 grs. |
| zingiberis..... | 1 to 3 grs. | Rheum..... | 2 to 30 grs. |
| Oleum copaiabæ..... | 8 to 15 mins. | Salicinum..... | 8 to 30 grs. |
| cubebæ..... | 15 to 30 mins. | Santonica..... | 8 to 60 grs. |
| eucalypti..... | 5 to 15 mins. | Santonium..... | 1 to 5 grs. |
| eucalypti..... | 10 to 30 mins. | Scammonium..... | 3 to 15 grs. |
| phosphoratum..... | 1 to 3 mins. | Senna..... | 8 to 60 grs. |
| sabina..... | 1 to 3 mins. | Sodii acetæ..... | 15 to 60 grs. |
| terebinth..... | 5 to 60 mins. | arsenias..... | 1 to 1 gr. |
| tiglii..... | 1 to 4 drops. | benzoas..... | 5 to 15 grs. |
| Opium (14 morphinæ)..... | 1 to 1½ grs. | bicarb..... | 8 to 30 grs. |

| REMEDY. | DOSE. | REMEDY. | DOSE. |
|--------------------------------|------------------------------------|--|------------------------------------|
| Sodii bisulphis..... | 8 to 30 grs. | Tinct. conii..... | 5 to 30 mins. |
| boras..... | 8 to 30 grs. | cubebæ..... | 1 to 2 fl. drs. |
| bromid..... | 8 to 30 grs. | digitalis..... | 5 to 15 mins. |
| carb..... | 8 to 30 grs. | ferri acet..... | 15 to 30 mins. |
| hypophosphis..... | 8 to 15 grs. | ferri chloridi..... | 15 to 60 mins. |
| hyposulphis..... | 8 to 30 grs. | ferri chloridi ather..... | 15 to 30 mins. |
| iodidum..... | 5 to 15 grs. | gallæ..... | $\frac{1}{2}$ to 2 fl. drs. |
| phosphas..... | 2 to 15 grs. | gelsemii..... | 5 to 10 mins. |
| salicylas..... | 5 to 30 grs. | guaiac..... | 30 to 60 mins. |
| santoninas..... | 2 to 10 grs. | guaiaci ammon..... | 30 to 60 mins. |
| sulphas..... | 1 to 2 drs. | hellebori..... | 10 to 15 mins. |
| sulphis..... | 8 to 30 grs. | humuli..... | 1 to 4 fl. drs. |
| Spir. æther..... | 30 to 60 mins. | hydrastis..... | 30 to 90 mins. |
| æther. nitrosi..... | $\frac{1}{2}$ to 2 fl. drs. | hyoscyami fol..... | 15 to 60 mins. |
| ammonia..... | 8 to 30 mins. | hyoscyami sem..... | 15 to 30 mins. |
| ammonia arom..... | 15 to 60 mins. | ignatia..... | 5 to 15 mins. |
| camphoræ..... | 8 to 30 mins. | iodi..... | 5 to 15 mins. |
| chloroformi..... | 15 to 60 mins. | ipeac. et opii..... | 5 to 15 mins. |
| lavend. comp..... | 30 to 60 mins. | jalapæ..... | $\frac{1}{2}$ to 2 fl. drs. |
| menth. pip..... | 30 to 60 mins. | kinæ..... | $\frac{1}{2}$ to 2 fl. drs. |
| Strychnina (and salts)..... | $\frac{1}{4}$ to $\frac{1}{2}$ gr. | krameria..... | $\frac{1}{2}$ to 2 fl. drs. |
| Sulphur..... | $\frac{1}{2}$ to 1 drs. | lavend. comp..... | $\frac{1}{2}$ to 2 fl. drs. |
| Syrupus acidi hydroiodici..... | 1 to 4 fl. drs. | lobelia..... | 15 to 45 mins. |
| alii..... | 1 to 4 fl. drs. | lupulini..... | $\frac{1}{2}$ to 2 fl. drs. |
| calci lactophos..... | 1 to 2 fl. drs. | maticæ..... | $\frac{1}{2}$ to 2 fl. drs. |
| calcis..... | 15 to 30 mins. | nuc. vomiceæ..... | 5 to 30 mins. |
| ferri bromidi..... | 15 to 60 mins. | opii..... | 5 to 25 mins. |
| ferri iodidi..... | 15 to 60 mins. | opii camph..... | 1 to 2 fl. drs. |
| ferri oxidi..... | 1 fl. drachm. | physostigmatis..... | 5 to 15 mins. |
| ferri hypophosphi..... | 1 fl. drachm. | quassia..... | $\frac{1}{2}$ to 2 fl. drs. |
| fer. quin et str. phos..... | 1 fl. drachm. | rhei..... | 1 to 8 fl. drs. |
| hypophosphit..... | 1 fl. drachm. | rhei arom..... | 30 to 75 mins. |
| ipeacæ..... | $\frac{1}{2}$ to 4 fl. drs. | rhei dule..... | 1 to 4 fl. drs. |
| krameria..... | $\frac{1}{2}$ to 4 fl. drs. | sanguinariae..... | 15 to 60 mins. |
| lactucarii..... | 1 to 3 fl. drs. | scilla..... | 8 to 60 mins. |
| rhei..... | 1 to 4 fl. drs. | serpentaria..... | $\frac{1}{2}$ to 2 fl. drs. |
| rhei arom..... | 1 to 4 fl. drs. | stramon. fol..... | 8 to 15 mins. |
| roseæ..... | 1 to 2 fl. drs. | stramon sem..... | 6 to 15 mins. |
| rubi..... | 1 to 2 fl. drs. | sumbul..... | 8 to 30 mins. |
| sarsap. comp..... | 1 to 4 fl. drs. | valer..... | $\frac{1}{2}$ to 2 fl. drs. |
| scilla..... | $\frac{1}{2}$ to 1 fl. dr. | valer. ammon..... | $\frac{1}{2}$ to 2 fl. drs. |
| scilla comp..... | 15 to 60 mins. | veratr. vir..... | 1 to 10 mins. |
| senega..... | 1 to 2 fl. drs. | zingiberis..... | 15 to 60 mins. |
| senna..... | 1 to 4 fl. drs. | Tritur. elaterini..... | $\frac{1}{2}$ to $\frac{1}{4}$ gr. |
| Tinct. aceniti fol..... | 5 to 16 mins. | Veratrina..... | 1 to $\frac{1}{2}$ gr. |
| aconiti rad..... | 1 to 5 mins. | Vin. aloes..... | 1 to 2 fl. drs. |
| acon. rad. Fleming's..... | $\frac{1}{2}$ to 2 fl. drs. | antim. $\left\{ \begin{array}{l} \text{expect. \& alt.} \\ \text{emet.} \end{array} \right.$ | 1 to 8 mins. |
| aloes (1880)..... | $\frac{1}{2}$ to 2 fl. drs. | colch. rad..... | 30 to 75 mins. |
| aloes et myrrha..... | 1 to 2 fl. drs. | colch. sem..... | 8 to 45 mins. |
| arnica flor..... | 8 to 30 mins. | ergotæ..... | 5 to 30 mins. |
| arnica rad..... | 15 to 30 mins. | ferri amar..... | 1 to 4 fl. drs. |
| asafoetida..... | 30 to 60 mins. | ferri amar..... | 1 fl. drachm. |
| belladonna..... | 5 to 15 mins. | ferri citrat..... | 1 fl. drachm. |
| bryonia..... | 15 to 30 mins. | ipeacæ $\left\{ \begin{array}{l} \text{expect.} \\ \text{emet.} \end{array} \right.$ | 5 to 15 mins. |
| calendula..... | 15 to 30 mins. | opii..... | 2 to 4 fl. drs. |
| calumbæ..... | 1 to 4 fl. drs. | opii..... | 5 to 10 mins. |
| cannabis ind..... | 15 to 30 mins. | rhei..... | 1 to 2 fl. drs. |
| cantharid..... | 5 to 15 mins. | Zinci acet..... | 1 to 2 grs. |
| capsici..... | 8 to 15 mins. | bromid..... | $\frac{1}{2}$ to 2 grs. |
| catechu comp..... | $\frac{1}{2}$ to 2 fl. drs. | iodid..... | $\frac{1}{2}$ to 3 grs. |
| emulsi-fuge..... | 30 to 60 mins. | oxid..... | 1 to 10 grs. |
| emulsiouæ..... | $\frac{1}{2}$ to 2 fl. drs. | phosphid..... | $\frac{1}{2}$ to $\frac{1}{4}$ gr. |
| cinchona comp..... | $\frac{1}{2}$ to 2 fl. drs. | sulphas, emet..... | 15 to 30 grs. |
| colchici rad..... | 5 to 15 mins. | valerianus..... | 1 to 6 grs. |
| colchici sem..... | 5 to 15 mins. | | |

THERAPEUTIC TABLE.

MODIFIED FROM TEXT-BOOK OF PHARMACOLOGY, THERAPEUTICS, AND MATERIA MEDICA.

BY T. LAUDER BRUNTON, M. D.

Abortion, Threatened.

REST.
OPIUM.
VIBURNUM PRUNIFOLIUM.

Abscess.

CALX SULPHURATA (SULPHIDE OF CALCIUM). Small doses frequently repeated, to hasten maturation or healing.
COD-LIVER OIL. In scrofulous cases.
HYDROGEN PEROXIDE.
IRON. As a tonic.

Acidity.

ACIDS. Before meals.
ALKALIES. After meals.
CREASOTE
NUX VOMICA. Before meals.
SILVER NITRATE.
TEREBENTHINE.

Acne.

ARSENIC. In chronic acne.
BISMUTH. As ointment or powder.
CALX SULPHURATA.
HOT FOMENTATIONS.
HOT SPOONGING.
IODIDE OF SULPHUR. In chronic cases.
SULPHUR. Internally and externally.
WATER. Hot sponging several times a day.

After-pains.

BELLADONNA. As ointment.
CAMPHOR. Combined with morphia
MORPHIA AND ATROPIA. Hypodermically, $\frac{1}{4}$ - $\frac{1}{2}$ gr. morphia with $\frac{1}{100}$ gr. of atropia.
POULTICES, warm, to the hypogastrium.

Alcoholism.

AMMONIA. Aromatic spirits of, as substitute for alcohol, to be taken when the craving comes on.
ARSENIC. To lessen vomiting
CAPSICUM. As substitute for alcohol.

Alcoholism (continued).

NUX VOMICA. Nervous and general tonic
OPIUM.

Alopecia.

ARSENIC. Internally
CANTHARIDES TINCT. One part to eight of castor oil, rubbed in roots of hair morning and night.

Amenorrhœa.

ALOES. Alone or with iron.
APIOL.
ARSENIC. With iron, in anæmia.
BATHS, hot foot and hip
MANGANESE and permanganate of potassium, j-ij. grs.

Anæmia.

ARSENIC. With, or in place of, iron.
IRON. Chalybeate waters
MANGANESE. Given with iron.
NUX VOMICA. With iron.
PANCREATIN. In feeble digestion.
PEPSIN. In feeble digestion.

Aneurism.

ALIMENT. Low diet; absolute rest.
ERGOT. Adjunct to rest
POTASSIC IODIDE. In doses of 30 grs.

Angina Pectoris.

ETHER. To diminish pain.
MORPHIA. Hypodermically
NITRITE OF AMYL. During paroxysms.
NITRO-GLYCERINE. Begin with one drop of 1 per cent. solution.
TURPENTINE. Locally to chest during paroxysms.

Anus, Fissure of.

BELLADONNA. Locally.
CASTOR OIL. To keep motions soft.
IODOFORM. Locally, to heal and relieve pain
OPIUM AND GALL OINTMENT. Locally.

Annus. Fissure of (*continued*).

SILVER NITRATE.
TANNIN. Locally.
ZINC CHLORIDE.

Aphthae.

BORIC ACID. In honey or glycerine, alone or with chlorate of potash.
MERCURY. Hydr. cum creta, to remove indigestion.
POTASSIC CHLORATE. As wash, 10 grs. to the oz., alone or with borax.
RHUBARB. To remove indigestion.
SULPHUROUS ACID. Locally.

Apoplexy.

ACONITE. To lower blood-pressure where pulse is strong.
BLEEDING.
COLD WATER. To head when face congested.
CROTON OIL. One or two drops on back of tongue.
ELATERIUM. In suppository or enema during attack.

Ascites.

ARSENIC. In old persons with feeble heart.
DIGITALIS. In cardiac dropsy.
ELATERIUM. As hydragogue cathartic.
JABORANDI. In anasarca.
SODIUM CHLORIDE SOLUTION.

Asthma.

ACONITE. In spasmodic cases, also in asthma consequent on nasal catarrh in children.
AMYL NITRITE. To check paroxysm.
ARSENIC. In small doses in cases associated with bronchitis or hay fever.
BELLADONNA. Internally, in large doses.
BROMIDES.
CHLORAL. During paroxysm.
CHLOROFORM. Inhaled.
GRINDELIA ROBUSTA.
MORPHIA. Hypodermically.
NUX VOMICA.
PEPSIN. To prevent attacks.
POTASSIC IODIDE. In large doses, when asthma due to acute bronchial catarrh.
POTASSIC NITRATE. Inhalation of fumes.
QUINIA. In intervals, when attacks are periodical.
STRAMONIUM. In cigarettes, or 20 grs. of dried leaves mixed with nitrate of potash, and fumes inhaled.
TOBACCO. Smoking sometimes beneficial.

Balanitis.

ASTRINGENT LOTIONS. Alum: sulphate of zinc. $\frac{1}{4}$ gr.-f3j.
MERCURY. $\frac{1}{4}$ gr. chlor. corrosiv. to an oz. lime-water, as lotion.

Bed Sores.

ALUM. With white of egg, as local application.
CHARCOAL. Dry, or in poultices.
IODOFORM.

Biliousness.

ALOES. In constipation.
EUONYMIN.
IRIDIN.
MERCURIAL CATHARTICS. In moderate doses night and morning.
MINERAL ACIDS. Nitro-hydrochloric especially useful.
PODOPHYLLUM. In place of mercury when stools dark.

Bladder, Irritable.

BELLADONNA.
BENZOIC ACID. In large prostate and alkaline conditions of urine.
CANTHARIDES. Not to be used in states of acute inflammation.

Bladder, Paralysis of.

CANNABIS INDICA. In retention, from spinal disease.
CANTHARIDES. In atonic bladder.
ERGOT. In paralysis of bladder or sphincter.
STRYCNINA.

Blepharitis.

BISMUTH. Equal parts of the subnitrate and glycerine locally.
MERCURY (UNG. HYDRARG. NITRATIS). If too strong, dilute with vaseline or simple ointment.

Boils.

ARSENIC.
BELLADONNA. Internally or locally.
CALX SULPHURATA (SULPHIDE OF CALCIUM). In small doses to abort or hasten maturation.
COLLODION. Painted over surface in papular stage, and over base, leaving centre free, in pustular stage.
OPIMUM. Locally to remove pain.
POULTICES.

Breath, Fetid.

BORIC ACID AND POTAS. CHLORATE.
CAMPHOR.
PERMANGANATE OF POTASH. As wash to mouth.

Bright's Disease.

BATHS. Hot water and air.
 BROOM INFUSION. As diuretic.
 CAFFEINE.
 ELATERIUM. As cathartic.
 GALLIC ACID. Lessens albumen.
 JABORANDI.
 LEAD. Lessens albumen.
 NITROUS ETHER.
 POTASSIC B. TARTRATE. Cathartic and diuretic.
 POTASSIC IODIDE.
 TURPENTINE. As diuretic.
 WATER. In large draughts.

Bronchiectasis.

CHLORINE. Inhalation, to lessen fetor.
 IODINE. Inhalation.
 PHOSPHATES AND HYPOPHOSPHITES.
 QUINIA.

Bronchitis, Acute.

ACONITE. $\frac{1}{4}$ -1 m. every hour at commencement.
 AMMONIUM CARBONATE. Where much expectoration and depression.
 IPECACUANHA. When expectoration scanty with dryness in chest.
 NITRIC ACID. When expectoration too copious.
 QUINIA. To reduce temperature.
 TARTAR EMETIC. In dry stage.
 TURPENTINE. When expectoration is profuse.

Bronchitis, Capillary.

AMMONIUM CARBONATE. Much expectoration and lividity.
 CAMPHOR. As expectorant.
 IPECACUANHA. As expectorant and emetic.
 MUSTARD. In poultices.
 POULTICES. Over whole chest.

Bronchitis, Chronic.

AMMONIA. For difficulty in expectoration.
 AMMONIUM. Useful for old people.
 ASAFOETIDA.
 BALSAM OF PERU, AND BALSAM OF TOLU. When expectoration copious.
 CAMPHOR.
 COD-LIVER OIL.
 DIGITALIS. Where heart is feeble.
 EUCALYPTUS.
 IODINE. By inhalation.
 IRON.
 OPIUM. To lessen secretion and cough.
 PHOSPHATES.
 SENEGA. Where expulsion is feeble.

Bronchitis, Chronic (continued).

SERPENTARIA. Like SENEGA.
 SQUILL. Where expectoration is thick.
 SULPHUR. Where exp. ctoration is copious.
 TAR. To lessen secretion and allay cough.
 TEREBENTH.
 TURPENTINE.

Bronchorrhœa.

ASAFOETIDA
 EUCALYPTUS.
 LEAD ACETATE. To lessen secretion

Bruises.

ACONITE, liniment
 CAPSICUM. To remove discoloration of bruise
 OPIUM. Locally applied.

Bubo.

ICE.
 IODINE.
 NITRIC ACID. Locally to indolent bubo.
 POTASSIO-TARTRATE OF IRON. Locally and internally.

Burns and Scalds.

BORACIC ACID. Saturated watery solution.
 CARBOLIC ACID. One to six of olive oil, locally.
 COCAINE.
 CREASOTE WATER.
 LEAD CARBONATE. In white pain.
 LIME. Lime-water with linseed oil
 SODA. Bicarbonate.
 WARM BATH. Whole body immersed for days in extensive burns or scalds. It relieves pain, although it may not save life.

Calculi, Biliary.

CHLOROFORM. Inhalation in paroxysm.
 COUNTER-IRRITANTS. Mustard poultices.
 MORPHIA. $\frac{1}{4}$ gr. with $\frac{1}{12}$ gr. atropia subcutaneously in paroxysm.
 SODIUM PHOSPHATE. In 20 or 30 gr. doses before meals as prophylactic.

Calculi, Renal and Vesical.

ALKALIES.
 AMMONIUM BENZOATE.
 ANÆSTHETICS. To relieve pain.
 CHLOROFORM.
 MORPHIA. Hypodermically.

Cancer.

BELLADONNA. Locally.
 CHARCOAL, poultices, and dry.
 CITRIC ACID. As lotion.

Cancer (continued).

IODOFORM. Locally.
OPIUM. Locally and internally.

Cancerum Oris.

ARSENIC. Internally.
BORIC ACID.
NITRIC ACID. Locally.
POTASSIC CHLORATE.

Carbuncle.

BELLADONNA EXTRACT. Locally.
COLLODION.
POULTICES.
STRAPPING. Leaving centre free.

Catarrh, Acute Nasal.

ACONITE. Internally at commencement.

ACONITE AND BELLADONNA. In sore throat and cold, one drop of tinct. aconite to two of belladonna every hour.

BELLADONNA. 5 m. of tinct., and afterward one or two every day, for profuse watery secretion.

CAMPBOR. As inhalation from steaming water.

COCAINE.

IODINE. By inhalation.

OPIUM. Dover's powder at commencement.

QUINIA. 10 gr. of quinia at commencement.

Cerebral Congestion.

ACONITE. In acute cases.

BELLADONNA.

BROMIDES.

CATHARTICS.

HYDROCYANIC ACID.

VENESECTION.

Chancre.

IODOFORM. Powdered.

Chancroid.

CAUSTICS. Rarely necessary.

IODOFORM. Very useful.

Cholera Asiatica.

ALCOHOL. Iced brandy, to stop vomiting.

CAMPBOR.

SALINE INJECTIONS.

Cholera Infantum.

BISMUTH

FOOD. PREDIGESTED.

MERCURY. Gray powder or calomel.

OPIUM.

POTASSIC BROMIDE. In nervous irritation.

ZINC OXIDE. With bismuth and pepsin.

Chorea.

ARSENIC.

CHLORAL.

CONIUM.

HYOSCIN HYDROBROMATE.

IRON. In anemia.

STRYCHNIA.

Colic, Intestinal.

ASAFETIDA.

CHLOROFORM.

ESSENTIAL OILS. ANISEED, CARDAMOM, CINNAMON, CLOVES, PEPPERMINT.

MORPHIA. Hypodermically.

POULTICES.

Colic, Lead.

ALUM.

CASTOR OIL.

IODIDE OF POTASSIUM.

MAGNESIUM SULPHATE.

Condylomata.

BISMUTH. Locally.

CALOMEL. Locally.

CHROMIC ACID. Locally.

MERCURY. Wash with chlorine-water or chlorinated soda, and dust with calomel.

Conjunctivitis.

BELLADONNA. Locally.

BLISTERS. Behind ear.

BORIC ACID. Solution.

CASTOR OIL. Drop in eye.

MERCURY. Citrine ointment.

SASSAFRAS PITH INFUSION.

Constipation.

ALOES.

BELLADONNA EXTRACT.

CASCARA SAGRADA.

COLOCYNTH COMPOUND EXTRACT.

LIQUORICE POWDER COMPOUND.

MERCURY.

NUX VOMICA. 5-10 $\mathfrak{m}$ in a glass of cold water before breakfast or before dinner.

PHYSOSTIGMA. 10 $\mathfrak{m}$ of tincture along with belladonna and nux vomica.

PODOPHYLLUM. In biliousness.

SALINE WATERS. Friedrichshalle, Hunyadi Janos, or Pullna.

SENNA

WATER. Draught in the morning before breakfast.

Convulsions, Infantile.

ASAFETIDA. A small dose in an enema arrests convulsions from teething.

BATHS. Warm, with cold effusions to the head.

BROMIDE OF POTASSIUM.

Convulsions. Infantile (*cont'd*).

CHLORAL. 5 grs. per mouth or rectum.

CHLOROFORM. To arrest fit

VALERIAN. When due to worms.

Cough. See *Bronchitis*.

ACONITE. In throat-cough

APOMORPHIA. In bronchitis, with deficient secretion

ASAPETIDA. In the after-cough from habit, and in the sympathetic whooping cough of mothers

CELESTIUM. In convulsive and spasmodic cough

HYDROCYANIC ACID. For irritable cough

HYOSCYAMUS. In tickling night-coughs

PRUNUS VIRGINIANA

VALERIAN. In hysterical cough.

Croup.

ALUM. Teaspoonful, with honey or syrup, every $\frac{1}{2}$ or $\frac{1}{4}$ hour until vomiting is induced

HOT APPLICATIONS TO NECK.

IPECACUANHA

SULPHUROUS ACID. As spray.

Cystitis.

ALKALIES. When urine is acid.

BELLADONNA. To allay irritability.

BORIC ACID. Injected

CANTHARIDES. In small doses.

NAPHTHALIN.

OPUM. By enema or suppository, to relieve pain.

TURPENTINE.

Delirium Tremens.

ALCOHOL. Sometimes necessary.

BEEF TEA.

BROMIDES. In large doses.

CAPSICUM.

DIGITALIS. In large doses has had some success

FOOD, nutritious, more to be depended upon than anything else.

HYOSCIN HYDROBROMATE.

Diabetes Mellitus.

ARSENIC.

CODEIA. Sometimes requires to be pushed to the extent of 18 grs. or more per diem

GLYCERINE. As remedy, and as food in place of sugar.

SKIM-MILK DIET.

Diarrhœa.

ACONITE. In high fever and cutting abdominal pains

BISMUTH.

CAMPHOR.

Diarrhœa (*continued*).

CASTOR OIL AND OPIUM. At commencement.

CATTIEU.

CORROSIVE SUBLIMATE. In small doses.

KINO.

LEAD ACETATE.

NUX VOMICA. In chronic cases.

OPUM.

ZINC OXIDE.

ZINC SULPHATE.

Diphtheria.

ALCOHOL.

ARSENIC.

CHLORINE-WATER. Internally and locally

HYDROCHLORIC ACID. Dilute as gargle, or strong as caustic.

IRON PERCHLORIDE.

LACTIC ACID. As solvent of false membrane

MERCURY. Calomel or corrosive sublimate internally

PAPAIN. As solvent of false membrane.

POTASSIUM CHLORATE. Internally and locally.

STRYCHNIA

SULPHUROUS ACID.

Dropsy. See *Ascites*.

BROOM.

ERYONIA. Purgative and diuretic.

COLCHICUM. In hepatic, cardiac, and scarlatinal dropsy.

DIGITALIS. Especially in cardiac dropsy; infusion is the best form.

ELATERIUM.

IRON.

JUNIPER

NITROUS ETHER.

POTASSIUM BITARTRATE. With compound jalap powder.

Dysentery.

ACONITE.

BELLADONNA.

CASTOR OIL. In small doses, with opium

CORROSIVE SUBLIMATE. In small doses in chronic cases

ENEMATA, of starch and laudanum, and of nitrate of silver

HAMAMELIS. Much blood in motions.

IPECACUANHA. In 30-gr. doses on empty stomach, with complete rest.

LEAD ACETATE. By mouth, enema, or suppository.

NUX VOMICA.

OPUM.

TANNIN.

Dysentery (*continued*).

ZINC OXIDE.
ZINC SULPHATE.

Dysmenorrhœa.

APIOL (Oil of Parsley). In neuralgic form.
ARSENIC. When membranous discharge.
BELLADONNA. In neuralgic form.
CANNABIS INDICA. Very useful.
CROTON CHLORAL. In neuralgic form.
GELSEMIUM.
IRON.
NUX VOMICA.
OPIUM. 3 to 5 ms. of tincture, alone or with 3 or 4 gr. of chloral.

Dyspepsia.

ACIDS. Especially nitro-muriatic acid
ALKALIES.
ARSENIC.
ASAFOETIDA.
BELLADONNA.
BISMUTH.
CAPSICUM.
CHARCOAL.
CHOLAGOGUES.
FOOD. Nutritious
HYDROCYANIC ACID.
LACTIC ACID.
LIME-WATER.
MAGNESIUM SULPHATE.
NUX VOMICA.
PANCREATIN.
PEPSIN.
SILVER NITRATE.
TARAXACUM.

Dysuria.

ALKALIES.
BELLADONNA.
CAMPHOR.
CANNABIS INDICA.
CANTHARIDES, TINCTURE.
ERGOT
GELSEMIUM.
NITROUS ETHER.
OPIUM.

Earache.

ALMOND OIL.
ATROPIA SOLUTION. Locally.
GLYCERINE.
OPIUM.
CARBOLIZED OIL.

Eczema.

ALKALIES. Weak solutions of, as constant dressing.
ARSENIC.
BENZOLIN. Locally.
BISMUTH. Locally.

Eczema (*continued*).

LEAD SALTS. Locally.
LIME-WATER. Alone or with glycerine or oil, locally.
ZINC OXIDE. Ointment.

Emphysema. See *Asthma* and *Bronchitis*.

Endocarditis.

ACONITE. At commencement.
BLISTERS.
OPIUM.

Endometritis.

CARBOLIC ACID.
GLYCERINE.
HOT-WATER INJECTIONS.
IODINE.
NITRIC ACID.

Epididymitis.

ACONITE. In small doses frequently repeated.
MERCURY AND MORPHIA. Locally as oleate.
OINTMENT OF MERCURY AND BELLADONNA.
PULSATILLA. In very small doses along with aconite.

Epilepsy.

ARSENIC.
ASAFOETIDA.
BELLADONNA.
BROMIDES OF POTASSIUM, SODIUM, LITHIUM, AND IRON.
HYDROBROMIC ACID.
NITRITE OF AMYL. By inhalation.
NITRO-GLYCERINE.
QUINIA.
STRYCHNIA.
VALERIAN.
ZINC OXIDE.

Epistaxis.

ALUM. In powder or solution, locally.
DIGITALIS.
GALLIC ACID.
VINEGAR.

Epithelioma.

LACTIC ACID.

Erysipelas.

ACONITE. At commencement.
IRON. Large doses of tincture of chloride frequently.
QUININE
STARCH and other dry powders.
SULPHUROUS ACID.

Fever. See *Intermittent*, *Typhoid*, etc.

ACIDS. To allay thirst.
ALCOHOL. Small doses frequently.

Fever (continued).

ANTIPYRINE.
 ARSENIC. In malarious fevers.
 BELLADONNA. In eruptive fevers.
 CAMPHOR. In adynamic fevers.
 DIGITALIS.
 SYRUP OF HYDRIODIC ACID.
 MERCURY. Small doses at commencement of typhoid or scarlet fever.
 SALICIN, { In rheumatic fevers or
 SALICYLATE OF SODA, { In hyper-
 SALICYLIC ACID, { pyrexia.
 TERPENTINE. As stimulant.

Fistula.

PEPPER. The confection, as laxative.

Flatulence.

ALKALIES. Before meals.
 ASAFOETIDA.
 CARMINATIVES.
 CHARCOAL.
 CREASOTE.
 NUX VOMICA.

Gangrene.

BALSAM OF PERU.
 BROMINE.
 CHARCOAL. As poultice or dry.
 NITRIC ACID.

Gastralgia.

ATROPIA.
 BISMUTH.
 CHLOROFORM. Two or three drops on sugar.
 MORPHIA. Subcutaneously or internally, combined with chloroform.

Gastric Ulcer.

ARSENIC.
 BISMUTH.
 GALLIC ACID.
 LEAD ACETATE. To check hemorrhage.
 MILK.
 MORPHIA.
 PEPSIN.
 SILVER NITRATE.
 SILVER OXIDE.

Gastritis.

ARSENIC.
 BISMUTH.
 NUX VOMICA.
 OPIUM.
 SILVER NITRATE.
 SILVER OXIDE.

Gleet.

CANTHARIDES.
 IRON.

Gleet (continued).

SANDAL-WOOD OIL. Locally and internally.
 TANNIN. GLYCERINE OF. Locally.
 TERPENTINE.

Gout.

IODINE. Internally and locally ointment or tincture, and by injection.
 MERCURI BIODIDE. As ointment.

Gonorrhoea.

ALKALINE salts or waters.
 ALUM. As an injection.
 CANTHARIDES.
 CYREBS. Alone or mixed with copaiba.
 SILVER NITRATE. As injection.
 ZINC SALTS. As injection.

Gout.

ACONITE.
 ALKALIES.
 ARSENIC.
 BELLADONNA.
 COLCHICUM.
 LITHIUM.
 VERATRIA. As ointment.

Hay Fever.

ACONITE.
 ARSENIC.
 IODIDE OF POTASSIUM. Internally and locally.

Hæmoglobinuria.

ARSENIC.

Headache.

ACONITE. When circulation excited.
 AMMONIA. Aromatic spirits $\frac{1}{2}$ -2 dr. doses.
 BELLADONNA. In frontal headache, especially at menstrual period, or from fatigue.
 BICARBONATE OF SODA. With bitters before meals in frontal headache.
 BROMIDE OF POTASSIUM.
 BRYONIA. In bilious headache.
 CAMPHOR. Internally and externally.
 CANNABIS INDICA. In neuralgic headache.
 CHLOROFORM, SPIRITS OF. In nervous headache.
 DIGITALIN. $\frac{\pi}{\pi}$ of a grain twice a day for congestive hemicrania.
 GUARANA.
 HEAT. As hot-water bag or poultice to nape of neck.

Headache (*continued*).

IODIDE OF POTASSIUM. In rheumatic headache, with tenderness of scalp.

MAGNESIA SULPHATE. For frontal headache, with constipation.

MENTHOL. Locally.

MERCURY. In bilious headache.

NUX VOMICA. In nervous or bilious headache.

SPECTACLES. Where the headache depends on astigmatism.

TEA.

VALERIAN. In nervous and hysterical cases.

Heart, Dilated.

ADONIDINE.

DIGITALIS.

MORPHIA.

Heart, Fatty.

ARSENIC.

BELLADONNA.

IRON.

STRYCHNIA.

Heart, Hypertrophied.

ACONITE. To be used with care when valvular disease is present.

CAMPHOR. In palpitation and dyspnoea.

DIGITALIS. In small doses.

VERATRUM VIRIDE.

Heart, Palpitation of.

ADONIDINE.

BELLADONNA. Internally, useful in cardiac strain.

BROMIDE OF POTASSIUM. In fluttering heart.

CAMPHOR.

DIGITALIS.

HYOSCYAMUS. In nervous palpitation.

VALERIAN. In nervous cases with dyspnoea.

Hematemesis.

ALUM.

GALLIC ACID.

ICE.

LEAD ACETATE.

SULPHURIC ACID.

TANNIC ACID.

TURPENTINE.

Hematuria.

ALKALINE DIURETICS.

ALUM.

BITARTRATE OF POTASH.

DIGITALIS.

GALLIC ACID.

LEAD ACETATE.

Hematuria (*continued*).

TANNIC ACID.

TURPENTINE.

Hemoptysis.

ALUM.

DIGITALIS.

ERGOT.

GALLIC ACID. Very useful.

LEAD ACETATE.

TURPENTINE.

Hemorrhage, Intestinal.

GALLIC ACID.

LEAD ACETATE.

TANNIC ACID.

TURPENTINE.

Hemorrhage, Post-partum.

ALCOHOL.

DIGITALIS.

ERGOT.

HOT WATER. Injection into uterus.

ICE. To abdomen, uterus, or rectum.

IRON PERCHLORIDE, DILUTED. 1:4 injected into uterus.

PRESSURE OVER UTERUS.

Hemorrhoids.

ALKALINE mineral waters useful

CARBOLIC ACID. Injection into piles.

ERGOT.

GALLS, OINTMENT. With opium.

IODOFORM. In ointment or suppository.

NUX VOMICA. Very useful.

SENNA. Confection or compound liquorice powder.

SULPHUR.

Hepatitis.

AMMONIUM CHLORIDE.

BRYONIA.

NITRO-HYDROCHLORIC ACID.

Herpes Zoster.

ACONITE AND OPIUM. Locally.

MORPHIA.

OIL OF PEPPERMINT. Locally

Hiccough.

AMYL NITRITE.

BELLADONNA.

CAMPHOR.

CANNABIS INDICA.

CHLORAL.

CHLOROFORM.

ETHER.

LAUREL-WATER.

MORPHIA. Hypodermically.

NUX VOMICA.

Hydrocephalus, Acute.

BLISTERS. To nape of neck.

Hydrocephalus, Acute (cont'd).

BROMIDE OF POTASSIUM.
 IODIDE OF POTASSIUM.
 LEECHES.
 MERCURIC CHLORIDE. Small doses

Hydrocephalus, Chronic.

BLISTERS.
 COD-LIVER OIL.
 IODIDE OF IRON.
 IODIDE OF POTASSIUM.
 MERCURY.

Hydrophobia.

CANNABIS INDICA.
 CHLOROFORM, and hypodermic of morphia $\frac{1}{4}$ grain.

Hydrothorax.

BLISTERS.
 BROOM.
 DIGITALIS.
 ELATERIUM.
 JABORANDI.
 SODIUM CHLORIDE.

Hypochondriasis.

ALCOHOL.
 ASAFCETIDA.
 BROMIDE OF POTASSIUM.
 HYOSCYAMUS.
 MUSK.

Hysteria.

ALOES. In constipation.
 AMMONIA, AROMATIC SPIRITS OF ASAFCETIDA.
 BROMIDE OF CAMPHOR.
 BROMIDE OF POTASSIUM.
 CAMPHOR.
 IGNATIA.
 IRON.
 SPIRITUS ETHERIS NITROSI.
 VALERIAN.
 ZINC VALERIANATE.

Ichthyosis.

BATHS.
 COD-LIVER OIL.
 ELM BARK DECOCTION.
 GLYCERINE.

Impetigo.

ARSENIC.
 COD-LIVER OIL.
 GUTTA-PERCHA. Locally, in solution.
 HYDROCYANIC ACID. To relieve itching.
 LAUREL-WATER. To relieve itching.
 QUININE.
 SULPHUR. Internally.
 ZINC OINTMENT.

Impotence.

ARSENATE OF IRON.
 CANTHARIDES.

Impotence (continued).

NUX VOMICA.
 PHOSPHORUS.
 ZINC PHOSPHATE.

Influenza.

OPIUM. With ipecacuanha
 POTASSIUM NITRATE. Freely diluted, as lemonade.
 QUININE. In later stages
 SPIRITUS ETHERIS NITROSI.

Insomnia.

ACONITE. 1 m. every quarter of an hour, when skin dry and harsh.
 ALCOHOL.
 CAMPHOR.
 CANNABIS INDICA. Alone, or with hyoscyamus
 CHLORAL.
 CROTON CHLORAL. If heart weak.
 DIGITALIS. When deficient tone of vaso-motor system
 FOOD at bed-time.
 GELSENIUM. In simple wakefulness.
 HYOSCIN HYDROBROMATE.
 HYOSCYAMUS. Alone, or with cannabis indica.
 PARALDEHYDE.
 POTASSIUM BROMIDE. In full doses.
 URETHAN.
 WARMTH. Internally and externally.
 WET COMPRESS.

Intermittent Fever.

ARSENIC.
 ATROPIA. Subcutaneously, to arrest or cut short cold stage
 CAMPHOR.
 CAPSICUM. With quinine.
 CHLOROFORM. To prevent or cut short cold stage
 GELSENIUM. Pushed until it produces dilated pupils or double vision
 IODINE, TINCTURE. To prevent recurrence of ague
 OPIUM. In full doses, to prevent chill.
 PEPPER. With quinine.
 QUININE.
 QUININE BROMOHYDRATE. Like quinine, and less liable to produce cinchonism.
 SODIUM CHLORIDE. Tablespoonful in glass of hot water at a draught on empty stomach.

Intertrigo.

BISMUTH. Locally.
 FULLER'S EARTH.
 LEAD LOTION.
 LIME-WATER.

Iritis.

ATROPIA. Locally.
 BELLADONNA. Internally and locally.
 DUBOISIA. Substitute for atropia.
 MERCURY. Most serviceable.

Jaundice.

ALKALINE mineral waters in catarrh of duodenum or bile-ducts.
 AMMONIUM CHLORIDE. In scruple doses.
 CALOMEL. Often very useful.
 ENEMATA. Large.
 ETONYMIN.
 IRIDIN.
 MAGNESIUM SULPHATE.
 NITRO-HYDROCHLORIC ACID.
 PODOPHYLLUM
 QUININE. In malarious cases.

Lactation, Defective.

CASTOR-OIL LEAVES. Poultice or infusion, to breast.
 GOSSEYPIUM. Decoction of seeds two or three times every hour.
 JABORANDI.

Lactation, Excessive.

BANDAGE.
 BELLADONNA. Internally and locally.
 CAMPHOR AND GLYCERINE. Locally.
 IODIDE OF POTASSIUM.

Laryngismus Stridulus.

ACONITE.
 BELLADONNA.
 BROMIDES.
 CHLORAL.
 CHLOROFORM. By inhalation.
 COLD SPONGING.
 EMETICS.
 MORPHIA. Hypodermically.
 QUININE.

Laryngitis, Acute.

ACONITE.
 COCAINE.
 MORPHIA.
 QUININE.
 SCARIFICATION OF LARYNX.
 SULPHUROUS ACID. As inhalation or spray.
 ZINC SULPHATE. As emetic.

Laryngitis, Chronic.

ALUM. As gargle.
 AMMONIUM CHLORIDE. As spray.
 BISMUTH. Locally.
 CARBOLIC ACID. As spray.
 FERRIC CHLORIDE. Locally.
 SILVER NITRATE. Solution locally.
 TANNIN. As gargle or spray.

Leucorrhœa.

ALUM. By injection.
 BALSAM OF PERU. Internally.
 BELLADONNA. Locally.
 BORAX. By injection.
 GLYCERINE.
 IODOFORM. Locally.
 IRON. Internally.
 POTASSIUM PERMANGANATE. Locally.
 TANNIN. By injection or suppository.

Locomotor Ataxia.

CANNABIS INDICA.
 ELECTRICITY.
 ERGOT.
 PHOSPHORUS.
 POTASSIUM IODIDE.
 STRYCHNIA.

Lumbago.

ACONITE. Internally and locally.
 CHLOROFORM, liniment.
 HOT DOUCHE. Or poultice.
 ICE.
 IODIDE OF POTASSIUM.
 MORPHIA.
 QUININE.

Lupus.

ARSENIC.
 IODIDE OF POTASSIUM.
 IODIDE OF SULPHUR.
 IODOFORM.
 LACTIC ACID.
 PYROGALLIC ACID.
 SODIUM ETHYLATE.
 ZINC CHLORIDE.

Mania.

CANNABIS INDICA.
 CHLORAL. As narcotic and carminative.
 DIGITALIS.
 HYOSCYAMUS.
 HYOSCYN HYDROBROMATE.
 IRON.
 OPIUM.
 PARALDEHYDE.
 PHYSOSTIGMA.
 URETHAN.
 WET PACK.

Mastitis.

AMMONIUM CHLORIDE. Locally.
 BELLADONNA. Locally.
 MERCURY AND MORPHIA OLEATE. Locally.
 SUPPORT, by bandaging or strapping.

Measles.

ACONITE.

Measles (continued).

DIGITALIS.
 POTASSIUM BROMIDE. For sleeplessness.
 QUININE.

Melancholia.

ARSENIC.
 BROMIDE OF POTASSIUM.
 CANNABIS INDICA.
 CHLORAL.
 HYOSCIN HYDROBROMATE.
 IRON.
 OPIUM.
 URETHAN.
 VALERIAN.

Meningitis, Cerebral.

ACONITE.
 BELLADONNA.
 BLISTERS. To nape of neck.
 BROMIDE OF POTASSIUM.
 COLD, to head.
 HYOSCYAMUS.
 MERCURY.
 OPIUM. In small doses, alone or with tartar emetic.
 POTASSIUM IODIDE.
 PURGATIVES.
 VENESECTION.

Meningitis, Cerebro-Spinal.

ACONITE.
 BELLADONNA.
 CAUTERY, to back.
 COLD, to spine.
 DIGITALIS.
 ERGOT.
 OPIUM.
 QUININE.
 VENESECTION.

Menorrhagia.

ARSENIC.
 BROMIDE OF POTASSIUM.
 CANNABIS INDICA.
 DIGITALIS.
 ERGOT.
 GALLIC ACID.
 HOT-WATER BAG to dorsal and lumbar vertebræ.
 IRON.
 QUININE.
 SULPHURIC ACID.

Metritis, Acute.

ACONITE.
 HOT WATER. Vaginal injection.
 OPIUM. By suppository or enema.
 POULTICES.
 TURPENTINE STUPES.

Myalgia.

AMMONIUM CHLORIDE
 ARNICA. Internally and locally

Myalgia (continued).

BELLADONNA LINIMENT. Locally and internally.
 CHLOROFORM LINIMENT.
 FRICTION.
 GELSEMIUM. Large doses.
 OPIUM.
 PACKING
 POTASSIUM IODIDE.
 SALICYLATE OF SODA.

Myelitis.

BELLADONNA.
 ERGOT
 PHOSPHORUS.
 SILVER NITRATE.

Nævus.

CHROMIC ACID.
 ELECTROLYSIS.
 GALVANO-CAUTERY.
 LIGATION.
 NITRIC ACID.

Nausea.

BISMUTH.
 CLOVES
 HYDROCYANIC ACID.
 ICI.
 IPECACUANHA. In sickness of pregnancy and chronic alcoholism: 1 ℥ of wine, repeated frequently.
 MORPHIA.
 PEPPERMINT.
 SPIRIT AMMON. AROMATIC.

Nephritis, Acute.

ACONITE. At commencement.
 BELLADONNA
 CANNABIS INDICA. As diuretic, especially in hæmaturia
 CANTHARIDES. 1 ℥ of tincture every three hours to stop hæmaturia after acute symptoms have subsided
 DIGITALIS. As diuretic.
 EUCALYPTUS.
 GALLIC ACID.
 JABORANDI.
 JUNIPER.
 POULTICES. Over loins
 TURPENTINE. $\frac{1}{2}$ -1 ℥ doses, every two to four hours.

Nervousness.

ACONITE. 1 ℥ of tincture at bedtime for restlessness.
 BROMIDE OF POTASSIUM.
 CAFFEINE.
 CAMPHOR.
 CHLORAL
 IGNATIA.
 STRYCHNINE

Neuralgia.

ACONITIA.
 AMMONIUM VALERIANATE.
 AMYL NITRITE.
 ATROPINE. In liniment or hypodermically
 BELLADONNA.
 BLISTERS.
 BROMIDE OF POTASSIUM.
 CAFFEIN.
 CANNABIS INDICA.
 CHLORAL AND CAMPHOR.
 CHLOROFORM. Locally, and by inhalation, when pain very severe.
 COD-LIVER OIL.
 CROTON CHLORAL.
 DIGITALIS.
 ELECTRICITY.
 ERGOT.
 GELSEMIUM.
 IODIDES.
 IRON.
 MORPHIA. Hypodermically.
 MUSTARD POUltICE.
 NITRO-GLYCERINE.
 NUX VOMICA.
 OIL OF CLOVES.
 PEPPERMINT.
 PULSATILLA.
 QUININE.
 SALICYLIC ACID.
 THERMO-CAUTERY.
 VALERIAN.
 VIBRATION.
 WATER. Hot or cold, locally.

Nightmare.

BROMIDE OF POTASSIUM.
 CAMPHOR-WATER.

Nipples, Sore.

BALSAM OF PERU.
 BENZOIN.
 BORAX.
 COLLODION.
 LEAD SALTS.
 TANNIN.

Nodes.

MERCURY.
 POTASSIUM IODIDE.

Nymphomania.

BROMIDE OF POTASSIUM.
 CAMPHOR.
 LUPULINE.

Obesity.

ALKALINE WATERS.
 AMMONIUM BROMIDE.
 BANTING'S SYSTEM—living on meat and green vegetables, and avoiding starch, sugars, and fats.

Onychia.

IODOFORM.
 POTASSIUM PERMANGANATE.
 STRAPPING.

Ophthalmia. See *Conjunctivitis*.**Orethritis.**

MERCURY AND BELLADONNA. Ointment.
 STRAPPING.

Otitis.

BORACIC ACID. Locally.
 DRY CLEANSING.
 HYDROGEN PEROXIDE.
 SILVER NITRATE. Solution.
 ZINC SULPHATE.

Ozena.

ALKALINE WASHES.
 IODINE. As inhalation; washing out nose with solution of common salt, to which a few drops of the tincture of iodine have been added
 IODOFORM.
 POTASSIUM PERMANGANATE. Solution.
 TANNIN, GLYCERINE OF.

Parotitis.

ACONITE.
 JABORANDI.
 MERCURY $\frac{1}{2}$ gr. of gray powder three or four times a day.
 POUltICE.

Pediculi.

MERCURY. As ointment or wash.

Pemphigus.

ARSENIC.
 COD-LIVER OIL.
 IODIDE OF POTASSIUM.
 MERCURY.
 PHOSPHORUS.

Pericarditis.

ACONITE.
 BLISTERS. Near heart.
 BRYONIA.
 DIGITALIS. Heart rapid and feeble, with cyanosis and dropsy.
 OPIUM. In grain doses every three to six hours.
 QUININE.
 VERATRUM VIRIDE.

Periostitis.

IODIDE OF POTASSIUM, or AMMONIUM.
 IODINE. Locally.
 POUltICES.

Peritonitis.

ACONITE. At commencement.
 BRYONIA. When exudation
 LEECHES.
 MERCURY.
 OPIUM. Freely used
 POUltICES.
 TURPENTINE.

Perspiration.

AROMATIC SULPHURIC ACID.
 ATROPIA
 BELLADONNA. In liniment, for local
 sweats
 JABORANDI.
 OPIUM
 QUININE
 SPONGING, very hot.
 STRYCHNINE.
 VINEGAR.

Pertussis.

ACONITE.
 ALUM.
 BELLADONNA.
 BROMIDE OF POTASSIUM.
 CHLORAL.
 CHLOROFORM. By inhalation.
 COD-LIVER OIL.
 CROTON CHLORAL.
 ERGOT.
 IPECACUANHA.
 MONOBROMIDE OF CAMPHOR.
 NITRITE OF AMYL.
 OPIUM
 VALERIAN.

Pharyngitis.

ACONITE.
 ALUM. Gargle.
 CAPSICUM. Gargle.
 COCAINE.
 FERRIC CHLORIDE. Locally and in-
 ternally
 GLYCERIN. Locally.
 POTASSIUM CHLORATE. Locally.
 QUININI
 SILVER NITRATE. Locally
 STRYCHNINE.
 TANNIN. As powder, or glycerine,
 locally.
 ZINC SULPHATE. Gargle.

Phlebitis.

HAMAMELIS.
 HOT FOMENTATIONS.
 OPIUM.
 REST, absolute.

Phlegmasia Alba.

AMMONIUM CARBONATE. In full
 doses, when much prostration.
 BELLADONNA EXTRACT. With mer-
 curial ointment locally.

Phlegmasia Alba (continued).

HAMAMELIS.
 OPIUM. Internally and locally.

Phthisis.

ALCOHOL. With food or cod-liver
 oil
 ANTIPYRINE. To reduce tempera-
 ture
 ARSENIC. To remove commencing
 consolidation, and when tongue is
 red and irritable
 ATROPIA. To check perspiration.
 BENZOIN. As inhalation to lessen
 cough and expectoration
 CHLORAL.
 CHLOROFORM, to check cough.
 COCAINE. Solution 4-20 per cent. to
 throat and mouth to relieve irrita-
 tion.
 COD-LIVER OIL.
 DIGITALIS
 GLYCERINE. In place of cod-liver
 oil.
 HYPOPHOSPHITES.
 IODINE LINIMENT.
 IRON
 MERCURIC CHLORIDE. In minute
 doses for diarrhoea.
 MORPHIA. WITH STARCH OR BIS-
 MUTH. To larynx.
 OPIUM.
 OXYGEN
 PHOSPHATE OF LIME.
 PILOCARPINE. To check sweats.
 QUININE.
 SALICYLIC ACID. For foul breath
 and offensive expectoration.

Pleurisy. See Hydrothorax.

ACONITE.
 BELLADONNA PLASTER.
 BLISTERS.
 BRYONIA. After aconite.
 DIGITALIS.
 IODIDE OF POTASSIUM. To aid ab-
 sorption.
 LOCAL WET PACK.

Pleurodynia.

BELLADONNA. Plaster or liniment.
 CHLORAL. With camphor locally.
 ETHER. Locally, in spray.
 IRON
 MUSTARD-LEAVES.
 OPIUM.
 POUltICES.
 QUININI
 STRAPPING.

Pneumonia.

ACONITE. At commencement.
 AMMONIUM CARBONATE. As stimu-
 lant toward end.

Pneumonia (*continued*).

BELLADONNA. At commencement.
 BRYONIA. When pleurisy present.
 DIGITALIS
 OPIUM.
 POULTICES.
 QUININE.
 SALICYLATE OF SODA. As anti-pyretic.
 TURPENTINE. As stimulant.

Polypus.

ALUM.
 TANNIN. By insufflation.

Prolapsus Ani.

ALUM. Locally.
 ICE.
 CAUTERIZATION.
 STRYCHNIA.
 TANNIN. Locally.

Prolapsus Uteri.

ALUM. By vaginal douche.
 GALLS. Decoction by injection.
 TANNIN.

Prostate, Enlarged.

ALKALIES. For irritation of bladder, with acid urine.
 AMMONIUM BENZOATE. For cystitis with alkaline urine.
 CONIUM.
 ERGOT.
 IODOFORM. By suppository.

Prurigo and Pruritus.

ARSENIC. Internally.
 BALSAM OF PERU.
 BORACIC ACID.
 CALOMEL. Ointment.
 CHLORAL AND CAMPHOR.
 COD-LIVER OIL. By inunction.
 CORROSIVE SUBLIMATE. Wash.
 GLYCERINE.
 GLYCEROLE OF TAR.
 HYDROCYANIC ACID. Locally.
 IODOFORM. Ointment.
 OPIUM.
 OIL OF CADE.
 TAR OINTMENT.

Psoriasis.

ALKALINE BATH.
 ARSENIC.
 CALOMEL. Ointment.
 CHRYSOPHANIC ACID.
 COD-LIVER OIL.
 GLYCERINE.
 INDIA-RUBBER SOLUTION.
 LEAD IODIDE. Locally.
 PHOSPHORUS. Substitute for arsenic.
 POTASSIUM IODIDE.

Psoriasis (*continued*).

SULPHUR. Internally.
 TAR. Ointment.
 ULMUS.
 VASELINE.

Ptyalism.

ACIDS. Small doses, and in gargles.
 BELLADONNA
 CHLORATE OF POTASH. Gargle
 TANNIN.

Puerperal Convulsions.

BELLADONNA.
 BLEEDING.
 CHLORAL. In full doses.
 CHLOROFORM. By inhalation.
 MORPHIA. Hypodermically.
 VERATRUM VIRIDE. Pushed to nausea.

Puerperal Fever.

ACONITE. At commencement.
 OPIUM. For wakefulness and delirium, very useful.
 QUININE. In large doses.
 TURPENTINE
 VERATRUM VIRIDE.

Puerperal Mania.

ANÆSTHETICS.
 BROMIDE OF POTASSIUM.
 CHLORAL.
 HYOSCYAMUS.
 OPIUM
 QUININE

Puerperal Peritonitis.

ACONITE.
 LAXATIVES.
 OPIUM.
 QUININE.
 TURPENTINE.

Purpura Hemorrhagica.

ERGOT.
 GALLIC ACID.
 IRON.
 NUX VOMICA.
 QUININE.
 TURPENTINE.

Pyæmia.

IRON.
 OPIUM.
 PERMANGANATE OF POTASH.
 QUININE.
 TURPENTINE.

Pyrosis.

ACIDS, MINERAL.
 ALKALIES.
 BISMUTH

Pyrosis (continued).

CAMPHOR.
CARBOLIC ACID.
CREASOTE.
GALLIC ACID.
GLYCERINE.
NUX VOMICA.

Rheumatism, Acute.

ACONITE.
ALKALIES.
AMMONIUM BROMIDE.
ANTIPYRIN.
BLISTERS.
COLD BATHS.
DIGITALIS.
IRON.
LIME-JUICE.
OPIUM.
QUININE.
SALICYLATE OF SODA. Relieves pain most quickly.

Rheumatism, Chronic.

BELLADONNA.
BROMIDE OF LITHIUM.
COD-LIVER OIL.
GUAIAC.
IODIDE OF POTASSIUM.
MERCURY AND MORPHIA. Oleate locally.
QUININE.
SULPHUR. Locally and as sulphides, or sulphur waters internally.
TURKISH BATHS.

Rickets.

CALCIUM PHOSPHATE.
COD-LIVER OIL.
COLD SPONGING.
IODIDE OF IRON.
QUININE.

Scabies.

BAKING of clothes to destroy ova.
BALSAM OF PERU. Locally, agreeable and effective.
MERCURY. White precipitate ointment.
PETROLEUM.
SULPHUR. Ointment.

Scarlet Fever. (Treatment chiefly symptomatic.)

ACONITE.
BELLADONNA.
CHLORINE-WATER. As gargle.
COLD COMPRESS to throat.
DIGITALIS.
FAT, Inunction over the whole body.
PACKING. Useful and comforting.
SALICYLATE OF SODA. As antipyretic.
SPONGING.

Sciatica.

ACUPUNCTURE.
ATROPIA.
BELLADONNA.
BLISTERS.
CAUTERY. Slight application.
CHLOROFORM. Liniment.
ELECTRICITY.
ETHER. Spray.
MORPHIA.
POULTICES.
TURPENTINE. In $\frac{1}{4}$ -oz. doses for 3 or 4 nights.

Serofula.

CALCIUM PHOSPHATE.
COD-LIVER OIL.
IODIDE OF IRON.

Scurvy.

ACIDS.
ALUM. With myrrh, for ulcerated gums.
CITRIC ACID.
LIME-JUICE.
QUINIA. With mineral acids.
VINEGAR.

Sea-Sickness.

BROMIDE OF POTASSIUM.
BROMIDE OF SODIUM.
CANNABIS INDICA.
CHAMPAGNE.
CHLORAL.
CHLOROFORM. 2-5 ms. on sugar.
COCAINE.
NITRO-GLYCERINE.
NUX VOMICA.

Seborrhœa.

BORAX, with GLYCERINE.

Spermatorrhœa.

ARSENIC.
BELLADONNA.
BROMIDES.
CAMPHOR.
CANTHARIDES.
ELECTRICITY.
NUX VOMICA.
PHOSPHORUS.
TURPENTINE.

Spinal Congestion.

ACONITE.
ERGOT.
NUX VOMICA.

Spinal Irritation.

ACONITE OINTMENT.
ARSENIC.
BELLADONNA.

Spinal Irritation (*continued*).

BLISTERS
CONIUM
COUNTER-IRRITATION.
DIGITALIS.
ERGOL.
NUX VOMICA.
OPIUM.
PHOSPHORUS.

Sprains.

AMMONIUM CHLORIDE.
ARNICA.
BANDAGING.
COMENTATIONS.
MASSAGE.

Stomatitis.

ACIDS, MINERAL.
ALUM.
BORAX.
CHLORATE OF POTASSIUM.
GLYCERINE OF TANNIN.
NITRIC ACID.

Sweating. See *Perspiration*.**Sycosis.**

CORROSIVE SUBLIMATE, solution 2-3
grs. to 15 of water.
EPILATION.
SULPHUROUS ACID.

Synovitis.

ACONITE.
ARNICA.
BANDAGE. OR STRAPPING.
BLISTERS
COD-LIVER OIL.
IODINE. Injection.
MASSAGE.

Syphilis.

ARSENIC.
BITTER TONICS.
COD-LIVER OIL.
IODIDES.
IODOFORM
IRON.
MERCURY.
POTASSIC IODIDE.

Tabes Mesenterica.

CHLORIDE OF LIME.
COD-LIVER OIL.
FATTY INUNCTION.
PHOSPHORIC ACID.
SARSAPARILLA.

Tetanus.

BROMIDES.
CHLORAL.
GELSEMIUM.
NEUROTOMY.
OPIUM.
PHYSOSTIGMA.

Tetanus (*continued*).

URETHAN.
WARM BATHS.

Tic Douloureux.

ACONITIN
AMMONII CHLORIDUM.
ARSENIC.
CANNABIS INDICA.
MORPHIA.
PHYSOSTIGMA.
VERATRIA.

Tinea Circinata.

COD-LIVER OIL.
IODINE, TINCTURE. Locally.
MERCURY. Locally.
SULPHUROUS ACID. Locally.

Tinea Favosa.

ARSENIC
EPILATION.
HYPOSULPHITES
MERCURY. Locally.
SULPHUROUS ACID.

Tinea Tonsurans.

CHRYSOGRAPHIC ACID.
COD-LIVER OIL.
EPILATION
MERCURY OLEATE.
PARASITICIDE LOTIONS.

Tonsillitis.

ACONITE
ALUM.
BELLADONNA.
BICARBONATE OF SODA. Dry, locally.
COCAINE.
GUAIAC.
ICE AND WET PACK.
MERCURY.
OPIUM
POTASSIC IODIDE.
TANNIC ACID.

Tonsils. Enlarged.

AMMONIUM IODIDE
IRON, tincture of chloride.
MASSAGE.
SILVER NITRATE.
TANNIN.

Tonsils. Ulcerated.

BICARBONATE OF SODA. Dry locally.
IRON, tincture.
MERCURIC IODIDE.
POTASSIUM CHLORATE.
SULPHUROUS ACID.

Toothache.

BICARBONATE OF SODA.
CARBOLIC ACID.
CLOVES, OIL.
CREASOTE.

Torticollis.

ARSENIC.
ELECTRICITY.
MASSAGE.
MERCURY.
NUX VOMICA.

Trismus.

ANESTHETICS.
BELLADONNA.
CHLORAL.
OPIUM.
PHYSOSTIGMA.

Tympanites.

ACIDS AND ALKALIES.
ASAFOETIDA.
CARBOLIC ACID.
HYOSCYAMUS.
NUX VOMICA.
TURPENTINE.
VEGETABLE CHARCOAL.

Typhlitis.

BELLADONNA.
ICE-BAG.
LEECHES.
MERCURY.
OPIUM.

Typhoid Fever.

ALCOHOL.
ANTIPYRETICS.
ANTIPYRIN.
ASAFOETIDA.
BATHS.
CALOMEL.
CAMPHOR.
CARBONATE OF AMMONIA.
COLD COMPRESS.
DIGITALIS.
HYDROCHLORIC ACID.
HYOSCYAMUS.
MURIATIC ACID.
OPIUM.
PEPSIN.
QUININE.
TURPENTINE.

Typhus Fever.

ALCOHOL.
CAMPHOR.
DIGITALIS.
HYOSCYAMUS.
MINERAL ACIDS.
OPIUM.
QUININE.
STRYCHNIA.

Varicose Veins.

HAMAMELIS.

Vomiting. See *Nausea*.

AMMONIUM SALTS AND AROMATIC
SPIRITS.

Vomiting *continued*.

ARSENIC.
BELLADONNA.
BISMUTH.
BLISTERS.
BROMIDES.
CALOMEL.
CARBOLIC ACID.
CERUUM OXALATE.
CHLOROFORM.
CINNAMON.
CLOVES.
CREASOTE.
COUNTER-IRRITATION.
HYDROCYANIC ACID.
ICE.
IPECACUANHA.
KOUSSO.
MORPHIA.
NUTRIENT ENEMATA.
NUX VOMICA.
OPIUM.
PEPSIN.
STRYCHNIA.
ZINC.

Warts.

ACIDS, MINERAL.
ALKALIES, CAUSTIC.
CHROMIC ACID.
SILVER NITRATE.

Worms.

ALUM. Injection for thread-worms.
CASTOR OIL.
EUCALYPTOL. Injection for thread-
worms.
FILIX MAS. For tape-worms.
IGNATIA.
IRON. Injection for thread-worms
and as a tonic.
KAMALA. For tape-worms.
KOUSSO. For tape-worms.
LIME-WATER. Injection for thread-
worms.
PEPO.
QUINIA.
QUASSIA. Injection for thread-
worms.
SANTONINE. For round-worms.
SODIUM CHLORIDE. Injection for
thread-worms.
TURPENTINE. For tape-worms.

Yellow Fever.

CALOMEL.
CAMPHOR.
CHLORODYNE.
IPECACUANHA.
LEAD ACETATE.
NUX VOMICA.
QUININE.
STIMULANTS.
TURPENTINE.

LIGATION OF ARTERIES.


THE accompanying plate indicates the positions in which the operation of ligating an artery in its continuity is most commonly performed. There is no point at which ligatures are applied which is not represented, with the exception of a few which are very rarely chosen. With the explanations which follow, this diagram will serve as a guide to the practitioner who may be called upon in an emergency to perform an operation in regard to which he has grown somewhat rusty from want of practice.

1. Ligation of the Temporal Artery.—The temporal artery is the continuation of the external carotid from the level of the lobe of the ear to the level of the eyebrow, where it divides into its terminal branches. The point at which it is ligated is a little in front of the anterior curve of the concha. A vertical incision, about an inch in length, is made in front of the concha. The artery is easy to find, as it is very superficial at this part of its course. Its accompanying vein lies to the outer side, and the temporal branch of the auriculo-temporal nerve lies in front of it.

2. Ligation of the Occipital Artery.—The occipital artery comes off from the posterior part of the external carotid at about the level of the angle of the lower jaw. It curves backward under and to the inner side of the mastoid process, at the back of which it comes out through the attachment of the trapezius muscle, and winds in a tortuous course upward over the occipital bone. The point at which it is ligated is about an inch behind the ear. An incision an inch and a half long, parallel to the posterior edge of the mastoid process, will expose it. Its accompanying vein lies to the outer side of it. (There are but few occasions when it is necessary to ligate this artery, as its position makes it very easy to stop the flow of blood through it by pressure with a suitable compress and bandage.)

3. Ligation of the Facial Artery.—The facial artery comes off from the external carotid just below the level of the angle of the lower jaw, and passes forward below this bone to a point just in front of the insertion of the masseter muscle. Here it sends a branch forward, the submental, while the main trunk curves up over the lower jaw and passes in a tortuous course toward the angle of the mouth. The point at which it is ligated is that at which it ascends in front of the masseter muscle. An incision an inch in length will expose it. Its accompanying vein lies to the outer side.

4. Ligation of the Lingual Artery.—The lingual artery comes off from the external carotid at the level of the great cornu of the hyoid bone; it passes forward under the posterior belly of the digastric muscle, and the styloglossus and hyoglossus muscles, to the root of the tongue, and thence to its further distribution. It is in some respects the most difficult artery to ligate in the whole body. The point at which it is ligated is the second part of its course, under the hyoglossus muscle. An incision is made about half an inch above, and parallel to, the great horn of the hyoid bone, and extending from just in front of the anterior edge of the sterno-cleido-mastoid muscle to near the middle line of the neck, half an inch below the base of the jaw. The skin and subcutaneous fat having been divided, the capsule of the submaxillary gland will be met. This capsule must be divided, and the gland carefully drawn up over the jaw. In doing this great care must be taken not to injure the facial artery and vein, which pass through the gland. The posterior part of the capsule of the gland must next be divided. This will expose the shining aponeurosis which constitutes the pulley of the digastric muscle and attaches it to the great horn of the hyoid bone, near the insertion of the stylo-hyoid muscle. Behind these passes the hypoglossal nerve. This must be delicately detached from the sheath of the hyoglossus muscle, which lies below it, and pushed up out of the way. As the next step, fix the pulley of the digastric with a tenaculum and gently insinuate a director behind the posterior edge of the hyoglossus muscle, and push it forward between this and the middle constrictor of the pharynx, which lies under



neath it, in a line parallel to, and just above, the greater horn of the hyoid bone. The hyoglossus muscle is then to be divided on the director, and the lingual artery is exposed. In placing the ligature the needle should be passed from above downward, so as not to injure the hypoglossal nerve. The important guides in this ligation are the hyoid bone, the pulley of the digastric muscle, and the hypoglossal nerve.

5 and 6. Ligation of the Common Carotid Artery.—For ligation, the common carotid artery is divided into two portions by the narrow omohyoid muscle, which crosses it near its middle. (Shown in the cut.) The upper portion is comparatively superficial, and occupies what on this account is called the triangle of election. The lower portion lies deeper, and occupies what for this reason is called the triangle of necessity. (The cut represents the incisions for both these ligations united.)

5. Ligation of the Common Carotid Artery in the Triangle of Election.—The patient should be recumbent, with the head extended and turned toward the opposite side. An incision should be made along the anterior edge of the sterno cleido-mastoid muscle, from about an inch below the angle of the lower jaw to a little below the cricoid cartilage. The superficial fascia, the platysma-myoides muscle, and the deep fascia are then to be divided on a director. The sterno-mastoid muscle must then be held back, and the deep layer of the deep fascia picked up and carefully divided on a director. This will expose the sheath of the blood vessels, and perhaps the *descendens noni* nerve. Avoiding the latter, the sheath of the vessels is to be picked up with forceps and nicked with a scalpel. A director is then to be introduced, and the sheath slit up upon it. The artery is now to be separated from the internal jugular vein, which lies to the outer side, and from the pneumogastric nerve, which lies back of and between the artery and vein. The ligature should be passed from without inward.

6. Ligation of the Common Carotid Artery in the Triangle of Necessity.—The position of the patient should be the same as for the operation just described. The incision should be made along the inner edge of the sterno mastoid muscle, and from the middle of the neck to the top of the sternum. The skin, superficial fascia, and platysma myoides muscle are to be divided. Then the deep fascia is to be divided on a director, avoiding injury to the anterior jugular vein. When the sides of the wound are drawn asunder, another layer of the deep fascia will be exposed, which must be lifted with forceps, nicked, and then carefully divided on a director. Under this lies the sheath of the blood-vessels, and upon it the *descendens noni* nerve. The subsequent steps of the operation are the same as those described for ligation in the triangle of election.

7. Ligation of the Subclavian Artery.—The point of election for ligation of this artery is the outer side of the *scalenus anticus* muscle. An incision is made parallel to and just above the clavicle, from the outer half of the clavicular insertion of the sterno cleido-mastoid muscle to the anterior edge of the trapezius muscle, taking care not to go deep enough to cut the external jugular vein. The skin, superficial fascia, and the platysma-myoides muscle are to be divided. Then the external jugular vein is to be drawn aside. If this be impossible, it must be divided between two ligatures. The loose connective tissue must now be carefully broken up with a director and a search made for the artery. It must be remembered that it lies to the outer side of the *scalenus anticus* muscle and to the inner side of the cords of the brachial plexus. When found, it is to be very cautiously separated from its attachments to the pleura, and the ligature passed from below upward. The greatest care is necessary to avoid injuring the vein or the pleura, or including a nerve in the ligature.

8 and 9. Ligation of the Axillary Artery.—This artery is ligated in its first portion just after passing over the first rib, and in its third portion in the axilla.

8. Ligation of the First Portion of the Axillary Artery.—The arm is drawn away from the side, and an incision about three inches in length is made in the line which marks the division between the upper and lower parts of the pectoralis major muscle. The superficial and deep

lumen must be divided on a director. The muscular interspace is then to be sought for, and the two parts of the muscle separated, working up toward the clavicle, and not directly backward. As this is done, the arm should be brought back to the side, so as to relax the parts. The costo-cervical membrane is next to be found and opened on a director. The artery lies beneath it. It is to be carefully isolated from the axillary vein, which lies on its inner, or lower, side, and the axillary nerves, which lie to its outer side. The ligature should be passed from within outward, or from below upward.

9. *Ligation of the Axillary Artery in the Axilla.*—This is the most accessible part of the artery. It is reached by drawing the arm away from the body, and making an incision about three inches long over the course of the vessel, at the junction of the anterior and middle thirds of the axilla. The deep fascia must be divided on a director, and the artery will be found beneath it, with the axillary vein on the ulnar side (toward the back of the arm), and the median nerve on the radial side (toward the front of the arm). The ligature should be passed from the ulnar toward the radial side, so as not to injure the vein.

10. *Ligation of the Brachial Artery, in the middle of the arm.*—An incision about two inches in length is to be made along the inner edge of the swell of the biceps muscle, at the middle of the arm. After dividing the skin and superficial fascia, and any fat that may be met with, the fibrous membrane which encloses the biceps muscle will be encountered. This is to be divided on a director, and the loose connective tissue along the edge of the muscle broken up, when the artery will be exposed. Care must be taken not to divide the median nerve, which crosses the brachial artery near its middle, passing from the radial toward the ulnar side, and not to include it in the ligature. The *venae comites* lie one on each side of the artery.

11. *Ligation of the Radial Artery, in the upper third.* An incision is to be made in a line drawn from the middle of the bend of the elbow to the point at which the pulse can be felt at the wrist. The median vein is to be avoided in incising the superficial fascia. When the deep fascia has been exposed, a yellow line, marking the division between the supinator longus, on the radial side, and the pronator radii teres, on the ulnar side, is to be looked for. The fascia is to be opened along this line, and the artery will be found below it. The radial nerve lies to the radial side of the artery, but so far away as to be in little danger of being injured.

12. *Ligation of the Ulnar Artery, in the lower third.* An incision is to be made in the depression between the flexor sublimis digitorum and the flexor carpi ulnaris, just above the carpus. The skin and superficial fascia are to be divided, and then the deep fascia is to be picked up with the forceps and divided on a director. The artery will be found lying on the pronator quadratus, alongside the tendon of the flexor carpi ulnaris, with the ulnar nerve on its ulnar side. The ligature should be passed from the ulnar toward the radial side, avoiding the nerve.

13. *Ligation of the External Iliac Artery.*—An incision is to be made nearly parallel to Poupart's ligament, and about an inch above it, extending from about an inch above the anterior superior spinous process of the ilium to a point near the external abdominal ring. This incision should curve slightly upward at both ends. The skin and superficial fascia are to be divided, and ligatures to be applied to any vessels which bleed. The tendon of the external oblique muscle is to be divided, and then the internal oblique and the transversalis muscles. The separation of the transversalis fascium is a work of the utmost delicacy, to be effected with the forceps and the finger, the latter restraining and protecting the peritoneum. A finger passed down into the pelvis through the opening will discover the artery. The accompanying iliac vein lies to the inner side of the artery, and the anterior crural nerve at some distance to its outer side. The ligature should be passed from within outward.

14. *Ligation of the Femoral Artery in Scarpa's Triangle.*—The point of election for ligation of the femoral artery is at the apex of Scarpa's triangle. To accomplish it, the leg should be flexed at the knee, and the thigh slightly flexed upon the abdomen, abducted and rotated

outward, so as to lie on its outer side. An incision should then be made, over the course of the artery, in a line bisecting Scarpa's triangle from the middle of its base to its apex. This incision should be about three inches long, and its middle should be over the apex of the triangle. The skin, superficial fascia, and fat having been divided, the inner edge of the sartorius muscle must be found and drawn outward. The artery will now be found, with the vein to the inner side and a little behind it, and several nerves on the outer side. The artery should be carefully isolated after opening its sheath, and the ligature passed from within outward.

15. Ligation of the Femoral Artery in Hunter's Canal.

This rare ligation is performed by placing the limb in the position for the former operation, and making an incision from the apex of Scarpa's triangle toward the back part of the internal condyle of the femur, avoiding the saphena vein. After passing through the skin, superficial fascia, and fat, the deep fascia will be seen, binding down the sartorius muscle. This membrane is to be opened on a director and the muscle drawn backward. This will disclose the aponeurotic sheath which covers the femoral artery at this point. This sheath must be opened, and the artery will be exposed. In ligating it, care must be taken not to injure the long saphenous nerve, which accompanies it, or the femoral vein, which here lies behind it and toward its outer side. The ligature should be passed very carefully from without inward.

16. Ligation of the Anterior Tibial Artery, in the middle

third.—An incision is to be made, about three inches long, in a line passing from a point midway between the tubercle of the tibia and the head of the fibula, to a point midway between the two malleoli. When the deep fascia has been reached, a yellow line may be found, indicating the septum between the tibialis anticus muscle and the extensor longus digitorum. Along this line the aponeurosis is to be divided on a director, and the two muscles carefully separated until the extensor proprius pollicis is met with, when this is to be separated from the tibialis anticus until the artery is reached. It is accompanied by two *venæ comites*, which it is not important to regard if they are difficult to separate from the artery. The anterior tibial nerve lies either in front or at the outer side of the artery. To avoid injuring it, it should be pressed to the outer side of the wound and the ligature should be passed from without inward.

17. Ligation of the Posterior Tibial Artery, in the middle

third.—An incision three inches long is to be made parallel to the inner border of the tibia and about half an inch back of it, and carried down to the superficial fascia. This is to be carefully opened, avoiding injury to the internal saphenous vein and nerve, which lie in the track of the wound. They must be found and drawn aside. Then the deep fascia is to be divided on a director, and the inner border of the soleus muscle is be pressed away from the tibia, when the vessel will be exposed. It is accompanied by two *venæ comites* and the posterior tibial nerve, which lies to the fibular side of it. The veins and nerve being separated from the artery, the ligature should be passed from the fibular toward the tibial side.

18. Ligation of the Posterior Tibial Artery behind the

Internal Malleolus.—A slightly curving incision an inch and a half long is to be made behind the internal malleolus, and about half-way between it and the tendo Achillis. The skin and superficial fascia having been divided, the deep fascia is to be lifted and divided on a director. The artery will then be exposed, accompanied by its *venæ comites*, and the posterior tibial nerve, which lies to the back of it and toward the fibular side. The ligature should be passed from behind forward.

19. Ligation of the Dorsalis Pedis Artery.

An incision an inch long is to be made on the back of the foot between the inner tendon of the extensor brevis digitorum and the outer side of the tendon of the extensor longus pollicis. (These tendons can be mapped out by manipulating the great toe.) After dividing the skin, the superficial fascia, and the deep fascia, the artery will be exposed. The continuation of the anterior tibial nerve lies to the fibular side of the artery. The ligature should be passed from the fibular toward the tibial side.

THE
MEDICAL NEWS
VISITING LIST.

RECORDS OF PRACTICE,
FOR THIRTY PATIENTS PER WEEK,
1887.

NAME.

Su. M. Tu. W. Th. F. Sa.

1

Record.

January, 1887.

51

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 2 | 3 | 4 | 5 | 6 | 7 | 8 |

Mr. Horn

Ct.

Q.

" (C. or Merton¹ & Pillsbury
 West. in Bellhop + 34 Market

C

C

MEMORANDA.

CHARGE.

Page 11

Ledger

| NAME. | Su. | M. | Tu. | W. | Th. | F. | Sa. |
|---|-----|----|---------|----|-----|----|-----|
| | 9 | 10 | 11 | 12 | 13 | 14 | 15 |
| a in Belknap | C | | Dec. 14 | | | | |
| Fallen. Grand Canyon. mile 7. | | | C | | | | |
| C. with D. 9.15 in. Miller. 530 Dauphin | | | | | | | C. |

Record.

January, 1887.

55

MEMORANDA.

CHARGE.

Page in
Ledge

Record.

January, 1887.

57

MEMORANDA.

CHARGE.

Page in
Lodge 36.

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 23 | 24 | 25 | 26 | 27 | 28 | 29 |

MEMORANDA.

CHARGE.

Page in
Ledger

NAME.

| Su | M. | Tu. | W. | Th. | F. | Sa |
|----|----|-----|----|-----|----|----|
| 30 | 31 | 1 | 2 | 3 | 4 | 5 |

the ... of ...
 ... the ...

to

MEMORANDA.

CHARGE.

Page in
Ledger.

Record.

February, 1887.

63

MEMORANDA.

CHARGE.

Page in

Lodge #100

Record.

February, 1887.

65

MEMORANDA.

CHARGE.

Page in
Ledger.

MEMORANDA.

| CHARGE.

Page in
Ledger.

Record.

February—March, 1887.

69

MEMORANDA.

CHARGE.

Page in
Ledger

| NAME. | Su. | Mo. | Tu. | W. | Th. | F. | Sa. |
|----------|-----|-----|-----|----|-----|----|-----|
| | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
| Nov. 11 | / | / | / | / | 7 | / | / |
| Nov. 12 | | | | | | | |
| Nov. 13 | | | | | | | |
| Nov. 14 | | | | | | | |
| Nov. 15 | | | | | | | |
| Nov. 16 | | | | | | | |
| Nov. 17 | | | | | | | |
| Nov. 18 | | | | | | | |
| Nov. 19 | | | | | | | |
| Nov. 20 | | | | | | | |
| Nov. 21 | | | | | | | |
| Nov. 22 | | | | | | | |
| Nov. 23 | | | | | | | |
| Nov. 24 | | | | | | | |
| Nov. 25 | | | | | | | |
| Nov. 26 | | | | | | | |
| Nov. 27 | | | | | | | |
| Nov. 28 | | | | | | | |
| Nov. 29 | | | | | | | |
| Nov. 30 | | | | | | | |
| Nov. 31 | | | | | | | |
| Nov. 32 | | | | | | | |
| Nov. 33 | | | | | | | |
| Nov. 34 | | | | | | | |
| Nov. 35 | | | | | | | |
| Nov. 36 | | | | | | | |
| Nov. 37 | | | | | | | |
| Nov. 38 | | | | | | | |
| Nov. 39 | | | | | | | |
| Nov. 40 | | | | | | | |
| Nov. 41 | | | | | | | |
| Nov. 42 | | | | | | | |
| Nov. 43 | | | | | | | |
| Nov. 44 | | | | | | | |
| Nov. 45 | | | | | | | |
| Nov. 46 | | | | | | | |
| Nov. 47 | | | | | | | |
| Nov. 48 | | | | | | | |
| Nov. 49 | | | | | | | |
| Nov. 50 | | | | | | | |
| Nov. 51 | | | | | | | |
| Nov. 52 | | | | | | | |
| Nov. 53 | | | | | | | |
| Nov. 54 | | | | | | | |
| Nov. 55 | | | | | | | |
| Nov. 56 | | | | | | | |
| Nov. 57 | | | | | | | |
| Nov. 58 | | | | | | | |
| Nov. 59 | | | | | | | |
| Nov. 60 | | | | | | | |
| Nov. 61 | | | | | | | |
| Nov. 62 | | | | | | | |
| Nov. 63 | | | | | | | |
| Nov. 64 | | | | | | | |
| Nov. 65 | | | | | | | |
| Nov. 66 | | | | | | | |
| Nov. 67 | | | | | | | |
| Nov. 68 | | | | | | | |
| Nov. 69 | | | | | | | |
| Nov. 70 | | | | | | | |
| Nov. 71 | | | | | | | |
| Nov. 72 | | | | | | | |
| Nov. 73 | | | | | | | |
| Nov. 74 | | | | | | | |
| Nov. 75 | | | | | | | |
| Nov. 76 | | | | | | | |
| Nov. 77 | | | | | | | |
| Nov. 78 | | | | | | | |
| Nov. 79 | | | | | | | |
| Nov. 80 | | | | | | | |
| Nov. 81 | | | | | | | |
| Nov. 82 | | | | | | | |
| Nov. 83 | | | | | | | |
| Nov. 84 | | | | | | | |
| Nov. 85 | | | | | | | |
| Nov. 86 | | | | | | | |
| Nov. 87 | | | | | | | |
| Nov. 88 | | | | | | | |
| Nov. 89 | | | | | | | |
| Nov. 90 | | | | | | | |
| Nov. 91 | | | | | | | |
| Nov. 92 | | | | | | | |
| Nov. 93 | | | | | | | |
| Nov. 94 | | | | | | | |
| Nov. 95 | | | | | | | |
| Nov. 96 | | | | | | | |
| Nov. 97 | | | | | | | |
| Nov. 98 | | | | | | | |
| Nov. 99 | | | | | | | |
| Nov. 100 | | | | | | | |

MEMORANDA.

CHARGE.

Page in
Ledger.

Cheque for \$1.0 to Henry Fairbank
455 Bernard St. in Cash
Guild 208 Broadway L.S.

March, 1887.

Daily

| NAME. | Su.
13 | M.
14 | Tu.
15 | W.
16 | Th.
17 | F.
18 | Sa.
19 |
|-----------------------------------|-----------|----------|-----------|----------|-----------|----------|-----------|
| Miss Ripple. | / | / | / | / | / | | / |
| 1. Gardner | | | 10 | | | | |
| Wm. B. Williams 22. 3rd Jan. 1877 | | | | | c | c | |
| Miss Thompson | | | | | | c | |
| Miss | | | | | | | |

65-

* fine little institute in my neighborhood

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 27 | 28 | 29 | 30 | 31 | 1 | 2 |

1. 10 1/2

2. 10 1/2

3. 10 1/2

4. 10 1/2

5. 10 1/2

1 1 1 1 1

c

c

c

c

60

MEMORANDA.

CHARGE.

Page in
Ledger

NAME

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 3 | 4 | 5 | 6 | 7 | 8 | 9 |

/

/

/

/

/

/

11 30

MEMORANDA.

CHARGE.

Page in
Ledger

NAME.

| Su | M | Tu | W | Th | F | Sa |
|----|----|----|----|----|----|----|
| 10 | 11 | 12 | 13 | 14 | 15 | 16 |

John C. Smith
John C. Smith
John C. Smith

John C. Smith
John C. Smith

MEMORANDA.

CHARGE.

Page in
Ledger.

50 cents to date

April, 1887.



Daily

| NAME. | Su. | M. | Tu. | W. | Th. | F. | Sa. |
|---------------|-----|----|-----|----|-----|----|-----|
| | 17 | 18 | 19 | 20 | 21 | 22 | 23 |
| Mrs. ... | / | / | / | / | / | / | X |
| Father ... | | | | | | | |
| Mr. Williams | | | | X | * | * | X |
| Mrs. D. Scott | | | | | | | |
| Mr. ... | | | | | | | / |

H. W.

Sus. ...

Record.

April, 1887.

83

MEMORANDA.

CHARGE.

Page in
Lodge

April, 1887.

Daily

| NAME. | Sa. | M. | Tu. | W. | Th. | F. | Sa. |
|-----------|-----|----|-----|----|---------|----|-----|
| | 24 | 25 | 26 | 27 | 28 | 29 | 30 |
| 1. | X | X | | | | | |
| 2. | / | | | - | 15 days | 2 | W |
| 3. | | | / | | | | |
| 4. | | | | | | | 1 |
| 5. | | | | | | | 0 |

Record.

April, 1887.

85

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 |

C C

4 10

Record.

May, 1887.

87

MEMORANDA.

CHARGE.

Page in
Ledger.

| NAME. | Su. | M. | Tu. | W. | Th. | F. | Sa. |
|---------------------------|-----|----|-----|----|-----|----|-----|
| | 8 | 9 | 10 | 11 | 12 | 13 | 14 |
| Dan. H. Lewis, Louisville | | | C. | | | | |
| to get a | | | C. | | | | |

20

Record.

May, 1887.

89

MEMORANDA.

1 CHARGE.

Page in
Ledger

NAME.

| Sa. | M | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 15 | 16 | 17 | 18 | 19 | 20 | 21 |

G. W.
M. Sellers

1/2 03

Record.

May, 1887.

91

MEMORANDA.

1 CHARGE.

Page in
Ledger.

May, 1887.

Daily

MEMORANDA.

CHARGE.

Page in
Ledger

MEMORANDA.

1 CHARGE.

Page in.

1.1.1.1

rich and beautiful. Forest of the country
high, the road ridge and the stream, in the
west of the land.

[illegible]

June, 1857.

Daily

| NAME. | Su | M. | Tu. | W | Th. | F. | Sa |
|--------------------------|----|----|-----|---|-----|----|----|
| | 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| 1. 1000 ft. to 1000 ft. | | | | | | | |
| 2. 1000 ft. to 1000 ft. | | | | | | | |
| 3. 1000 ft. to 1000 ft. | | | | | | | |
| 4. 1000 ft. to 1000 ft. | | | | | | | |
| 5. 1000 ft. to 1000 ft. | | | | | | | |
| 6. 1000 ft. to 1000 ft. | | | | | | | |
| 7. 1000 ft. to 1000 ft. | | | | | | | |
| 8. 1000 ft. to 1000 ft. | | | | | | | |
| 9. 1000 ft. to 1000 ft. | | | | | | | |
| 10. 1000 ft. to 1000 ft. | | | | | | | |
| 11. 1000 ft. to 1000 ft. | | | | | | | |
| 12. 1000 ft. to 1000 ft. | | | | | | | |
| 13. 1000 ft. to 1000 ft. | | | | | | | |
| 14. 1000 ft. to 1000 ft. | | | | | | | |
| 15. 1000 ft. to 1000 ft. | | | | | | | |
| 16. 1000 ft. to 1000 ft. | | | | | | | |
| 17. 1000 ft. to 1000 ft. | | | | | | | |
| 18. 1000 ft. to 1000 ft. | | | | | | | |
| 19. 1000 ft. to 1000 ft. | | | | | | | |
| 20. 1000 ft. to 1000 ft. | | | | | | | |
| 21. 1000 ft. to 1000 ft. | | | | | | | |
| 22. 1000 ft. to 1000 ft. | | | | | | | |
| 23. 1000 ft. to 1000 ft. | | | | | | | |
| 24. 1000 ft. to 1000 ft. | | | | | | | |
| 25. 1000 ft. to 1000 ft. | | | | | | | |
| 26. 1000 ft. to 1000 ft. | | | | | | | |
| 27. 1000 ft. to 1000 ft. | | | | | | | |
| 28. 1000 ft. to 1000 ft. | | | | | | | |
| 29. 1000 ft. to 1000 ft. | | | | | | | |
| 30. 1000 ft. to 1000 ft. | | | | | | | |
| 31. 1000 ft. to 1000 ft. | | | | | | | |
| 32. 1000 ft. to 1000 ft. | | | | | | | |
| 33. 1000 ft. to 1000 ft. | | | | | | | |
| 34. 1000 ft. to 1000 ft. | | | | | | | |
| 35. 1000 ft. to 1000 ft. | | | | | | | |
| 36. 1000 ft. to 1000 ft. | | | | | | | |
| 37. 1000 ft. to 1000 ft. | | | | | | | |
| 38. 1000 ft. to 1000 ft. | | | | | | | |
| 39. 1000 ft. to 1000 ft. | | | | | | | |
| 40. 1000 ft. to 1000 ft. | | | | | | | |
| 41. 1000 ft. to 1000 ft. | | | | | | | |
| 42. 1000 ft. to 1000 ft. | | | | | | | |
| 43. 1000 ft. to 1000 ft. | | | | | | | |
| 44. 1000 ft. to 1000 ft. | | | | | | | |
| 45. 1000 ft. to 1000 ft. | | | | | | | |
| 46. 1000 ft. to 1000 ft. | | | | | | | |
| 47. 1000 ft. to 1000 ft. | | | | | | | |
| 48. 1000 ft. to 1000 ft. | | | | | | | |
| 49. 1000 ft. to 1000 ft. | | | | | | | |
| 50. 1000 ft. to 1000 ft. | | | | | | | |
| 51. 1000 ft. to 1000 ft. | | | | | | | |
| 52. 1000 ft. to 1000 ft. | | | | | | | |
| 53. 1000 ft. to 1000 ft. | | | | | | | |
| 54. 1000 ft. to 1000 ft. | | | | | | | |
| 55. 1000 ft. to 1000 ft. | | | | | | | |
| 56. 1000 ft. to 1000 ft. | | | | | | | |
| 57. 1000 ft. to 1000 ft. | | | | | | | |
| 58. 1000 ft. to 1000 ft. | | | | | | | |
| 59. 1000 ft. to 1000 ft. | | | | | | | |
| 60. 1000 ft. to 1000 ft. | | | | | | | |
| 61. 1000 ft. to 1000 ft. | | | | | | | |

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 12 | 13 | 14 | 15 | 16 | 17 | 18 |

Mrs. Lippin

€

Ed. Lillard & Co. 5000

€

Mrs. Walden

/

/

/

/

MEMORANDA.

CHARGE.

Page in
Ledger

22

1. The first of these is the fact that the

1874

[illegible]

2. I have the right, per se, to give away the - intellectual
 creation.

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 19 | 20 | 21 | 22 | 23 | 24 | 25 |

MEMORANDA.

CHARGE.

Page in
Ledger

June 30 Balance
100 00

NAME.

| Su | M. | Tu. | W. | Th. | F | Sa. |
|----|----|-----|----|-----|---|-----|
| 26 | 27 | 28 | 29 | 30 | 1 | 2 |

June 26. Sunday. Clear. 70-80.

June 27. Monday. Clear. 70-80.

June 28. Tuesday. Clear. 70-80.

June 29. Wednesday. Clear. 70-80.

June 30. Thursday. Clear. 70-80.

MEMORANDA.

| CHARGE.

|| Page in
|| Ledger

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 3 | 4 | 5 | 6 | 7 | 8 | 9 |

Jim Webb
Doyle
in Kansas

1 1 1 1
2 0
2

at clerk in Salt Co. Dark. Comp. her hair
Anemic, dark beneath the eyes. ...
when a fellow clerk

Had been bitten on the hip by a dog (not wild) 16 years ago &
18 yrs ago got a scratch on the right thumb from a
dog, which is still alive. Was ever extremely
nervous about the same the statement of the clerk.
Has dwelt with & felt certain ... was going mad
Has had pain in the ... & ... & ...
beliefs. ... not lost a day's work. Set, & ...
as usual. ... slate
ment about the ... of it being ... & ...
been ... He went away ... On the eve of
9th returned with his wife. ... had
red ... could not drink a glass of water & ...
cliffers about the neck. ... at ...
"went ... one crazy" today. ...
... the place ...
...

Record.

July, 1887.

15

MEMORANDA.

CHARGE.

Page in
Ledger

NAME.

| Su. | M | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 10 | 11 | 12 | 13 | 14 | 15 | 16 |

Jan - 5 of 1887

Jan 1887

Miss Grace Lewis, 501 Marlborough St.

received from the Rev. Dr. J. H. ...
1887

Record.

July, 1887.

107

MEMORANDA.

CHARGE.

Page in
Ledger

NAME.

| Sa. | M | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 17 | 18 | 19 | 20 | 21 | 22 | 23 |

Record.

July, 1887.

109

MEMORANDA.

CHARGE.

Page in
Ledger

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 24 | 25 | 26 | 27 | 28 | 29 | 30 |

Record.

July, 1887.

111

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Su | M. | Tu. | W | Th. | F | Sa. |
|----|----|-----|---|-----|---|-----|
| 31 | 1 | 2 | 3 | 4 | 5 | 6 |

Record.

July—August, 1887.

113

MEMORANDA.

CHARGE.

Page in
Ledger

NAME.

| Su | M. | Tu. | W. | Th. | F. | Sa. |
|----|----|-----|----|-----|----|-----|
| 7 | 8 | 9 | 10 | 11 | 12 | 13 |

Record.

August, 1887.

115

MEMORANDA.

CHARGE.

Page in
Ledger

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 14 | 15 | 16 | 17 | 18 | 19 | 20 |

Record.

August, 1887.

117

MEMORANDA.

CHARGE.

Page in
Ledger

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 21 | 22 | 23 | 24 | 25 | 26 | 27 |

2

Record.

August, 1887.

119

MEMORANDA.

CHARGE.

Page in
Ledger

NAME.

| Su. | M | Tu. | W. | Th. | F | Sa. |
|-----|----|-----|----|-----|---|-----|
| 28 | 29 | 30 | 31 | 1 | 2 | 3 |

Record.

August—September, 1887.

121

MEMORANDA.

CHARGE.

Page in
Ledger

September, 1887.

Daily

[illegible]

September, 1887.

Daily

NAME.

| Sa. | M | Tu. | W. | Th. | F | Sa. |
|-----|----|-----|----|-----|----|-----|
| 11 | 12 | 13 | 14 | 15 | 16 | 17 |

Mrs. Pardo

c c c c

Mrs. Sullivan

c

Record.

September, 1887.

125

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Su. | M | Tu. | W | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 18 | 19 | 20 | 21 | 22 | 23 | 24 |

[Faint handwritten notes and numbers, possibly a ledger entry]

[Faint handwritten notes at the bottom of the page]

Record.

September, 1887.

127

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Su | M. | Tu. | W | Th. | F | Sa. |
|----|----|-----|----|-----|----|-----|
| 25 | 26 | 27 | 28 | 29 | 30 | 1 |

Mrs. and Mr.

Mrs. and Mr. 1812 1814

C.

C. 1814 1816

C.

Mrs. and Mr. 1814 1816

C.

C.

John W. 1814 1816

C.

C.

Record.

September—October, 1887.

129

MEMORANDA.

Charge.

Page 46

Letter.

| NAME. | Su | M. | Tu. | W. | Th. | F. | Sa. |
|-------------------------------|----|----|-----|----|-----|----|-----|
| | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| John Emmer 2230 Palms St | C | | | | | | |
| Geo J. Ford. Jackson St 215 | C | | | | | | C |
| Wm. A. H. H. H. | | C | | C | | | |
| C. D. H. H. H. | | | | | | | C |
| C. C. H. H. H. 7th & Col. Ave | | | | | | | C |

Neurotic. ordered Tr. H. H. H.

Record.

October, 1887.

131

MEMORANDA.

CHARGE.

Page 11.

Ledger.

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 9 | 10 | 11 | 12 | 13 | 14 | 15 |

F. P. Davis 15 in and one

c

1. Carter 10 in, 1. Carter 10 in

c

1. Carter 10 in

c

1. Carter 10 in

c

1. Carter 10 in

c

c

Record.

October, 1887.

123

MEMORANDA.

CHARGE.

Page in
Ledger

NAME.

| Sa. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 16 | 17 | 18 | 19 | 20 | 21 | 22 |

Gen Mess, (22 June) /
 Gen Mess 5-5-8 N 17.5

C

C

Record.

October, 1887

125

MEMORANDA.

CHARGE.

Page in
Ledger.

October, 1887.

Dail

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 23 | 24 | 25 | 26 | 27 | 28 | 29 |

C. with Dr. Kelly, Fall of 1887 C.

C. with Dr. Hunt

C.

Record.

October, 1887.

137

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

Su. M. Tu. W. Th. F. S.

30 31

1

2

3

4

5

| | | | | | | |
|--------------------------------|----|--|--|--|--|---|
| Edwards, S. L. - Kane | c. | | | | | |
| Mr. Bunker 1000 Portland | c. | | | | | |
| Mr. S. L. Bellum | c. | | | | | |
| Mr. Filer 2504 Continental Ave | c. | | | | | |
| James 1203 | | | | | | c |
| Isaac Harvey 125 S 23rd | | | | | | c |
| Mr. Carroll Lancaster Ave | | | | | | c |

Record.

October - November, 1887.

139

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Su. | M. | Tu. | W. | Th. | F. | S. |
|-----|----|-----|----|-----|----|----|
| 6 | 7 | 8 | 9 | 10 | 11 | |

C. Smith at noon 60¢ Peter
 Geo - Fred. & D. Sutcliff

C.

L C

Record.

November, 1887.

141

MEMORANDA.

CHARGE.

Page in
Ledger.

66

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa. | MEM. |
|-----|----|-----|----|-----|----|-----|------|
| 13 | 14 | 15 | 16 | 17 | 18 | 19 | |

Miss L.
 from her home (Lafayette Hotel)

C.

C.

Record.

November, 1887.

113

MEMORANDA.

CHARGE.

Page in
Ledger.

Nov 4

November, 1887.

Dail

NAME.

| Su | M | Tu | W | Th | F | Sa |
|----|----|----|----|----|----|----|
| 20 | 21 | 22 | 23 | 24 | 25 | 26 |

20 21 22 23 24 25 26

MEMO

the family

; /

Record.

November, 1887.

115

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Su | M. | Tu. | W | Th. | F. | S |
|----|----|-----|----|-----|----|---|
| 27 | 28 | 29 | 30 | 1 | 2 | 3 |

MEMO

Mr. [unclear] [unclear]

Mr. [unclear] [unclear]

Capt. S. [unclear] [unclear]

Mr. [unclear] [unclear]

[unclear]

MEMORANDA.

| CHARGE.

Page in
Ledger

NAME.

| Su. | M. | Tu. | W | Th. | F. | Sa. |
|-----|----|-----|---|-----|----|-----|
| 4 | 5 | 6 | 7 | 8 | 9 | 10 |

MEM.

Record.

December, 1887.

149

MEMORANDA.

CHARGE.

Paid in
Ledger

NAME.

| Su. | M | Tu. | W. | Th. | F. | Sa |
|-----|----|-----|----|-----|----|----|
| 11 | 12 | 13 | 14 | 15 | 16 | 17 |

MEMO

Record.

December, 1887.

151

MEMORANDA.

CHARGE.

Page in
Ledger.

December, 1887.

Daily

NAME.

| Su. | M. | Tu. | W. | Th. | F. | Sa | MEMO |
|-----|----|-----|----|-----|----|----|------|
| 18 | 19 | 20 | 21 | 22 | 23 | 24 | |

Record.

December, 1887.

153

MEMORANDA.

CHARGE.

Page in
Ledger.

NAME.

| Sa. | M. | Tu. | W. | Th. | F. | Sa. |
|-----|----|-----|----|-----|----|-----|
| 25 | 26 | 27 | 28 | 29 | 30 | 31 |

MEMO

MEMORANDA.

CHARGE.

Page in
Ledger.

Pathological Society, 1886-87

- Oct 26 Hematoma in the brain
Nov 11 2 Cases of Pulmonary Valves
incompleteness papillary muscles
inversion of Thoracic Aorta
Dec 4 Slices of Liver 3 cases

Section of Stomach (Farley)
Section of Pylorus (Smith)
Section of Colon (Hay)
Section of Pulmonary Valve
Section of parietal peritoneum.

Jan 13th. Accutons in sub. capsule. Abscess. degenerated

Jan 27th. Tumor of brain
Hyp & Dilat of the heart

Feb 10th
1. Contraction of liver. Hemorrhage
2. Intestine of sub. adhesion

~~March 10th~~

March 10th

1. Perforated appendix
2. Intestine
3. Adenoma with choleliths

Section of the intestine

DATE.

PULSE.

TEMP.

RESP.

URINE.

STOOLS.

TR

TREATMENT.—CONDITION.—REMARKS.

June 9 Enrol not certified

Subcom 12 1/2 1911

old record. Transcribing from old record

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

1911 1911 1911 1911 1911 1911 1911 1911 1911 1911

TREATMENT.—CONDITION.—REMARKS.

DATE.

PULSE.

TEMP.

RESP.

URINE.

STOOLS.

TREATMENT. — CONDITION. — REMARKS.

DATE.

PULSE.

TEMP.

RESP.

URINE.

STOOLS.

TREATMENT.—CONDITION.—REMARKS.

TREATMENT.—CONDITION.—REMARKS.

DATE.

PULSE.

TEMP.

RESP.

URINE.

STOOLS.

TREATMENT.—CONDITION.—REMARKS.

| DATE. | PULSE. | TEMP. | RESP. | URINE. | STOOLS. |
|-------|--------|-------|-------|--------|---------|
|-------|--------|-------|-------|--------|---------|

TREATMENT.—CONDITION.—REMARKS.

| DATE. | PULSE. | TEMP. | RESP. | URINE. | STOOLS. |
|-------|--------|-------|-------|--------|---------|
|-------|--------|-------|-------|--------|---------|

TREATMENT.—CONDITION.—REMARKS.

TREATMENT.—CONDITION.—REMARKS.

DATE.

PULSE.

TEMP.

RESP.

URINE.

STOOLS.

TREATMENT.—CONDITION.—REMARKS

DATE,

PULSE.

TEMP.

RESP.

URINE.

STOOLS.

Record.

177

TREATMENT.—CONDITION.—REMARKS.

| DATE. | PULSE. | TEMP. | RESP. | URINE. | STOOLS. |
|-------|--------|-------|-------|--------|---------|
|-------|--------|-------|-------|--------|---------|

TREATMENT.—CONDITION.—REMARKS.

DATE.

PULSE.

TEMP.

RESP.

URINE.

STOOLS.

TREATMENT. CONDITION. — REMARKS.

DATE.

PULSE.

TEMP.

RESP.

URINE.

STOOLS.

TREATMENT. CONDITION.—REMARKS.

DATE.

TIME.

PATIENT'S NAME AND ADDRESS.

ATTENDING PHYSICIAN.

REMARKS.

DATE.

TIME.

PATIENT'S NAME AND ADDRESS.

PHYSICIAN.

ATTENDING PHYSICIAN.

REMARKS.

DATE.

TIME.

PATIENT'S NAME AND ADDRESS.

ATTENDING PHYSICIAN.

REMARKS.

PATIENT'S NAME.

End of Last
Menstruat'n.Date Exp'd
ConfinementDate of
Delivery

PATIENT'S NAME.

PATIENT'S NAME.

| End of Last
Menstruat'n. | Date Exp'd
Confinement. | Date of
Delivery. |
|-----------------------------|----------------------------|----------------------|
|-----------------------------|----------------------------|----------------------|

PATIENT'S NAME.

End of Last
Menstruat'n.Date Exp'd
ConfinementDate of
Delivery.

PATIENT'S N

Obstetric Engagements.

193

PATIENT'S NAME.

End of Last
Menstruat'n.Date Exp'd
ConfinementDate of
Delivery.

Obstetric Engagements.

PATIENT'S NAME.

End of Last Date Exp'd
Menstruat'n. ConfinementDate of PATIENT'S NA
Delivery

Obstetric Engagements.

195

PATIENT'S NAME.

| End of Last
Menstruat'n. | Date Exp'd
Confinement | Date of
Delivery. |
|-----------------------------|---------------------------|----------------------|
|-----------------------------|---------------------------|----------------------|

DATE.

NAME OF CHILD.

SEX.

No. OF
CHILD.

COLOR.

MOTHER'S NAME.

PARENTS' NAMES, OCCUPATION AND ADDRESS.--REMARKS.

DATE.

NAME OF CHILD.

SEX.

No. OF
CHILD.

COLOR.

CHILD'S N.

PARENTS' NAMES, OCCUPATION AND ADDRESS.—REMARKS.

DATE.

NAME OF CHILD.

SEX.

No. OF
CHILD.

COLOR.

PARENTS' NAMES, OCCUPATION AND ADDRESS.—REMARKS

| DATE. | NAME OF CHILD. | SEX. | No. OF
CHILD. | COLOR. | REMARKS. |
|-------|----------------|------|------------------|--------|----------|
|-------|----------------|------|------------------|--------|----------|

PARENTS' NAMES, OCCUPATION AND ADDRESS.—REMARKS.

DATE.

NAME OF CHILD.

SEX.

NO. OF
CHILD.

COLOR.

WEIGHT.

PARENTS' NAMES, OCCUPATION AND ADDRESS.—REMARKS.

DATE.

NAME OF CHILD.

SEX.

NO. OF
CHILD.

COLOR.

EYES.

PARENTS' NAMES, OCCUPATION AND ADDRESS.—REMARKS.

Vaccinations.

NAME.

DATE.

RESULT.

Vaccinations.

209

| NAME. | DATE. | RESULT. |
|-------|-------|---------|
|-------|-------|---------|

NAME.

DATE.

RESULT.

Vaccinations.

211

Page

NAME.

DATE.

RESULT.

NAME AND ADDRESS OF DECEASED.

AGE.

Y.

M.

D.

COLOR.

single,
wid, wed.

CAUSE OF DEATH.

DATE.

NAME AND ADDRESS OF DECEASED.

AGE.

Y. M. D.

COLOR.

Mr's, Single,
or Widowed.

CAUSE OF DEATH.

DATE.

ADDRESSES OF PATIENTS.

15.

W. H. H. H.

2011

ADDRESSES OF PATIENTS.

ADDRESSES OF PATIENTS.

ADDRESSES OF PATIENTS.

NURSES' ADDRESSES.

NURSES' ADDRESSES.

| | | | |
|-----------|----------|----------|------------------------|
| 850 | <i>m</i> | | |
| 107 9 | \$ | 5 0 0 | |
| 12 | | 1 00 00 | |
| 117 Jan 5 | | 1 00 00 | - 10 to May 15: roomed |
| March 3 | | 2 00 00 | |
| May 18 | | 5 00 00 | |
| | \$ | 10 00 00 | |

Cash Account.

| BILL
SENT. | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|---------------|------------------------------|--------------------|-----------------|---------------|
| | From October 1st. com ATC | | \$768 | 00 |
| Jan 2nd | E. F. Horn | | 20 | 00 |
| " | John McCord | | 60 | 00 |
| " | J B Lippincott & Co Ed. art | | 3 | 75 |
| 7 | R. T. Sharpley | | 10 | 00 |
| " | Am. news. Oct 5. " 1st AS | | 10 | 00 |
| 10 | S. Hume | | 15 | 00 |
| 3 | Ulege end grade | | 333 | 33 |
| 1-1 | C. Hill D. G. B. McC. Muller | | 10 | 00 |
| 25 | Mrs. Mellen | | 10 | 00 |
| 25 | Mad. News. Bk. 31 at 5- | | 10 | 00 |
| 31 | Mrs. Corby. (Carl Hole) | | 5 | 00 |
| Feb 20 | W. S. Fox | | 100 | 00 |
| " | " " | | 101 | 25 |
| 10 | Ben C. S. W. Wm | | 10 | 00 |
| " | H. Adams Kunkel | | 25 | 00 |
| " | " " " " " " " " | | 10 | 00 |
| 16 | Philips P. L. Le | | 10 | 00 |
| " | P. Sh. Wm | | 10 | 00 |
| " | C. Russell | | 10 | 00 |
| " | H. | | 10 | 00 |
| 23 | Leri | | 10 | 00 |
| 22 | News. 95 Monday 23rd 5- | | 10 | 00 |
| March | Lippincott - Ed. art | | 3 | 80 |
| " | Sid. | | 333 | 33 |
| " | " " " " " " " " | | 10 | 00 |
| " | Glo. Mac Donald Toronto | | 20 | 00 |
| " 10 | C. H. Muller | | 25 | 00 |
| " 12 | J. Conn. & Kelley | | 10 | 00 |
| 3/4 | Mrs. Sykes ... | | 10 | 00 |
| | | | \$2023 | 46 |

Cash Account.

223

| BILL
NO. | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|-------------|-------|--------------------|-----------------|---------------|
| 1 | ... | | 22 | ... |
| 2 | ... | | 10 | ... |
| 3 | ... | | 10 | ... |
| 4 | ... | | 10 | ... |
| 5 | ... | | 10 | ... |
| 6 | ... | | 10 | ... |
| 7 | ... | | 10 | ... |
| 8 | ... | | 10 | ... |
| 9 | ... | | 10 | ... |
| 10 | ... | | 10 | ... |
| 11 | ... | | 10 | ... |
| 12 | ... | | 10 | ... |
| 13 | ... | | 10 | ... |
| 14 | ... | | 10 | ... |
| 15 | ... | | 10 | ... |
| 16 | ... | | 10 | ... |
| 17 | ... | | 10 | ... |
| 18 | ... | | 10 | ... |
| 19 | ... | | 10 | ... |
| 20 | ... | | 10 | ... |
| 21 | ... | | 10 | ... |
| 22 | ... | | 10 | ... |
| 23 | ... | | 10 | ... |
| 24 | ... | | 10 | ... |
| 25 | ... | | 10 | ... |
| 26 | ... | | 10 | ... |
| 27 | ... | | 10 | ... |
| 28 | ... | | 10 | ... |
| 29 | ... | | 10 | ... |
| 30 | ... | | 10 | ... |
| 31 | ... | | 10 | ... |
| 32 | ... | | 10 | ... |
| 33 | ... | | 10 | ... |
| 34 | ... | | 10 | ... |
| 35 | ... | | 10 | ... |
| 36 | ... | | 10 | ... |
| 37 | ... | | 10 | ... |
| 38 | ... | | 10 | ... |
| 39 | ... | | 10 | ... |
| 40 | ... | | 10 | ... |
| 41 | ... | | 10 | ... |
| 42 | ... | | 10 | ... |
| 43 | ... | | 10 | ... |
| 44 | ... | | 10 | ... |
| 45 | ... | | 10 | ... |
| 46 | ... | | 10 | ... |
| 47 | ... | | 10 | ... |
| 48 | ... | | 10 | ... |
| 49 | ... | | 10 | ... |
| 50 | ... | | 10 | ... |
| 51 | ... | | 10 | ... |
| 52 | ... | | 10 | ... |
| 53 | ... | | 10 | ... |
| 54 | ... | | 10 | ... |
| 55 | ... | | 10 | ... |
| 56 | ... | | 10 | ... |
| 57 | ... | | 10 | ... |
| 58 | ... | | 10 | ... |
| 59 | ... | | 10 | ... |
| 60 | ... | | 10 | ... |
| 61 | ... | | 10 | ... |
| 62 | ... | | 10 | ... |
| 63 | ... | | 10 | ... |
| 64 | ... | | 10 | ... |
| 65 | ... | | 10 | ... |
| 66 | ... | | 10 | ... |
| 67 | ... | | 10 | ... |
| 68 | ... | | 10 | ... |
| 69 | ... | | 10 | ... |
| 70 | ... | | 10 | ... |
| 71 | ... | | 10 | ... |
| 72 | ... | | 10 | ... |
| 73 | ... | | 10 | ... |
| 74 | ... | | 10 | ... |
| 75 | ... | | 10 | ... |
| 76 | ... | | 10 | ... |
| 77 | ... | | 10 | ... |
| 78 | ... | | 10 | ... |
| 79 | ... | | 10 | ... |
| 80 | ... | | 10 | ... |
| 81 | ... | | 10 | ... |
| 82 | ... | | 10 | ... |
| 83 | ... | | 10 | ... |
| 84 | ... | | 10 | ... |
| 85 | ... | | 10 | ... |
| 86 | ... | | 10 | ... |
| 87 | ... | | 10 | ... |
| 88 | ... | | 10 | ... |
| 89 | ... | | 10 | ... |
| 90 | ... | | 10 | ... |
| 91 | ... | | 10 | ... |
| 92 | ... | | 10 | ... |
| 93 | ... | | 10 | ... |
| 94 | ... | | 10 | ... |
| 95 | ... | | 10 | ... |
| 96 | ... | | 10 | ... |
| 97 | ... | | 10 | ... |
| 98 | ... | | 10 | ... |
| 99 | ... | | 10 | ... |
| 100 | ... | | 10 | ... |

| BILL
SENT. | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|---------------|---------------|--------------------|-----------------|---------------|
| 1 | Is S. L... | | 50 | |
| 2 | ... | | 5 | |
| 4 | ... | | | |
| 6 | ... | | 10 | |
| 8 | ... | | 50 | 00 |
| 10 | ... | | 900 | 00 |
| 12 | ... | | 10 | 00 |
| 14 | ... | | | |
| July 2 | Leas | | 33 | 00 |
| 4 | B J Doyle | | 5 | 00 |
| 12 | C. D. Farrell | | 10 | 00 |
| 13 | Capt Griffin | | 15 | 00 |
| 14 | Con D. Hines | | 10 | 00 |
| 10 | C. D. Farrell | | 15 | 00 |
| 11 | C. D. Farrell | | 15 | 00 |
| 12 | C. D. Farrell | | 10 | 00 |
| 13 | C. D. Farrell | | 10 | 00 |
| 14 | C. D. Farrell | | 10 | 00 |
| 15 | C. D. Farrell | | 10 | 00 |
| 21 | ... | | 10 | 00 |
| 22 | ... | | 10 | 00 |

Cash Account.

225

| DATE | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|------|-------|--------------------|-----------------|---------------|
|------|-------|--------------------|-----------------|---------------|

Cash Account.

227

| | PAYEE. | AMOUNT CHARGED | AMOUNT PAID. | WHEN PAID. |
|----|-----------------------------|----------------|--------------|------------|
| 76 | John Dennis | 10 00 | | |
| | G T R Bell | 2 00 | | |
| | T H Hutchins | 10 00 | | |
| | Philip White | 00 00 | | |
| | D S Stone | 5 00 | | |
| | Ben Thompson | 10 00 | | |
| | John Mc Card | 60 00 | | |
| | C F Muller | 25 00 | | |
| | Ben Mellon | 10 00 | | |
| | Ward Crump | 20 00 | | |
| | Sirley Gilegan 315 Spgs | 10 00 | | |
| | Ken Schlemmer 1425 West Ave | 10 00 | | |
| | Wm Chapley | 10 00 | | |
| | Wm Parker | 10 00 | | |
| | Wm ... | 20 00 | | |
| 7 | A M Rehnig 834 Market | 10 00 | | |
| | Eggs H & Grafton | 15 00 | | |
| | Shank Potters 50 North | 10 00 | | |
| | C. B. Mc Miller 50 South | 10 00 | | |
| | Mrs Ditch Chas Hill | 20 00 | | |
| | Mrs Phillips | 10 00 | | |
| | Mr Camp. Cent. Rd | 5 00 | | |
| | Mr Kerr Chestnut H. C. B | 10 00 | | |
| | J. W. ... | 20 00 | | |
| | Mr Dale | | | |
| 7 | Bushwick St. ... | 10 00 | | |
| | F. A. ... | 10 00 | | |
| | Levin | 10 00 | | |
| | E. M. ... | 10 00 | | |
| | ... | 50 00 | | |

| BILL
SENT. | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|---------------|----------------------------------|--------------------|-----------------|---------------|
| 10/24/11 | Mr. Macdonald | | 20 00 | |
| 5 | Mr. Chapley | | 10 00 | PA |
| 5 | Wm. J. J. 20 21 St. Coll. Ave. | | 10 00 | PA |
| 12 | Cony. S. 100 Kelly | | 50 00 | PA |
| 11 | Wm. J. J. 20 21 St. Coll. Ave. | | 5 00 | |
| 14 | Wm. B. Robins 2211 St. James Ave | | 30 00 | PA |
| 15 | Mr. Capstone 15 15th | | 10 00 | PA |
| 16 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 17 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 18 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 19 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 20 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 21 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 22 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 23 | Canada Life Ins. Co | | 50 00 | PA |
| 24 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 25 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 26 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 27 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 28 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 29 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 30 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 31 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 32 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 33 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 34 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 35 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 36 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 37 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 38 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 39 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 40 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 41 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 42 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 43 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 44 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 45 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 46 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 47 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 48 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 49 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 50 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 51 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 52 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 53 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 54 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 55 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 56 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 57 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 58 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 59 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 60 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 61 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 62 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 63 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 64 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 65 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 66 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 67 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 68 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 69 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 70 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 71 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 72 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 73 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 74 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 75 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 76 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 77 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 78 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 79 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 80 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 81 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 82 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 83 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 84 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 85 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 86 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 87 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 88 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 89 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 90 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 91 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 92 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 93 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 94 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 95 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 96 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 97 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 98 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 99 | Mr. J. J. 333 21 12th | | 10 00 | PA |
| 100 | Mr. J. J. 333 21 12th | | 10 00 | PA |

Cash Account.

229

[illegible]

| BILL
SENT. | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|--------------------|------------------------------------|--------------------|-----------------|---------------|
| From Oct 18 / 1887 | | | | |
| Oct. | | | | |
| - | Mr Mitchell 46 N 19th | | 20 | 00 |
| 1 | John Wolffskill. | | 30 | 00 |
| 2 | John Gromes 2230 Ralegh St. | | 25 | 00 |
| 2 | Geo Jordan Ford. Jacksonville Mo | | 25 | 00 |
| 8 | W. H. H. H. | | 5 | 00 |
| 6 | Mr J. 7th & 1st ave | | 5 | 00 |
| 10 | 1 Paris 15 2nd & 1st ave | | 10 | 00 |
| 10 | D C Wharton Smith Darlington Mo | | 10 | 00 |
| 11 | Barney 12 Tugaw & 1st ave | | 5 | 00 |
| 17 | C. Smith & Son | | 5 | 00 |
| 21 | Mr Bates 55 & N 17th | | 10 | 00 |
| 23 | C. Smith & Son 17th & 1st ave | | 10 | 00 |
| 27 | C. Smith & Son | | 5 | 00 |
| 1 | E E Lewis Solomon City Kansas | | 15 | 00 |
| | W. H. H. H. | | 5 | 00 |
| 1 | Mr Seibert Kellheim | | 10 | 00 |
| 2 | Mr H. H. 2504 Cumberland ave | | 5 | 00 |
| 4 | Mr H. H. | | 10 | 00 |
| 4 | Mr H. H. 15 23 | | 10 | 00 |
| 4 | Mr Carroll Leicestershire 12th St. | | 15 | 00 |
| 9 | Mr J. H. H. 600 N. H. | | 10 | 00 |
| 11 | Geo J Ford Baltimore | | 15 | 00 |
| 16 | Mr H. H. | | 5 | 00 |
| 19 | Mr H. H. 12th & 1st | | 10 | 00 |
| 29 | Mr H. H. York Pa | | 10 | 00 |
| " | Mr H. H. 12th & 1st | | 5 | 00 |
| 30 | Capt Griffin Pollstown | | 10 | 00 |
| " | Mr H. H. Cleveland | | 20 | 00 |

Cash Account.

231

| PAID
BY | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|------------|-------|--------------------|-----------------|---------------|
|------------|-------|--------------------|-----------------|---------------|

Cash Account.

| BILL
SENT. | DATE. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|---------------|----------------------|--------------------|-------------------|---------------|
| | S.B.C. | | 500 00 | |
| | Travel exp. - dinner | | 107 | Nov 11 |
| | Travel exp. - meals | | 60 00 | Oct 13 |
| | Travel exp. - | | 34 00 | Nov 7 |
| | Travel exp. - | | 213.80 | |
| | Overseas | | 9 ^m 8. | |
| | Exp. - | | 52 00 | |
| | Exp. - | | 35 | |
| | Exp. - | | 14 15 | |
| | Exp. - | | 25 00 | |
| | Exp. - | | 10 00 | |
| | Exp. - | | 6 35 | |
| | Exp. - | | 2 00 | |
| | Exp. - | | 227.00 | |
| | Exp. - | | 5 00 | |
| | Exp. - | | 9 91 | Oct 10 |
| | Exp. - | | 34 95 | |
| | Exp. - | | 6 15 | |
| | Exp. - | | 128 35 | Nov 1 |
| | Exp. - | | 11 00 | |
| | Exp. - | | 2 95 | Oct 12 |
| | Exp. - | | 10 00 | |
| | Exp. - | | 24 74 | Oct 13 |
| | Exp. - | | 54 00 | Nov. |
| | Exp. - | | 11 00 | |
| | Exp. - | | 83 00 | |
| | Exp. - | | 70 | |

Cash Account.

BILL
SENT.

NAME.

AMOUNT
CHARGED.AMOUNT
PAID.WHEN
PAID.

Cash Account.

225

| DATE | NAME. | AMOUNT
CHARGED. | AMOUNT
PAID. | WHEN
PAID. |
|------|-------|--------------------|-----------------|---------------|
|------|-------|--------------------|-----------------|---------------|

Cash Account.

BILL
SENT.

NAME.

AMOUNT
CHARGED.AMOUNT
PAID.WHEN
PAID.

GENERAL MEMORANDA.

GENERAL MEMORANDA.

GENERAL MEMORANDA.

GENERAL MEMORANDA.

The Medical News Visiting List.

— (ERASABLE TABLET.) —


Lea Brothers & Co. Philadelphia


George W. Carter


Chicago


James W. Carter

St. Louis














MS

Acc. no. 302

v. 3

